

EFFECTIVE SOLUTION OF THE $D(-1)$ -QUADRUPLE CONJECTURE

ANDREJ DUJELLA, ALAN FILIPIN, AND CLEMENS FUCHS

ABSTRACT. The $D(-1)$ -quadruple conjecture states that there does not exist a set of four positive integers such that the product of any two distinct elements is one greater than a perfect square. It is proved that if $\{a, b, c, d\}$ is such a set, then $\max\{a, b, c, d\} < 10^{10^{23}}$, hence a proof of the conjecture is given up to a completely determined, finite (but currently computationally infeasible large) set of possible exceptions to check.

2000 *Mathematics Subject Classification*: 11D09, 11D45 (primary), 11B37, 11J68 (secondary).

Keywords: Diophantine m -tuples, Pell equation, Diophantine approximation

1. INTRODUCTION

Let n be an integer. A set of m positive integers is called a Diophantine m -tuple with the property $D(n)$ or simply $D(n)$ - m -tuple, if the product of any two of them increased by n is a perfect square.

A lot of work has been done on $D(n)$ - m -tuples in the last decade, but in fact the problem is much older. It was first studied by Diophantus in the case $n = 1$. He found a set of four positive rationals with the above property: $\{\frac{1}{16}, \frac{33}{16}, \frac{17}{4}, \frac{105}{16}\}$. The first $D(1)$ -quadruple however, the set $\{1, 3, 8, 120\}$, was found by Fermat. Later Euler was able to add the fifth positive rational, $\frac{777480}{8288641}$, to the Fermat's set (see [6], [7, pp. 103-104, 232], [8, pp. 141-145] and [24, pp. 177-181]). Recently, Gibbs [22] found examples of sets of six positive rationals with the property of Diophantus. The folklore conjecture is that there does not exist a $D(1)$ -quintuple. In 1969, Baker and Davenport [2] proved that the Fermat's set cannot be extended to a $D(1)$ -quintuple. Recently, the first author proved that there does not exist a $D(1)$ -sextuple and there are only finitely many $D(1)$ -quintuples (see [15], this refined previous results [19, 12]). The last result is effective, namely if $\{a, b, c, d, e\}$ is a $D(1)$ -quintuple, then $d < 10^{2171}$ and $e < 10^{10^{26}}$ (cf. [15, Corollary 4, p. 210]). This proves the $D(1)$ -quintuple conjecture in an effective way, leaving open a completely determined finite set to check.

For general n , it is easy to see, by considering congruences modulo 4, that if $n \equiv 2 \pmod{4}$ then there does not exist a $D(n)$ -quadruple (see [5, 23, 29]), while the first author proved in [9] that if $n \not\equiv 2 \pmod{4}$ and $n \notin S = \{-4, -3, -1, 3, 5, 8, 12, 20\}$, then there exists at least one Diophantine quadruple with the property $D(n)$. The conjecture is that for $n \in S$ there does not exist a Diophantine quadruple with the property $D(n)$.

The best known upper bounds for the size of sets with the property $D(n)$ are logarithmic in $|n|$ (see [14, 16]). Moreover, for n prime it was shown recently that the size is bounded by the absolute constant 2^{170} (cf. [18]).

Therefore, in the case $n = -1$ the conjecture is that there does not exist a $D(-1)$ -quadruple. In analogy to above this is known as the $D(-1)$ -quadruple conjecture and it appeared explicitly in [10] for the first time. It is known that some particular $D(-1)$ -triples cannot be extended to $D(-1)$ -quadruples, namely this was verified for the triples $\{1, 2, 5\}$ (by Brown in [5], see also [32, 25, 31, 26]), $\{1, 5, 10\}$ (by Mohanty and Ramasamy in [28]), $\{1, 2, 145\}$, $\{1, 2, 4901\}$, $\{1, 5, 65\}$, $\{1, 5, 20737\}$, $\{1, 10, 17\}$, $\{1, 26, 37\}$ (by Kedlaya [25]) and $\{17, 26, 85\}$ (again by Brown in [5]). Moreover, Brown proved that the infinite families $\{x^2 + 1, (x + 1)^2 + 1, (2x + 1)^2 + 4\}$, if $x \not\equiv 0 \pmod{4}$, $\{2, 2x^2 + 2x + 1, 2x^2 + 6x + 5\}$, if $x \equiv 1 \pmod{4}$ of $D(-1)$ -triples cannot be extended to quadruples. The first author proved the conjecture in [11] for all triples of the form $\{1, 2, c\}$.

Very recently, Dujella and Fuchs [17] proved that there does not exist a $D(-1)$ -quintuple. More precisely, they proved that there does not exist a $D(-1)$ -quadruple $\{a, b, c, d\}$ with $2 \leq a < b < c < d$. This implies that we are left by considering the case $a = 1$, i.e. $D(-1)$ -quadruples of the form $\{1, b, c, d\}$. In [17] it was remarked that the case $a = 1$ seems more involved and much harder and that it can be compared with the strong version of the quintuple conjecture for $n = 1$, which says that every $D(1)$ -triple can be extended to a $D(1)$ -quadruple in an essentially unique way.

In the meantime the non-extendibility of $\{1, b, c\}$ was confirmed for $b = 5$ (in [1]), for $b = 10$ (by the second author [20]), and for $b = 17, 26, 37, 50$ (by Fujita in [21]).

The aim of the present paper is to go further and to prove that in fact there are at most finitely many $D(-1)$ -quadruples $\{1, b, c, d\}$. We will prove the following result:

Theorem 1. *Let $\mathcal{D} = \{1, b, c\}$ be a $D(-1)$ -triple. Then:*

- (i) *If $c > b^9$, then \mathcal{D} cannot be extended to a $D(-1)$ -quadruple $\{a, b, c, d\}$ such that $d > c$.*
- (ii) *If $11b^6 \leq c \leq b^9$, then \mathcal{D} cannot be extended to a $D(-1)$ -quadruple.*

Assume that $\{1, b, c, d\}$ with $1 < b < c < d$ is an extension of \mathcal{D} to a $D(-1)$ -quadruple.

- (iii) If $b^3 < c < 11b^6$, then $c < 10^{238}$, $d < 10^{10^{23}}$,
- (iv) if $b^{1.1} < c \leq b^3$, then $c < 10^{491}$, $d < 10^{10^{23}}$,
- (v) if $3b < c \leq b^{1.1}$, then $c < 10^{94}$, $d < 10^{10^{21}}$,
- (vi) if $c = 1 + b + 2\sqrt{b-1}$, then $c < 10^{74}$, $d < 10^{10^{21}}$.

Note that, by [17, Lemma 7], if $c > b$ and $c \neq 1 + b + 2\sqrt{b-1}$ then it follows that $c > 3b$. Hence, the theorem covers all possible $D(-1)$ -triples $\{1, b, c\}$ such that $1 < b < c$.

To sum up, we get the following effective solution of the $D(-1)$ -quadruple conjecture:

Theorem 2. *There are only finitely many $D(-1)$ -quadruples. Moreover, if $\{a, b, c, d\}$ is a $D(-1)$ -quadruple, then $\max\{a, b, c, d\} < 10^{10^{23}}$.*

As in the case for the $D(1)$ -quintuple conjecture, this proves the $D(-1)$ -quadruple conjecture in an effective way. Unfortunately, the remaining set is too large to be checked by the means of a computer program at the moment.

This is also the first nontrivial result (for integers $\not\equiv 2 \pmod{4}$) related to the following conjecture

Conjecture 1. *If n is not a perfect square, then there exist only finitely many $D(n)$ -quadruples.*

Since, by [9, Remark 3], all elements of a $D(-4)$ -quadruple are even, Theorem 2 implies that Conjecture 1 is valid for $n = -1$ and $n = -4$.

The case $n = -1$ deserves special attention because this case is closely connected with another old problem investigated by Diophantus and Euler. Namely, Diophantus studied the problem of finding numbers such that the product of any two increased by the sum of these two gives a square. He found two triples $\{4, 9, 28\}$ and $\{\frac{3}{10}, \frac{21}{5}, \frac{7}{10}\}$ satisfying this property. Euler found a quadruple $\{\frac{5}{2}, \frac{9}{56}, \frac{9}{224}, \frac{65}{224}\}$ and asked if there is an integer solution of this problem (see [6], [7, pp. 85-86, 215-217], [8, pp. 101-104] and [24, pp. 162-164, 344-347]). In [13] an infinite family of rational quintuples with the same property was given. Since

$$(1) \quad xy + x + y = (x+1)(y+1) - 1,$$

we see that the problem of finding integer m -tuples with the property that for any two distinct elements the product plus their sum is a perfect square is equivalent to finding $D(-1)$ - m -tuples. In fact the main result in [17] completely solved this problem, namely it was shown that there does not exist a set of four positive integers such that the product of any two

distinct elements plus their sum is a perfect square.

By the equivalence (1), we get the following corollaries to the above theorems, which extend the problem solved in [17].

Corollary 1. *There are at most finitely many sets $\{a^2, b^2, c^2\}$, $0 < a < b < c$ of three positive perfect squares such that the product of any two of its distinct elements plus their sum is a perfect square.*

Corollary 2. *There are at most finitely many sets of four integers with the property that the product of any two of its distinct elements plus their sum is a perfect square.*

In the next section we will start by collecting useful information about this problem (especially from [17]). The strategy of proof follows the same lines as the proofs of almost all other recent results on non-extendability of $D(n)$ - m -tuples. First we reduce the problem of finding d which extends $\{1, b, c\}$ to a $D(-1)$ -quadruple to a system of simultaneous Pellian equations, which leads to the consideration of intersections of linear recurring sequences. We know that the indices of these sequences have the same parity. Moreover, we have a very important congruence relation. E.g. this relation implies that the sequences cannot have intersections for small indices. Moreover, by the gap principle, which was the main improvement in the proof in [17], we get precise information on the initial terms of the recurring sequences.

If $c \geq 11b^6$, in Section 3 we obtain (Proposition 1) by Bennett's theorem on simultaneous approximations of square roots which are close to 1 (in a slightly refined version for our context by Fujita [21]) parts (i) and (ii) (the case of *very large solutions*) of our main theorem (Theorem 1).

In Section 4 we will compute some general upper bound for the indices of the recurring sequences in terms of c by using Baker's theory of linear forms in logarithms of algebraic numbers (in fact we will use Matveev's result [27]).

In Section 5 we will give the proof of part (iii)-(v) (*large, medium size, and small solutions*, respectively) of the theorem (see Propositions 2, 3, 4), where we have to get lower bounds for n in terms of some power of c . For *medium size* (part (iv) of the theorem, i.e. $b^{1.1} < c < b^4$) and *small* (part (v) of the theorem, i.e. $3b < c < b^{1.1}$) *solutions* we first have to refine the congruence relations. In fact this is the most important new part of the proof.

Finally, we will give the proof of the case of *very small solutions* ($c \leq 3b$), namely part (vi) of the theorem, in Section 6 (Proposition 5).

2. PRELIMINARIES

Let $\{1, b, c\}$, where $1 < b < c$, be a $D(-1)$ -triple and let r, s, t be positive integers defined by

$$b - 1 = r^2, \quad c - 1 = s^2, \quad bc - 1 = t^2.$$

In this paper, the symbols r, s, t will always have this meaning. Assume that there exists a positive integer $d > c$ such that $\{1, b, c, d\}$ is a $D(-1)$ -quadruple. We have

$$(2) \quad d - 1 = x^2, \quad bd - 1 = y^2, \quad cd - 1 = z^2,$$

with integers x, y, z . Eliminating d from (2) we obtain the following system of Pellian equations

$$(3) \quad z^2 - cx^2 = c - 1,$$

$$(4) \quad bz^2 - cy^2 = c - b.$$

By combining these two equations we additionally have $y^2 - bx^2 = b - 1$. We will describe the sets of solutions of equations (3) and (4) in the following lemma.

Lemma 1. *If (z, x) and (z, y) , with positive integers x, y, z , are solutions of (3) and (4) respectively, then there exist integers z_0, x_0 and z_1, y_1 with*

- (i) (z_0, x_0) and (z_1, y_1) are solutions of (3) and (4) respectively,
- (ii) the following inequalities are satisfied:

$$(5) \quad |x_0| < s, \quad 0 < z_0 < c,$$

$$(6) \quad |y_1| < t, \quad 0 < z_1 < c,$$

and there exist integers $m, n \geq 0$ such that

$$(7) \quad z + x\sqrt{c} = (z_0 + x_0\sqrt{c})(s + \sqrt{c})^{2m},$$

$$(8) \quad z\sqrt{b} + y\sqrt{c} = (z_1\sqrt{b} + y_1\sqrt{c})(t + \sqrt{bc})^{2n}.$$

Proof. This is [17, Lemma 1]. □

From (7) we conclude that $z = v_m$, for some (z_0, x_0) with the above properties and integer $m \geq 0$, where

$$(9) \quad v_0 = z_0, \quad v_1 = (2c - 1)z_0 + 2sx_0, \quad v_{m+2} = (4c - 2)v_{m+1} - v_m.$$

Hence for varying $m \geq 0$ the solutions z form a binary recurrent sequence $(v_m)_{m \geq 0}$ whose initial terms are found by solving equation (7) for z when $m = 0$ and 1, and whose characteristic equation has the roots $(s + \sqrt{c})^2$ and

$(s - \sqrt{c})^2$. In the same manner, from (8), we conclude that $z = w_n$ for some (z_1, y_1) with the above properties and integer $n \geq 0$, where

$$(10) \quad w_0 = z_1, \quad w_1 = (2bc - 1)z_1 + 2tcy_1, \quad w_{m+2} = (4bc - 2)w_{n+1} - w_n.$$

Our system of equations (3) and (4) is thus transformed to finitely many equations of the form $z = v_m = w_n$.

From (9) and (10) we get by induction

$$\begin{aligned} v_m &\equiv (-1)^m z_0 \pmod{2c}, \\ w_n &\equiv (-1)^n z_1 \pmod{2c}, \\ v_m &\equiv (-1)^m (z_0 - 2cm^2 z_0 - 2csmx_0) \pmod{8c^2}, \\ w_n &\equiv (-1)^n (z_1 - 2bcn^2 z_1 - 2ctny_1) \pmod{8c^2}. \end{aligned}$$

(see [17, Lemma 2]). So if the equation $v_m = w_n$ has a solution, then we have $z_0 = z_1$. Moreover, we have the following properties:

Lemma 2. *If $v_m = w_n, n \neq 0$, then*

- (i) $m \equiv n \pmod{2}$,
- (ii) $n \leq m \leq 2n$,
- (iii) $m^2 z_0 + smx_0 \equiv bn^2 z_1 + tny_1 \pmod{4c}$.

Proof. Part (i) and (iii) follow at once from $z_0 = z_1$ and the above congruence relations. Statement (ii) is [17, Lemma 5]. Observe that it was remarked in [17, Section 3] that this statement holds for arbitrary $D(-1)$ -quadruples and not only in the special situation considered there. \square

Moreover, we have the following bounds for v_m, w_n , respectively.

Lemma 3. *We have*

$$\begin{aligned} (c - 1)(4c - 3)^{m-1} &< v_m < 4c^2(4c - 2)^{m-1}, \quad \text{for } m \geq 1, \\ (c - b)(4bc - 3)^{n-1} &< w_n < 4bc^2(4bc - 2)^{n-1}, \quad \text{for } n \geq 1. \end{aligned}$$

Proof. This is part of the proof of [17, Lemma 5] (cf. (3.1) in [17] and the equation just before that). \square

We collect some useful gap principles for the elements of $D(-1)$ -triples and quadruples.

Lemma 4. *Let $\{a, b, c\}$ be a $D(-1)$ -triple and $0 < a < b < c$, then $c = a + b + 2r$ or $c > 3ab \geq 3b$.*

Let $\{1, b, c, d\}$ be a $D(-1)$ -quadruple and $1 < b < c < d$, then we have $d > c^9$.

Proof. The first part of the Lemma is [17, Lemma 7]. The second part is a consequence of [17, Lemma 13]. Observe that in fact there it is proved for the $D(-1)$ -quadruple $\{1, a, b, c\}$ with $1 < a < b < c$. The assumption that $\{a, b, c, d\}$ with $2 \leq a < b < c < d$ and d minimal under all such quadruples was only needed to ensure that $\{1, a, b, c\}$ is really a $D(-1)$ -quadruple again (by the use of [17, Lemma 3]). \square

Lemma 4 has the following important implication on the fundamental solutions of (9) and (10) described in Lemma 1.

Lemma 5. *Let the integers z_0, z_1, x_0, y_1 be as in Lemma 1. If $c \leq b^9$, then*

$$z_0 = z_1 = s, x_0 = 0, y_1 = \pm\sqrt{b-1} = \pm r.$$

Proof. Let us define d_0 by

$$d_0 = \frac{z_0^2 + 1}{c}.$$

It follows that

$$d_0 - 1 = x_0^2, bd_0 - 1 = y_1^2, cd_0 - 1 = z_0^2$$

and $d_0 < c$ (this construction was the key lemma in [17], in fact it was [17, Lemma 3]). If $d_0 > 1$, then $\{1, b, d_0, c\}$ is again a $D(-1)$ -quadruple and by the second part of Lemma 4 it follows that $c > b^9$, a contradiction. Hence, $d_0 = 1$, which implies our assertion (cf. also [17, Lemma 4] for this final conclusion). \square

It is easy to show that the equation $v_m = w_n$ cannot have solutions with small indices.

Lemma 6. *We have $v_1 \neq w_1, v_2 \neq w_2, v_4 \neq w_2$.*

Proof. See [17, Lemma 8, Lemma 10 and Lemma 11]. Observe again that the conclusions of these lemmata are independent of the existence of a minimal extension, which was assumed there. \square

We have collected all information we need to prove our main theorem, which will be done in the following sections, respectively.

3. VERY LARGE SOLUTIONS

The proof will follow from a lower bound for m, n in terms of some power of c , together with an logarithmic upper bound obtained from Bennett's theorem. The idea for obtaining the lower bound is that if $v_m = w_n$, then we can study the congruence from Lemma 2(iii). We will show that if b, m, n are small compared with c , these congruences are equations, which are in contradiction to the equation $v_m = w_n$.

Lemma 7. *If $v_m = w_n$, $n \neq 0, 1$ and $c \geq 11b^6$, then $n > c^{\frac{1}{6}}$.*

Proof. From Lemma 2(iii) we have $m^2z_0 + smx_0 \equiv bn^2z_0 + tny_1 \pmod{4c}$. Assume now that $n \leq c^{\frac{1}{6}}$. It is easy to see that $\max\{|m^2z_0|, |smx_0|, |tny_1|\} < |bn^2z_0|$. We first consider the case $c > b^9$. Let d_0 be such that $cd_0 - 1 = z_0^2$. If $d_0 > 1$ then $\{1, b, d_0, c\}$ is a $D(-1)$ -quadruple and we get by the second part of Lemma 4 that $c > (\max\{b, d_0\})^9 > b^6d_0^3$. Then we have

$$|bn^2z_0| < b \cdot c^{\frac{1}{3}} \cdot (cd_0)^{\frac{1}{2}} < c^{\frac{1}{3}} \cdot c^{\frac{1}{2}} \cdot (b^6d_0^3)^{\frac{1}{6}} < c.$$

If $d_0 = 1$ then we get as in the proof of Lemma 5 that $z_0 = s$. Thus we immediately get

$$|bn^2z_0| < c^{\frac{1}{9}} \cdot c^{\frac{1}{3}} \cdot c^{\frac{1}{2}} < c.$$

Now we consider the case $11b^6 \leq c \leq b^9$. Here we can use Lemma 5 to get

$$|bn^2z_0| < c^{\frac{1}{6}} \cdot c^{\frac{1}{3}} \cdot c^{\frac{1}{2}} = c.$$

Therefore we can conclude that $m^2z_0 + smx_0 = bn^2z_0 + tny_1$. We have

$$m^2z_0 + smx_0 < m^2\sqrt{cd_0} + m\sqrt{cd_0} < m(m+1)\sqrt{\frac{c}{b}}\sqrt{|y_1|^2 + 1}$$

and

$$bn^2z_0 - tny_1 > \sqrt{bc}|y_1|n^2 - tn|y_1| > \sqrt{bc}|y_1|n^2 - \sqrt{bc}|y_1|n = \sqrt{bc}|y_1|n(n-1).$$

This implies

$$m(m+1)\sqrt{|y_1|^2 + 1} > b|y_1|n(n-1),$$

and so, by the relation $m \leq 2n$ and $b \geq 65$, we get

$$5 \geq \frac{2n+1}{n-1} > \frac{65|y_1|}{2\sqrt{|y_1|^2 + 1}} > \frac{65}{2\sqrt{2}}.$$

We therefore get a contradiction that proves the statement. \square

Next we need an upper bound for n in terms of c . To obtain it we use the following slightly modified version of a special case of Bennett's theorem [4] (it is also a modified version of Rickert's theorem [30]), which is Theorem 3.5 in [21].

Lemma 8. *Let b and N be integers with $b \geq 5$ and $N \geq 2.39b^7$. Then the numbers*

$$\theta_1 = \sqrt{1 + \frac{1-b}{N}} \text{ and } \theta_2 = \sqrt{1 + \frac{1}{N}}$$

satisfy

$$\max \left\{ \left| \theta_1 - \frac{p_1}{q} \right|, \left| \theta_2 - \frac{p_2}{q} \right| \right\} > \left[32.1 \frac{b^2(b-1)^2}{2b-1} N \right]^{-1} q^{-1-\lambda}$$

for all integers p_1, p_2, q with $q > 0$, where

$$\lambda = \frac{\log \frac{16.1b^2(b-1)^2N}{2b-1}}{\log \frac{3.37N^2}{b^2(b-1)^2}} < 1.$$

We apply this theorem with the numbers $N = t^2 = bc - 1 \geq 11b^7 - 1 > 2.39b^7$, $p_1 = bsx$, p_2bz , $q = ty$, i.e. to

$$\theta_1 = \frac{s\sqrt{b}}{t}, \quad \theta_2 = \frac{\sqrt{bc}}{t}.$$

First we show that the solutions of our problem induce good approximations to the roots θ_1, θ_2 .

Lemma 9. *All positive integers solutions x, y, z of (3) and (4) satisfy*

$$\max \left\{ \left| \theta_1 - \frac{bsx}{ty} \right|, \left| \theta_2 - \frac{bz}{ty} \right| \right\} < \frac{b-1}{y^2}.$$

Proof. Since $f(x) = b(x-1)/(bx-1)$ is an increasing function we get

$$\left| \theta_1 - \frac{bsx}{ty} \right| = \frac{s\sqrt{b}}{t} \left| 1 - \frac{x\sqrt{b}}{y} \right| = \frac{s\sqrt{b}}{t} \left| 1 - \frac{bx^2}{y^2} \right| \left| 1 + \frac{x\sqrt{b}}{y} \right|^{-1} < \frac{b-1}{y^2}.$$

Moreover, since $bz/y > \sqrt{bc}$ we have

$$\begin{aligned} \left| \theta_2 - \frac{bz}{ty} \right| &= \frac{1}{t} \left| \sqrt{bc} - \frac{bz}{y} \right| = \frac{b}{t} \left| c - \frac{bz^2}{y^2} \right| \left| \sqrt{bc} + \frac{bz}{y} \right|^{-1} \\ &< \frac{bc-b}{t} \frac{1}{y^2} \frac{1}{2\sqrt{bc}} < \frac{1}{2y^2} \frac{bc-1}{\sqrt{bc}(bc-1)} < \frac{b-1}{y^2}. \end{aligned}$$

This proves the lemma. \square

Now we are ready to prove the first two parts of Theorem 1. Namely, we show the following proposition.

Proposition 1. *Let $\{1, b, c\}$ with $1 < b < c$ be a $D(-1)$ -triple. If $c \geq 11b^6$, then there does not exist an integer $d > c$ such that $\{1, b, c, d\}$ is a $D(-1)$ -quadruple. If $11b^6 \leq c \leq b^9$, then the triple $\{1, b, c\}$ cannot be extended to a $D(-1)$ -quadruple.*

Proof. Combining the upper bound of Lemma 9 with the lower bound from Lemma 8 we get

$$\left[32.1 \frac{b^2(b-1)^2}{2b-1} t^2 \right]^{-1} (ty)^{-1-\lambda} < \frac{b-1}{y^2},$$

which implies

$$y^{1-\lambda} < 32.1 \frac{b^2(b-1)^3}{2b-1} t^{3+\lambda} < \frac{32.1b^2(b-1)^3(bc-1)^2}{2b-1}.$$

We have

$$\frac{1}{1-\lambda} = \frac{\log \frac{3.37(bc-1)^2}{b^2(b-1)^2}}{\log \frac{3.37(2b-1)(bc-1)}{16.1b^4(b-1)^4}},$$

which leads to

$$\log y < \frac{\log \frac{3.37(bc-1)^2}{b^2(b-1)^2} \log \frac{32.1b^2(b-1)^3(bc-1)^2}{2b-1}}{\log \frac{3.37(2b-1)(bc-1)}{16.1b^4(b-1)^4}}.$$

We have (observe that $c \geq 11b^6$ and $b \geq 65, c \geq 82$)

$$\begin{aligned} \log \frac{3.37(bc-1)^2}{b^2(b-1)^2} &< \log \frac{3.37b^2c^2}{b^2(b-1)^2} < 2 \log \frac{1.9c}{b-1} < 2 \log 0.03c, \\ \log \frac{32.1b^2(b-1)^3(bc-1)^2}{2b-1} &< \log (16.18b^6c^2) < 3 \log 1.14c, \\ \log \frac{3.36(2b-1)(bc-1)}{16.1b^4(b-1)^4} &> \log \frac{2.29(2b^2c - bc - 2b + 1)}{b^2c} > 1.51. \end{aligned}$$

Consequently,

$$\log y < 3.98 \log 0.03c \log 1.14c.$$

Moreover, from (4) and the first part of Lemma 3, we get

$$cy^2 = bz^2 - c + b > b(c-1)^2(4c-3)^{2m-2} - c > bc(4c-3)^{2m-2} - c > c(4c-3)^{2m-2}$$

and therefore $y > (4c-3)^{m-1}$.

Assume that $n \neq 0, 1$. Then by Lemma 2 and Lemma 7 we have $m \geq n > c^{\frac{1}{6}}$, which implies

$$(11) \quad \left(c^{\frac{1}{6}} - 1\right) \log(4c-3) < 3.98 \log 0.03c \log 1.14c.$$

If we define the function

$$f(c) = 3.98 \log 0.03c \log 1.14c - \left(c^{\frac{1}{6}} - 1\right) \log(4c-3),$$

then it is easy to check that this function is decreasing for $c > 0.55 \cdot 10^{12}$. Since $c \geq 11b^6 \geq 11 \cdot 65^6 > 0.82 \cdot 10^{12}$, inequality (11) cannot be fulfilled. Hence, we conclude that $n = 0$ or $n = 1$. Now, by Lemma 6, we have that $z = v_0 = w_0$. But then $d = (z^2 + 1)/c < c$. Moreover, if $11b^6 \leq c \leq b^9$, then Lemma 5 implies that $z = s$ and $d = 1$. This proves the proposition. \square

4. A DIOPHANTINE APPROXIMATION RESULT

All other parts of Theorem 1 will be obtained by applying the theory of linear form of logarithms of algebraic numbers instead of Bennett's approximation result. In this way it will not be possible to exclude any solution, but we will get effective upper bounds. In this section we will prepare the approximation tool that will be used in the proofs in case of solutions that are not very large.

In fact we will use a result by Matveev [27], which we quote in a suitable simplified version.

Lemma 10. *Let Λ be a linear form in logarithm of l multiplicatively independent totally real algebraic numbers $\alpha_1, \dots, \alpha_l$ with rational integer coefficients b_1, \dots, b_l ($b_l \neq 0$). Let $h(\alpha_j)$ denote the absolute logarithmic height of α_j , $1 \leq j \leq l$. Define the number $D, A_j, 1 \leq j \leq l$, and B by $D = [\mathbb{Q}(\alpha_1, \dots, \alpha_l) : \mathbb{Q}]$, $A_j = \max\{Dh(\alpha_j), |\log \alpha_j|\}$,*

$$B = \max \left\{ 1, \max \left\{ \frac{|b_j| A_j}{A_l} : 1 \leq j \leq l \right\} \right\}.$$

Then

$$(12) \quad \log \Lambda > -C(l)C_0W_0D^2\Omega,$$

where

$$C(l) = \frac{8}{(l-1)!}(l+2)(2l+3)(4e(l+1))^{l+1}, \quad W_0 = \log(1.5eBD \log(eD)),$$

$$C_0 = \log(e^{4.4l+7}l^{5.5}D^2 \log(eD)), \quad \Omega = A_1 \cdots A_l.$$

Proof. See [27, Theorem 2.1]. □

Solving the recurrences (3) and (4) we get

$$v_m = \frac{s}{2} [(s + \sqrt{c})^{2m} + (s - \sqrt{c})^{2m}],$$

$$w_n = \frac{s\sqrt{b} \pm r\sqrt{c}}{2\sqrt{b}} [(t + \sqrt{bc})^{2n} + (t - \sqrt{bc})^{2n}].$$

We now turn the equation $v_m = w_n$ into an inequality for a linear form in three logarithms, to which Lemma 10 will be applied.

Lemma 11. *If $v_m = w_n, n \neq 0$, then*

$$0 < 2n \log(t + \sqrt{bc}) - 2m \log(s + \sqrt{c}) + \log \frac{s\sqrt{b} \pm r\sqrt{c}}{2\sqrt{b}} < (3.96bc)^{-n+1}.$$

Proof. Let

$$P = s(s + \sqrt{c})^{2m}, \quad Q = \frac{s\sqrt{b} \pm r\sqrt{c}}{\sqrt{b}}(t + \sqrt{bc})^{2n}.$$

Then $v_m = w_n$ implies $P + (c-1)P^{-1} = Q + \frac{c-b}{b}Q^{-1}$. Furthermore, we have

$$\begin{aligned} P &> 4s(c-1) = 4\sqrt{c-1}(c-1) > c-1, \\ Q &> \frac{s\sqrt{b} - r\sqrt{c}}{\sqrt{b}}(2bc - 1 + 2\sqrt{bc}) \geq 2c \geq 2. \end{aligned}$$

We get

$$\begin{aligned} P - Q &= \frac{c-b}{b}Q^{-1} - (c-1)P^{-1} < (c-1)Q^{-1} - (c-1)P^{-1} \\ &= (c-1)(P-Q)P^{-1}Q^{-1}, \end{aligned}$$

and therefore we have $Q > P$. Moreover, $P > Q - (c-1)P^{-1} > Q - 1$, which implies that $\frac{Q-P}{Q} < Q^{-1} \leq \frac{1}{2}$. Hence,

$$0 < \log \frac{Q}{P} = -\log \frac{P}{Q} = -\log \left(1 - \frac{Q-P}{Q} \right) < \frac{Q-P}{Q} + \left(\frac{Q-P}{Q} \right)^2.$$

From this we conclude

$$\begin{aligned} 0 &< \log \frac{Q}{P} < \frac{1}{Q} + \frac{1}{Q^2} < \frac{2}{Q} = \frac{2\sqrt{b}}{s\sqrt{b} \pm \sqrt{cr}}(t + \sqrt{bc})^{-2n} \\ &< \frac{2\sqrt{b}}{s\sqrt{b} - r\sqrt{c}}(3.96bc)^{-n} = \frac{2\sqrt{b}(s\sqrt{b} + r\sqrt{c})}{c-b}(3.96bc)^{-n} \\ &< \frac{4b\sqrt{c}}{c-b}(3.96bc)^{-n} < (3.96bc)^{-n+1}. \end{aligned}$$

This proves the lemma. \square

We use this result to prove a principle to obtain an upper bound for n in all remaining cases.

Lemma 12. *If $v_m = w_n$, then*

$$\frac{n-1}{\log \left(\frac{160.36n \log(2.02bc)}{\log c} \right)} < 0.33 \cdot 10^{12} \cdot \log c \cdot \log(bc).$$

Proof. We have $l = 3$, $D = 4$ and therefore $C(3) \leq 0.65 \cdot 10^9$ and $C_0 \leq 29.89$. Moreover, we have $\alpha_1 = s + \sqrt{c}$, $\alpha_2 = t + \sqrt{bc}$, and

$$\alpha_3 = \frac{s\sqrt{b} \pm r\sqrt{c}}{s\sqrt{b}}.$$

The numbers $\alpha_1, \alpha_2, \alpha_3$ satisfy the following algebraic relations: $\alpha_1^2 + (4c - 2)\alpha_1 + 1 = 0$, $\alpha_2^2 + (4bc - 2)\alpha_2 + 1 = 0$, and $b(c - 1)\alpha_3^2 - 2b(c - 1)\alpha_3 + c - b = 0$. Therefore, $A_1 = 2 \log(s + \sqrt{c}) < 2 \log(2\sqrt{c}) < 1.08 \log c$, $A_2 = 2 \log(t + \sqrt{bc}) < \log(\sqrt{2bc}) < 1.05 \log(bc)$, and

$$A_3 \leq 2 \log bc \frac{s\sqrt{b} + r\sqrt{c}}{s\sqrt{b}} < 2 \log(2.02bc) < 2 \log(3.96bc).$$

We also have

$$A_3 \geq \log b(c - 1) \geq \log(64.2c) > 1.95 \log c.$$

Thus,

$$B \leq \frac{4m \log(2.02bc)}{1.95 \log c} \leq \frac{2.06m \log(2.02bc)}{\log c}.$$

By Lemma 11 and Lemma 10, we therefore get

$$n - 1 < 0.33 \cdot 10^{12} \cdot \log \left(\frac{80.18m \log(2.02bc)}{\log c} \right) \log c \log(bc).$$

Finally, we have

$$\frac{n - 1}{\log \left(\frac{160.36n \log(2.02bc)}{\log c} \right)} < 0.33 \cdot 10^{12} \cdot \log c \cdot \log(bc),$$

which is the statement. \square

Our remaining goal is to calculate a lower bound for n in terms of small powers of c as above. Together with our gap conditions between b and c , Lemma 12 will enable us to get effective upper bounds for c in every case. This will be done in the next sections.

5. LARGE, MEDIUM AND SMALL SOLUTIONS

We now start with the remaining cases in Theorem 1. Observe that we always have $c \leq 11b^6$ and therefore by Lemma 5 we have $z_0 = z_1 = s, x_0 = 0, y_1 = \pm r$.

We start with the case of large solutions, i.e. with $b^3 \leq c < 11b^6$. We have the following lower bound for n .

Lemma 13. *If $v_m = w_n, n \neq 0, 1$ and $b^3 \leq c < 11b^6$, then $n > c^{\frac{1}{12}}$.*

Proof. We again start with the relation in Lemma 2(iii), which reads here as $m^2s \equiv bn^2s \pm tnr \pmod{4c}$. Assume that $n \leq c^{\frac{1}{12}}$. Since $\max\{m^2s, tnr\} < bn^2s$, and

$$bn^2s < c^{\frac{1}{3}} \cdot c^{\frac{1}{6}} \cdot c^{\frac{1}{2}} = c,$$

we have equality in the congruence above, thus $m^2s = bn^2s \pm tnr$. This implies $4n^2s > n(bsn - tr)$ and therefore

$$\frac{b\sqrt{c}}{2}n < (b-4)sn < tr < b\sqrt{c},$$

which gives a contradiction. \square

Combining the previous lemma with the result from Baker's theory from Lemma 12, we get part (iii) of Theorem 1.

Proposition 2. *Let $\{1, b, c, d\}$ with $1 < b < c < d$ be a $D(-1)$ -quadruple. If $b^3 \leq c \leq 11b^6$, then $c < 0.22 \cdot 10^{238}$ and $d < 10^{10^{23}}$.*

Proof. Since $b^3 \leq c$, we get $\log(2.02bc) < 1.50 \log c$, and therefore it follows by Lemma 12 and 13 that

$$\frac{n-1}{\log(240.54n)} < 0.29 \cdot 10^{13} \cdot \frac{4}{3}(\log c)^2 < 0.56 \cdot 10^{15}(\log n)^2.$$

Hence, $n < 0.60 \cdot 10^{20}$ and therefore $c < 0.22 \cdot 10^{238}$.

From Lemma 3 we get $z = v_m < 4c^2(4c-2)^{m-1} \leq 4^m c^{m+1}$. Consequently,

$$d \leq \frac{z^2 + 1}{c} \leq \frac{4^{2m} c^{2m+2}}{c} = 16^m c^{2m+1}.$$

Thus, $\log_{10} d \leq m \log_{10} 16 + (2m+1) \log_{10} c \leq 0.58 \cdot 10^{23}$ and therefore $d \leq 10^{10^{23}}$. \square

Next we handle the case of medium size solutions, i.e. with $b^{1.1} \leq c < b^3$, and therefore prove part (iv) of Theorem 1.

Lemma 14. *If $v_m = w_n, n \neq 0, 1, b^{1.1} \leq c < b^3$, and $c > 10^{100}$, then $n \geq c^{0.04}$.*

Proof. We will consider the congruence of Lemma 2(iii). We have

$$(13) \quad s(m^2 - bn^2) \equiv \pm rtn \pmod{4c}.$$

Let us define integers A and B by

$$\begin{aligned} A &= 2bc - 2rst - c, \\ B &= 2bc + 2rst - c, \end{aligned}$$

then $AB = c^2 + 4b^2c - 4b - 4c + 4$. Clearly, $B < 4bc$ and $A > \frac{4b^2c}{4bc} = b$. On the other hand, $A < \frac{c^2 + 4b^2c}{2bc} < c$. This implies that $2rst > 2bc - 2c$, $B > 4bc - 3c$ and

$$A < \frac{c^2 + 4b^2c}{4bc - 3c} < \frac{c}{3b} + \frac{4}{3}b < 0.334c^{0.67} + 1.334c^{0.91} < c^{0.92},$$

which shows that A is small compared with c . Observe that here we use that $c > 10^{100}$.

Multiplying both sides of (13) by $2s$, we get

$$2(m^2 - bn^2) \equiv \mp An \pmod{c}.$$

Assume now that $n \neq 0$ and $n < c^{0.04}$. Then $An < c$ and $|2(m^2 - bn^2)| \leq 2bn^2 < 2c^{0.91+0.08} < c$. Hence, we have an equality:

$$(14) \quad 2bn^2 - 2m^2 = An.$$

The relation (14) and the inequalities $n \leq m \leq 2n$ (cf. Lemma 2(ii)) imply

$$(15) \quad 2(b-4)n \leq A \leq 2(b-1)n.$$

Let us define β by

$$c = 4b^2(2n-1) + \beta.$$

We claim that $|\beta| < c^{0.56}$.

Let us first estimate the quantity γ defined by $\gamma = 4bA - c - 4b^2$. From $AB = c^2 + 4b^2c - 4b - 4c + 4$ and $4bc - 3c < B < 4bc$, we obtain

$$\begin{aligned} \gamma &> \frac{1}{4bc}(-16b^2 - 16bc + 16b) > -8, \\ \gamma &< \frac{3c + 12b^2}{3b} = \frac{c}{b} + 4b. \end{aligned}$$

Therefore, $|\gamma| < c^{0.92}$. Furthermore, from (15) we have $4bA = 8b^2n + \delta$, where $|\delta| \leq 32bn < c^{0.966}$. Since $\beta = \delta - \gamma$, we get by putting these estimates together that

$$|\beta| < c^{0.966} + c^{0.92} < 2c^{0.966} < c^{0.97}.$$

Hence,

$$(16) \quad c = 4b^2(2n-1) + \beta, \quad |\beta| < c^{0.97}.$$

Analyzing (16) we will improve this estimate for $|\beta|$. Namely, (16) clearly implies $4b^2 < c < 16nb^2$. It follows that $|\gamma| < \frac{c}{b} + 4b < 20bn < 10n\sqrt{c} < c^{0.55}$, and also $|\delta| < 32bn < c^{0.553}$. Hence, $|\beta| < c^{0.56}$, as we claimed.

Now we are ready to finish our argument. By squaring (14), we get

$$4(bn^2 - m^2)^2 \equiv 4(b-1)n^2 \pmod{c},$$

which implies

$$(17) \quad -n^4\beta \equiv (2n-1)(4(b-1)n^2 - 4m^4 + 8bm^2n^2) \pmod{c}.$$

But the absolute values of both sides of (17) are less than $c^{0.72}$. Hence, we have the equality in (17), and this implies (assuming that $n \neq 0$)

$$n^2c = (2n-1)(A^2 - 4(b-1)).$$

Since $A^2 \equiv 4(b-1) \pmod{c}$, we find that $(2n-1)|n^2$, which is possible only if $n = 0$ or $n = 1$. We obtain a contradiction, and hence we have

proved that $n \geq c^{0.04}$ □

Combining this lower bound with the upper bound which follows from Lemma 12, we will again obtain that c is bounded by an absolute constant.

Proposition 3. *Let $\{1, b, c, d\}$ with $1 < b < c < d$ be a $D(-1)$ -quadruple. If $b^{1.1} \leq c < b^3$, then $c < 0.39 \cdot 10^{491}$ and $d < 10^{10^{23}}$.*

Proof. Since $b \leq c^{0.91}$, we get $\log(2.02bc) \leq 1.93 \log c$ and $\log(bc) \leq 1.91 \log c$. Now, by Lemma 14 and Lemma 12 we get

$$\frac{n-1}{\log(309.50n)} < 0.40 \cdot 10^{15} (\log n)^2.$$

Hence, $n < 0.42 \cdot 10^{20}$ and thus $c < 0.39 \cdot 10^{491}$.

As in the proof of Proposition 2, we get by Lemma 3 that $\log_{10} d \leq 2n \log_{10} 16 + (4n+1) \log_{10} c \leq 0.83 \cdot 10^{23}$ and therefore $d \leq 10^{10^{23}}$. □

Finally, we turn to part (v) of Theorem 1. This is the case of small solutions, i.e. with $3b < c < b^{1.1}$.

To get the lower bounds for m, n in terms of some small power of c in the case of small solutions, we will use the following very useful construction, which in fact is the essence of the gap principle in Lemma 4.

Lemma 15. *Let $\{a, b, c\}$ be a $D(-1)$ -triple. Define*

$$e = -(a + b + c) + 2abc - 2rst.$$

Then there exist integers u, v, w such that

$$ae + 1 = u^2, be + 1 = v^2, ce + 1 = w^2$$

and

$$c = a + b - e + 2(abe + ruv).$$

Proof. This lemma is a special case of Lemma 3 in [14]. □

Using this lemma we now prove:

Lemma 16. *If $v_m = w_n, n \neq 0, 1, 2$ and $3b < c < b^{1.1}$, then $n \geq 0.25 \cdot c^{0.2}$.*

Proof. With the same notation as in Lemma 14, again we have the congruence

$$(18) \quad 2(m^2 - bn^2) \equiv \mp An \pmod{c}.$$

In this case we get

$$A < \frac{c^2 + 4b^2c}{4bc - 3c} = b + \frac{c + 3b}{4b - 3} < b + \frac{2c}{3b}.$$

Hence, $A = b + \alpha$, where $0 < \alpha < c^{0.1}$.

By Lemma 15, there exist integers e, u, v, w such that

$$e + 1 = u^2, \quad be + 1 = v^2, \quad ce + 1 = w^2,$$

and

$$b = 1 + c - e + 2ce - 2suw.$$

Hence, $(b + e - 1)^2 \equiv 4(c - 1)(e + 1)(ce + 1) \equiv -4(e + 1) \pmod{c}$, and

$$(19) \quad b^2 + 2b(e - 1) + e^2 + 2e + 5 \equiv 0 \pmod{c}.$$

Moreover, e is small compared with c . Indeed, from $c = 1 + b - e + 2be + 2ruv > 2be$, we find that $e < \frac{c}{2b} < c^{0.1}$.

Now (18) implies

$$(20) \quad 2m^2 \pm \alpha n \equiv b(2n^2 \mp n) \pmod{c}.$$

From (20), we also get

$$(21) \quad (2m^2 \pm \alpha n)^2 \equiv b^2(2n^2 \mp n)^2 \pmod{c}$$

and

$$(22) \quad 2(e - 1)(2n^2 \mp n)(2m^2 \pm \alpha n) \equiv 2(e - 1)b(2n^2 \mp n)^2 \pmod{c}.$$

By summing (21) and (22), and taking into account (19), we obtain

$$(23) \quad (2n^2 \mp n)^2(e^2 + 2e + 5) + (2m^2 \pm \alpha n)^2 + 2(e - 1)(2n^2 \mp n)(2m^2 \pm \alpha n) \equiv 0 \pmod{c}.$$

Assume now that $n < 0.25 \cdot c^{0.2}$.

Since $n \geq 3$, $e \geq 3$, $m \leq 2n$ and $c \geq 82$, we have the following estimates:

$$\begin{aligned} |(2n^2 \mp n)^2(e^2 + 2e + 5)| &\leq 6n^4 \cdot 3e^2 \cdot \frac{18c^{0.8}}{256} \cdot \frac{c^2}{4b^2} < 0.02c, \\ (2m^2 \pm \alpha n)^2 &\leq 8m^4 + \alpha^2 n^2 \leq \frac{c^{0.8}}{2} + \frac{c^{0.6}}{8} < 0.23c, \end{aligned}$$

and

$$|2(e - 1)(2n^2 \mp n)(2m^2 \pm \alpha n)| \leq 6en^2(2m^2 + \alpha n) \leq \frac{3c^{0.5}c^{0.4}}{8} < 0.13c.$$

Therefore, we have the equality in (23), i.e.

$$(24) \quad (e^2 + 2e + 5)X^2 + 2(e - 1)(e - 1)XY + Y^2 = 0,$$

where $X = (2n^2 \mp n)$, $Y = 2m^2 \pm \alpha n$. Since the discriminant of (24) is negative (it is equal to $-16u^2$), we conclude that by multiplying (24) with $4(e^2 + 2e + 5)$ we get

$$(2(e^2 + 2e + 5)X + 2(e - 1)^2Y)^2 + 16u^2Y^2 = 0.$$

This implies $X = Y = 0$, which in turn gives $n = 0$, a contradiction. Hence, we proved that $n \geq 0.25 \cdot c^{0.2}$. \square

Again by using Lemma 12 we will obtain that c is bounded by an absolute constant.

Proposition 4. *Let $\{1, b, c, d\}$ with $1 < b < c < d$ be a $D(-1)$ -quadruple. If $3b < c < b^{1.1}$, then $c < 0.78 \cdot 10^{94}$ and $d < 10^{10^{21}}$.*

Proof. Since $b < \frac{c}{3}$, we have $\log(2.02bc) \leq 2\log(0.83c) < 2\log c$ and $\log(bc) \leq 2\log \frac{c}{9}$. The inequality $c \leq 1024n^5$ (cf. Lemma 16) and Lemma 12 imply that

$$\frac{n-1}{\log(320.72n)} < 0.66 \cdot 10^{12} \cdot \log(1024n^5) \log(113.78n^5).$$

Hence, $n \leq 0.15 \cdot 10^{19}$ and $c \leq 0.78 \cdot 10^{94}$.

By Lemma 3 we get $\log_{10} d \leq 2n \log_{10} 16 + (4n+1) \log_{10} c \leq 0.57 \cdot 10^{21}$ and therefore $d \leq 10^{10^{21}}$. \square

It remains to prove the smallest possible case, i.e. when we have $c < 3b$. This will be done in the following section.

6. VERY SMALL SOLUTIONS

Now we consider the case $c = 1 + b + 2\sqrt{b-1}$ of very small solution (part (vi) in Theorem 1). Observe that we are left to consider $c \leq 3b$. By the gap principle (Lemma 4), this implies that we have indeed $c = 1 + b + 2r$. We can write everything parametrized in r . To avoid confusion, we set $k = r$ in this section. Then we have to consider the $D(-1)$ -triple

$$\{1, k^2 + 1, (k+1)^2 + 1\}.$$

Moreover, we have

$$s = k+1, t = k^2 + k + 1, z_0 = z_1 = k+1, x_0 = 0, y_1 = \pm k,$$

and $k \geq 8$.

If $v_m = w_n, n \neq 0, 1$ then the congruence from Lemma 2(iii) has the form

$$(25) \quad \begin{aligned} m^2(k+1) &\equiv n^2(k^2+1)(k+1) \pm n(k^2+k+1)k \\ &\equiv n^2(k+3) \mp n(k+1) \pmod{(k^2+2k+2)}. \end{aligned}$$

We start by proving an upper bound for k in terms of n .

Lemma 17. *In this case, if $v_m = w_n, n \neq 0, 1$, then $k < 4n^2$.*

Proof. Assume that $n^2 \leq \frac{1}{4}k$. We have the following estimates

$$m^2(k+1) \leq 4n^2(k+1) < k(k+1) < k^2 + 2k + 2,$$

$$|n^2(k+3) \pm n(k+1)| < \frac{1}{4}k(k+3) + \frac{1}{2}\sqrt{k}(k+1) < k^2 + 2k + 2.$$

Therefore we have an equality in (25), which gives

$$m^2(k+1) = n^2(k+3) \mp n(k+1).$$

This implies $m^2 \equiv 3n^2 \mp n \pmod{k}$. Since $m^2 \leq 4n^2 < k$ and $|3n^2 \mp n| < \frac{3}{4}k + \frac{1}{2}\sqrt{k} < k$, we have again an equality. This implies $m^2 = 3n^2 \mp n$. Moreover, we now also have the equation $m^2 = n^2 \mp n$. Combining these two facts, we get $2n^2 = \pm n$ and therefore $n = 0$, a contradiction. This proves the lemma. \square

We conclude the proof by combining Lemma 17 with Lemma 12 as in all cases above, to get an effective upper bound for k and therefore for c and d .

Proposition 5. *Let $\{1, k^2 + 1, (k+1)^2 + 1, d\}$ be a $D(-1)$ -quadruple. Then $k < 0.14 \cdot 10^{38}$ and therefore $c = (k+1)^2 + 1 < 0.19 \cdot 10^{75}$ and $d < 10^{10^{21}}$.*

Proof. We have $\log(2.02bc) = \log(2.02(k^2 + 1)(k^2 + 2k + 2)) \leq 4.47 \log k$ and $\log(k^2 + 2k + 2) > \log k$. By combining Lemma 17 and Lemma 12 we get

$$\frac{n-1}{\log(716.81n)} < 0.33 \cdot 10^{12} \log((4n^2 + 1)^2 + 1) \\ \cdot \log((4n^2)^2 + 1)((4n^2 + 1)^2 + 1).$$

This implies $n < 0.90 \cdot 10^{18}$ and therefore $k < 0.33 \cdot 10^{37}$.

Moreover, we get $c = (k+1)^2 + 1 < 0.11 \cdot 10^{74}$ and by Lemma 3 we have $\log_{10} d \leq 2n \log_{10} 16 + (4n+1) \log_{10} c \leq 0.27 \cdot 10^{21}$ and finally this gives $d < 10^{10^{21}}$. \square

Altogether, by combining Propositions 1, 2, 3, 4 and 5 above, we get Theorem 1. Theorem 2, as well as Corollary 1 and 2 are immediate consequences.

7. ACKNOWLEDGEMENTS

This paper was partly written in the frame of a joint Austrian-Croatian project granted by the Croatian Ministry of Science, Education and Sports and the Austrian Exchange Service (Nr. 20/2004); the authors would like to thank them for this support. Moreover, the first and the second author were supported by the Ministry of Science, Education and Sports, Republic of Croatia, grants 0037110 and 037-0372781-2821, the second author

also by the Austrian Science Foundation FWF, grant NFN S9611, and the third author was supported by the Austrian Science Foundation FWF, grant J2407-N12.

REFERENCES

- [1] F. S. Abu Muriefah and A. Al-Rashed, *On the extendibility of the Diophantine triples $\{1, 5, c\}$* , Int. J. Math. Math. Sci. **33** (2004), 1737-1746.
- [2] A. Baker and H. Davenport, *The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford Ser. (2) **20** (1969), 129-137.
- [3] A. Baker and G. Wüstholz, *Logarithmic forms and group varieties*, J. Reine Angew. Math. **442** (1993), 19-62.
- [4] M. A. Bennett, *On the number of solutions of simultaneous Pell equations*, J. Reine Angew. Math. **498** (1998), 173-199.
- [5] E. Brown, *Sets in which $xy + k$ is always a square*, Math. Comp. **45** (1985), 613-620.
- [6] L. E. Dickson, *History of the Theory of Numbers* Vol. 2, Chelsea, New York, 1966, 513-520.
- [7] Diophantus of Alexandria, *Arithmetics and the Book of Polygonal Numbers*, I. G. Bashmakova (ed.), Nauka, Moscow, 1974 (in Russian).
- [8] Diophantus von Alexandria, *Die Arithmetik und die Schrift ber Polygonalzahlen*, G. Wertheim (ed.), Teubner, Leipzig, 1890.
- [9] A. Dujella, *Generalization of a problem of Diophantus*, Acta Arith. **65** (1993), 15-27.
- [10] A. Dujella, *On the exceptional set in the problem of Diophantus and Davenport*, in: Application of Fibonacci Numbers Vol. 7, G. E. Bergum, A. N. Philippou, A. F. Horadam (eds.), Kluwer, Dordrecht, 1998, 69-76.
- [11] A. Dujella, *Complete solution of a family of simultaneous Pellian equations*, Acta Math. Inform. Univ. Ostraviensis **6** (1998), 59-67.
- [12] A. Dujella, *An absolute upper bound for the size of Diophantine m -tuples*, J. Number Theory **89** (2001), 126-150.
- [13] A. Dujella, *An extension of an old problem of Diophantus and Euler II*, Fibonacci Quart. **40** (2002), 118-123.
- [14] A. Dujella, *On the size of Diophantine m -tuples*, Math. Proc. Cambridge Philos. Soc. **132** (2002), 23-33.
- [15] A. Dujella, *There are only finitely many Diophantine quintuples*, J. Reine Angew. Math. **566** (2004), 183-214.
- [16] A. Dujella, *Bounds for the size of sets with the property $D(n)$* , Glas. Mat. Ser. III **39** (2004), 199-205.
- [17] A. Dujella and C. Fuchs, *Complete solution of a problem of Diophantus and Euler*, J. London Math. Soc. (2) **71** (2005), 33-52.
- [18] A. Dujella and F. Luca, *Diophantine m -tuples for primes*, Intern. Math. Research Notices **47** (2005), 2913-2940.
- [19] A. Dujella and A. Pethő, *Generalization of a theorem of Baker and Davenport*, Quart. J. Math. Oxford Ser. (2) **49** (1998), 291-306.
- [20] A. Filipin, *Non-extendibility of $D(-1)$ -triples of the form $\{1, 10, c\}$* , Int. J. Math. Math. Sci. **35** (2005) 2217-2226.
- [21] Y. Fujita, *The extendibility of $D(-1)$ -triples $\{1, b, c\}$* , Publ. Math. Debrecen **70** (2007), 103-117.

- [22] P. Gibbs, *Some rational Diophantine sextuples*, Glas. Mat. Ser. III **41** (2006), 195-203.
- [23] H. Gupta and K. Singh, *On k -triad sequences*, Int. J. Math. Math. Sci. **5** (1985), 799-804.
- [24] T. L. Heath, *Diophantus of Alexandria: A study in the history of Greek Algebra. With a supplement containing an account of Fermat's theorems and problems connected with Diophantine analysis and some solutions of Diophantine problems by Euler*, Cambridge, England, 1910; Powell's Bookstore, Chicago; Martino Publishing, Mansfield Center, 2003.
- [25] K. S. Kedlaya, *Solving constrained Pell equations*, Math. Comp. **67** (1998), 833-842.
- [26] O. Kihel, *On the extendibility of the set $\{1, 2, 5\}$* , Fibonacci Quart. **38** (2000), 464-466.
- [27] E. M. Matveev, *An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic number. II*, Izv. Ross. Akad. Nauk. Ser. Math. **64** (2000), no. 6, 125-180 (in Russian).
- [28] S. P. Mohanty and A. M. S. Ramasamy, *The simultaneous Diophantine equations $5y^2 - 20 = x^2$ and $2y^2 + 1 = z^2$* , J. Number Theory **18** (1984), 356-359.
- [29] S. P. Mohanty and A. M. S. Ramasamy, *On $P_{r,k}$ sequences*, Fibonacci Quart. **23** (1985), 36-44.
- [30] J. H. Rickert, *Simultaneous rational approximation and related Diophantine equations*, Math. Proc. Cambridge Philos. Soc. **113** (1993), 461-472.
- [31] P. G. Walsh, *On two classes of simultaneous Pell equations with no solutions*, Math. Comp. **68** (1999), 385-388.
- [32] D. X. Zheng, *On the systems of Diophantine equations $y^2 - 2x^2 = 1, z^2 - 5x^2 = 4$ and $y^2 - 5x^2 = 4, z^2 - 10x^2 = 9$* , Sichuan Daxue Xuebao **24** (1987), 25-29.

ANDREJ DUJELLA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB
 BIJENIČKA CESTA 30
 10000 ZAGREB, CROATIA
 E-MAIL: duje@math.hr

ALAN FILIPIN

FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF ZAGREB
 FRA ANDRIJE KAČIĆA-MIOŠIĆA 26
 10000 ZAGREB, CROATIA
 E-MAIL: filipin@grad.hr

Current address: Department for Analysis and Computational Number Theory, TU Graz, Steyrergasse 30, 8010 Graz, Austria

CLEMENS FUCHS

DEPARTMENT OF MATHEMATICS, ETH ZURICH
 RAEMISTRASSE 101
 8092 ZURICH, SWITZERLAND
 E-MAIL: clemens.fuchs@math.ethz.ch