

ON CONVERGENTS FORMED FROM DIOPHANTINE EQUATIONS

CARSTEN ELSNER, TAKAO KOMATSU AND IEKATA SHIOKAWA

University of Applied Sciences, Hannover, Germany, Hirosaki University and
Keio University, Japan

ABSTRACT. We compute upper and lower bounds for the approximation of certain values ξ of hyperbolic and trigonometric functions by rationals x/y such that x, y satisfy Diophantine equations. We show that there are infinitely many coprime integers x, y such that

$$|y\xi - x| \ll \frac{\log \log y}{\log y}$$

and a Diophantine equation holds simultaneously relating x, y and some integer z . Conversely, all positive integers x, y with $y \geq c_0$ solving the Diophantine equation satisfy

$$|y\xi - x| \gg \frac{\log \log y}{\log y}.$$

Moreover, we approximate $\sin(\pi\alpha)$ and $\cos(\pi\alpha)$ by rationals in connection with solutions of a quadratic Diophantine equation when $\tan(\pi\alpha/2)$ is a Liouville number.

1. INTRODUCTION AND STATEMENT OF THE RESULTS IN THE CASE OF HYPERBOLIC FUNCTIONS

Let p_n/q_n denote the n th convergent of the number

$$e = \exp(1) = [2, \overline{1, 2k, 1}]_{k=1}^{\infty}.$$

2000 *Mathematics Subject Classification.* 11D09, 11D25, 11J04, 11J70.

Key words and phrases. Diophantine approximation, Diophantine equations, trigonometric and hyperbolic functions.

The first author was supported in part by the Fund for the Promotion of International Scientific Research B-3 (2007) represented by Takao Komatsu.

The second author was supported in part by the Grant-in-Aid for Scientific Research (C) (No. 18540006), the Japan Society for the Promotion of Science.

Put $P_k = p_{3k+1}$, $Q_k = q_{3k+1}$ ($k = 0, 1, 2, \dots$), $P_0 = 3$, $P_{-1} = 1$, $P_{-2} = 1$, $Q_0 = 1$, $Q_{-1} = 1$, $Q_{-2} = -1$, $P_{-k} = P_{k-3}$, and $Q_{-k} = -Q_{k-3}$ ($k = 3, 4, 5, \dots$). By [5, Theorem 1.1], we know that the following identities hold.

$$(1.1) \quad P_{n+2} = 2(2n+5)P_{n+1} + P_n \quad \text{and} \quad Q_{n+2} = 2(2n+5)Q_{n+1} + Q_n \quad (n \in \mathbb{Z}).$$

A similar result can be proven for the leaping convergents of the number

$$e^{1/s} = \exp\left(\frac{1}{s}\right) = [1, \overline{s(2k-1)-1, 1, 1}]_{k=1}^{\infty} \quad (s \geq 2).$$

Put $P_0 = 1$, $P_1 = p_3 = 2s+1$, $P_k = p_{3k}$ ($k = 2, 3, \dots$), $Q_0 = 1$, $Q_1 = q_3 = 2s-1$, and $Q_k = q_{3k}$ ($k = 2, 3, \dots$). Then by [11, Theorem 1], we have

$$(1.2) \quad P_{n+2} = 2s(2n+3)P_{n+1} + P_n \quad \text{and} \quad Q_{n+2} = 2s(2n+3)Q_{n+1} + Q_n \quad (n \geq 0).$$

The preceding recurrence relations, (1.1) and (1.2), imply $P_n Q_n \equiv 1 \pmod{2}$ for all n and for all $s \geq 1$. Let $h(x)$ be a function with

$$h \in C^{(1)}[1+\delta, 3] \longrightarrow \mathbb{R}, \quad \min_{1+\delta \leq t \leq 3} |h'(t)| > 0,$$

where δ is an arbitrary small positive number. In particular, $h'(x)$ takes its minimum and maximum for $1+\delta \leq x \leq 3$. In our applications we choose $h(x)$ as rational functions such that at rational points p/q the functions h take the form

$$h\left(\frac{p}{q}\right) = \frac{g_1(p, q)}{g_2(p, q)}$$

where $g_1, g_2 \in \mathbb{Z}[p, q]$. Then we had the following.

LEMMA 1.1 ([7, Theorem 3]). *Let $s \geq 1$ be an integer and let P_n, Q_n , and h be as above. Then the inequalities*

$$C_1 \frac{\log \log Q_n}{Q_n^2 \log Q_n} \leq \left| h(e^{1/s}) - h\left(\frac{P_n}{Q_n}\right) \right| \leq C_2 \frac{\log \log Q_n}{Q_n^2 \log Q_n} \quad (n \geq 3)$$

hold, where C_1 and C_2 are effectively computable positive constants depending only on s and the function h .

In [7] the application of this Lemma to various functions h leads to the following approximation results. The basic idea is initiated in [6]. In what follows, all the constants C_3, C_4, \dots, C_{18} appearing in the rest of this section depend only on s .

PROPOSITION 1.2. *Let s be a positive integer and x and $y (\geq 3)$ relatively prime integers with $y \equiv 0 \pmod{2}$ such that $x^2 + y^2$ is a square. Then*

$$\left| y \sinh\left(\frac{1}{s}\right) - x \right| > C_3 \frac{\log \log y}{\log y}.$$

On the other hand, there are infinitely many pairs x, y as just described satisfying

$$\left| y \sinh \left(\frac{1}{s} \right) - x \right| < C_4 \frac{\log \log y}{\log y}.$$

PROPOSITION 1.3. Let $s \geq 1$ be an integer and x and $y (\geq 3)$ relatively prime integers with $y \equiv 0 \pmod{2}$ such that $x^2 - y^2$ is a square. Then

$$\left| y \cosh \left(\frac{1}{s} \right) - x \right| > C_5 \frac{\log \log y}{\log y}.$$

On the other hand, there are infinitely many pairs x, y as just described satisfying

$$\left| y \cosh \left(\frac{1}{s} \right) - x \right| < C_6 \frac{\log \log y}{\log y}.$$

PROPOSITION 1.4. Let $s \geq 1$ be an integer and x and $y (\geq 3)$ relatively prime integers with $x \equiv 1 \pmod{2}$ such that $y^2 - x^2$ is a square. Then

$$\left| y \tanh \left(\frac{1}{s} \right) - x \right| > C_7 \frac{\log \log y}{\log y}.$$

On the other hand, there are infinitely many pairs x, y as just described satisfying

$$\left| y \tanh \left(\frac{1}{s} \right) - x \right| < C_8 \frac{\log \log y}{\log y}.$$

Theorem	Diophantine equation	ξ_s
Theorem 1.5	$x^2 + y^2 = z^4$	$\frac{1}{2} \left(\sinh \left(\frac{1}{s} \right) - \operatorname{cosech} \left(\frac{1}{s} \right) \right)$
Theorem 1.6	$x^2 + y^2 = 2z^2$	$\frac{\sinh \left(\frac{1}{s} \right) - 1}{\sinh \left(\frac{1}{s} \right) + 1}$
Theorem 1.7	$x^3 + 4y^3 = z^2$	$\frac{\sinh \left(\frac{4}{s} \right) + \cosh \left(\frac{4}{s} \right) + 4 \sinh \left(\frac{1}{s} \right) + 4 \cosh \left(\frac{1}{s} \right)}{1 - 2 \sinh \left(\frac{3}{s} \right) - 2 \cosh \left(\frac{3}{s} \right)}$
Theorem 1.8	$x^2 + xy + y^2 = z^2$	$\frac{2 \sinh \left(\frac{1}{s} \right)}{2 + \cosh \left(\frac{1}{s} \right) - \sinh \left(\frac{1}{s} \right)}$
Theorem 1.9	$x^2 + y^2 = u^4 - v^2$	$\frac{1}{2} \sinh \left(\frac{2}{s} \right)$

TABLE 1. Theorems dealing with hyperbolic functions

One goal of this paper is to treat more Diophantine equations and the corresponding hyperbolic functions. A similar paper [1] appeared without giving bounds. Our emphasis here is to give computable bounds, too. We organize this paper as follows: first, we give more examples of values of *hyperbolic functions*, which can be approximated by rationals satisfying Diophantine equations (Theorems 1.5 - 1.9). Then, in the final section 6, we generalize our results to the approximation of values of *trigonometric functions* at specific rational points (Theorem 6.1). Finally, we treat the approximations of $\sin(\pi\alpha)$ and $\cos(\pi\alpha)$ by rationals with numerators and denominators solving the Pythagorean equation $x^2 + y^2 = z^2$, when additionally $\tan(\pi\alpha/2)$ is assumed to be a Liouville number (Theorems 6.4, 6.5). The Table 1 gives an overview on the subsequent theorems dealing with *hyperbolic functions*. For any rational function h let h^{-1} be the inverse function, always defined in an interval centered around some β with $h'(\beta) \neq 0$.

THEOREM 1.5. *Let $s \geq 1$ be an integer and let*

$$\xi_s := \frac{1}{2} \left(\sinh\left(\frac{1}{s}\right) - \operatorname{cosech}\left(\frac{1}{s}\right) \right).$$

Then there are infinitely many triplets (x, y, z) of integers satisfying simultaneously

$$|y\xi_s - x| < C_9 \frac{\sqrt{y} \log \log y}{\log y} \quad \text{and} \quad x^2 + y^2 = z^4.$$

Conversely, for any integer $s \geq 1$ and for given integers $x, y (\geq 3), z$ with $x^2 + y^2 = z^4$, we have the inequality

$$|y\xi_s - x| > C_{10} \frac{\sqrt{y} \log \log y}{\log y}.$$

THEOREM 1.6. *Let $s \geq 1$ be an integer and let*

$$\xi_s := \frac{\sinh(1/s) - 1}{\sinh(1/s) + 1}, \quad h(t) := \frac{t^2 - 2t - 1}{t^2 + 2t - 1}.$$

Then there are infinitely many triplets (x, y, z) of integers satisfying simultaneously

$$|y\xi_s - x| < C_{11} \frac{\log \log y}{\log y} \quad \text{and} \quad x^2 + y^2 = 2z^2.$$

Conversely, for any integer $s \geq 1$ and for given positive integers $x, y (\geq 3), z$ with $y > x$, $h^{-1}(x/y) > \sqrt{2} - 1$, and $x^2 + y^2 = 2z^2$, we have the inequality

$$|y\xi_s - x| > C_{12} \frac{\log \log y}{\log y}.$$

THEOREM 1.7. *Let $s \geq 1$ be an integer and let*

$$\xi_s := \frac{\sinh(4/s) + \cosh(4/s) + 4 \sinh(1/s) + 4 \cosh(1/s)}{1 - 2 \sinh(3/s) - 2 \cosh(3/s)}.$$

Then there are infinitely many triplets (x, y, z) of integers satisfying simultaneously

$$|y\xi_s - x| < C_{13} \frac{\sqrt{|y|} \log \log |y|}{\log |y|} \quad \text{and} \quad x^3 + 4y^3 = z^2.$$

Conversely, for $s \geq 1$ and for given integers x, y, z with $-y \geq 3$ and $x^3 + 4y^3 = z^2$, we assume that

$$(x, y, z) \in \left\{ (p(p^3 + 4q^3), q(q^3 - 2p^3), p^6 - 10p^3q^3 - 2q^6) : p, q \in \mathbb{Z}, \right. \\ \left. q(q^3 - 2p^3) \leq -3 \right\}$$

and that

$$\frac{x}{y} \leq -\frac{(\sqrt[3]{5+3\sqrt{3}})(3+\sqrt{3})}{3+2\sqrt{3}}, \\ \frac{p}{q} > \sqrt[3]{5+3\sqrt{3}} \quad (s=1), \quad \sqrt[3]{1/2} < \frac{p}{q} < \sqrt[3]{5+3\sqrt{3}} \quad (s>1).$$

Then we additionally have the inequality

$$|y\xi_s - x| > C_{14} \frac{\sqrt{|y|} \log \log |y|}{\log |y|}.$$

THEOREM 1.8. *Let $s \geq 1$ be an integer and let*

$$\xi_s := \frac{2 \sinh(1/s)}{2 + \cosh(1/s) - \sinh(1/s)}.$$

Then there are infinitely many triplets (x, y, z) of integers satisfying simultaneously

$$|y\xi_s - x| < C_{15} \frac{\log \log y}{\log y} \quad \text{and} \quad x^2 + xy + y^2 = z^2.$$

Conversely, for any integer $s \geq 1$ and for given positive integers $x, y (\geq 3), z$ with $x^2 + xy + y^2 = z^2$, we have the inequality

$$|y\xi_s - x| > C_{16} \frac{\log \log y}{\log y}.$$

THEOREM 1.9. *Let $s \geq 1$ be an integer and let*

$$\xi_s := \frac{1}{2} \sinh \left(\frac{2}{s} \right).$$

Then there are infinitely many quadruplets (x, y, u, v) of integers satisfying simultaneously

$$|y\xi_s - x| < C_{17} \frac{\sqrt{y} \log \log y}{\log y} \quad \text{and} \quad x^2 + y^2 = u^4 - v^2.$$

Conversely, for any integer $s \geq 1$ and for given positive integers $x, y (\geq 3), u, v$ with $x^2 + y^2 = u^4 - v^2$, we assume that

$$(x, y, u, v) \in \left\{ (p^4 - q^4, 4p^2q^2, p^2 + q^2, 2pq(p^2 - q^2)) : p, q \in \mathbb{Z}^+, p > q \right\}.$$

Then we additionally have the inequality

$$|y\xi_s - x| > C_{18} \frac{\sqrt{y} \log \log y}{\log y}.$$

As can be seen from these results and their proofs, $|q\xi_s - x|$ tends to zero when the parametric representation of the solutions x, y of the corresponding Diophantine equations are given by homogeneous forms of degree two. In Theorems 1.7, 1.9, and 6.1, it is hard to say whether all solutions of the Diophantine equations are given by the above mentioned parameterizations. Therefore, we preferred to deal with stronger conditions for the lower bounds of $|y\xi_s - x|$.

2. AN AUXILIARY LEMMA

First, we mention that for every rational function

$$R(\alpha) = \frac{g_t \alpha^t + g_{t-1} \alpha^{t-1} + \cdots + g_0}{h_u \alpha^u + h_{u-1} \alpha^{u-1} + \cdots + h_0}$$

we have

$$(2.1) \quad R(e^{1/s}) = \frac{\sum_{\nu=1}^t g_\nu \sinh(\nu/s) + \sum_{\nu=0}^t g_\nu \cosh(\nu/s)}{\sum_{\mu=1}^u h_\mu \sinh(\mu/s) + \sum_{\mu=0}^u h_\mu \cosh(\mu/s)},$$

which follows immediately from the identity $e^{1/s} = \sinh(1/s) + \cosh(1/s)$. We shall apply (2.1) with integral coefficients, and t, u not exceeding 4.

LEMMA 2.1. *Let $s \geq 1$ be an integer and let $h(t) \in \overline{\mathbb{Q}}(t) \setminus \mathbb{Q}$. Then there exists a closed interval $I_s = [e^{1/s} - \delta, e^{1/s} + \delta]$ centered around $e^{1/s}$ such that for any positive coprime integers p, q with $q \geq 3$ the following holds.*

$$(2.2) \quad \frac{p}{q} \in I_s \quad \implies \quad |h(e^{1/s}) - h(p/q)| > C \frac{\log \log q}{q^2 \log q},$$

where δ and C are positive constants depending possibly on s and the function h .

PROOF. Since $h(t)$ is a non-constant rational function with algebraic coefficients, we know by Lindemann's theorem that $h(e^{1/s}) \neq 0$ and $h'(e^{1/s}) \neq 0$. Hence there is a closed interval $I_s = [e^{1/s} - \delta, e^{1/s} + \delta]$ with $\delta > 0$ such that

$$(2.3) \quad h(t) \in C^{(1)}(I_s) \quad \text{and} \quad h'(t) \neq 0 \quad (t \in I_s).$$

Then, by the mean-value theorem, we have

$$(2.4) \quad C_{23}|e^{1/s} - t| < |h(e^{1/s}) - h(t)| < C_{24}|e^{1/s} - t| \quad (t \in I_s)$$

with

$$C_{23} := \min_{t \in I_s} |h'(t)| > 0 \quad \text{and} \quad C_{24} := \max_{t \in I_s} |h'(t)| > 0.$$

Let p, q be integers with $q \geq 3$ and $p/q \in I_s$. We can assume without loss of generality that

$$(2.5) \quad |h(e^{1/s}) - h(p/q)| \leq \frac{\log \log q}{q^2 \log q},$$

since otherwise for such rationals p/q the inequality in (2.2) is already satisfied with $C \leq 1$. Thus, by (2.4) and (2.5), there exists a constant C_{25} such that

$$(2.6) \quad \left| e^{1/s} - \frac{p}{q} \right| < \frac{\log \log q}{C_{23} q^2 \log q} < \frac{1}{3q^2} \quad (q \geq C_{25}).$$

Then from the well-known properties of the convergents of simple continued fractions, we find $p/q = p_k/q_k$ for some $k > 0$, where p_k/q_k is the k th convergent of $e^{1/s}$. If the $(k+1)$ th partial quotient a_{k+1} of the continued fraction expansion of $e^{1/s}$ is 1, we have for $q \geq q_n$ that

$$\left| e^{1/s} - \frac{p}{q} \right| = \left| e^{1/s} - \frac{p_k}{q_k} \right| > \frac{1}{(2 + a_{k+1})q_k^2} = \frac{1}{3q_k^2} \geq \frac{1}{3q^2},$$

which contradicts (2.6). Hence, by the definition of P_n and Q_n , i.e.,

$$(2.7) \quad \frac{P_n}{Q_n} = \frac{p_{3n+1}}{q_{3n+1}} \quad (\text{if } s = 1) \quad \text{or} \quad \frac{P_n}{Q_n} = \frac{p_{3n}}{q_{3n}} \quad (\text{if } s \geq 2),$$

we have $p/q = P_n/Q_n$ for some n , and so $p = P_n$ and $q = Q_n$ since $\gcd(p, q) = 1$. Therefore, by the left-hand inequality in Lemma 1.1, we get

$$\left| h(e^{1/s}) - h\left(\frac{p}{q}\right) \right| = \left| h(e^{1/s}) - h\left(\frac{P_n}{Q_n}\right) \right| > C_1 \frac{\log \log Q_n}{Q_n^2 \log Q_n} = C_1 \frac{\log \log q}{q^2 \log q}$$

for $q \geq C_{25}$. Hence, for some $0 < C < \min\{1, C_1\}$, the lemma is proven. \square

REMARK 2.2. Lemma 2.1 also holds without the condition $\gcd(p, q) = 1$, since then we have to deal with $Q_n \leq q$.

3. PARAMETER SOLUTIONS OF DIOPHANTINE EQUATIONS

The lemmata in this section can be proven by straightforward computations. Therefore, the details are left to the reader.

LEMMA 3.1 ([3, p.466], [4, p.256]). *All positive integral solutions of*

$$(3.1) \quad x^2 + y^2 = z^4$$

are given by

$$\begin{aligned} x &= (p^2 - q^2)^2 - (2pq)^2 = p^4 - 6p^2q^2 + q^4, \\ y &= 4pq(p^2 - q^2), \\ z &= p^2 + q^2 \end{aligned}$$

(up to exchange of x and y), where $p, q \in \mathbb{Z}$. Moreover, if we put

$$h(t) := \frac{(t^2 - 1)^2 - 4t^2}{4t(t^2 - 1)},$$

we have $x/y = h(p/q)$ for any solution $x, y (\neq 0)$ of the above equation (3.1). The function $h(t)$ is monotonously increasing for $t > 1$, and $h \in C^{(1)}(1, \infty)$.

LEMMA 3.2 ([2, p. 353, Corollary 6.3.14], [12, p. 13]). *All positive integral solutions of*

$$(3.2) \quad x^2 + y^2 = 2z^2$$

are given by

$$\begin{aligned} x &= p^2 - q^2 - 2pq, \\ y &= p^2 - q^2 + 2pq, \\ z &= p^2 + q^2 \end{aligned}$$

(up to exchange of x and y), where $p, q \in \mathbb{Z}$. Moreover, if we put

$$h(t) := \frac{t^2 - 2t - 1}{t^2 + 2t - 1},$$

we have $x/y = h(p/q)$ for any solution $x, y (\neq 0)$ of the above equation (3.2). The function $h(t)$ is monotonously increasing for $t > \sqrt{2} - 1$, and $h \in C^{(1)}(\sqrt{2} - 1, \infty)$.

LEMMA 3.3 ([9]). *A set of integral solutions of*

$$(3.3) \quad x^3 + 4y^3 = z^2,$$

where x, y, z are relatively prime in pairs, is given by

$$\begin{aligned} x &= p(p^3 + 4q^3), \\ y &= q(q^3 - 2p^3), \\ z &= p^6 - 10p^3q^3 - 2q^6, \end{aligned}$$

where $p, q \in \mathbb{Z}$. Moreover, if we put

$$h(t) := t \frac{4 + t^3}{1 - 2t^3},$$

we have $x/y = h(p/q)$ for any solution $x, y (\neq 0)$ of the above equation (3.3). The function $h(t)$ is monotonously increasing for $0.7937\dots = \sqrt[3]{1/2} < t < \sqrt[3]{5 + 3\sqrt{3}} = 2.1684\dots$, monotonously decreasing for $t > \sqrt[3]{5 + 3\sqrt{3}}$, and $h \in C^{(1)}(\sqrt[3]{1/2}, \infty)$.

LEMMA 3.4 ([4, p. 406]). All positive integral solutions of

$$(3.4) \quad x^2 + xy + y^2 = z^2$$

are given by

$$\begin{aligned} x &= p^2 - q^2, \\ y &= 2pq + q^2, \\ z &= p^2 + pq + q^2 \end{aligned}$$

(up to exchange of x and y), where $p, q \in \mathbb{Z}$ with $p > q$. Moreover, if we put

$$h(t) := \frac{t^2 - 1}{2t + 1},$$

we have $x/y = h(p/q)$ for any solution $x, y (\neq 0)$ of the above equation (3.4). The function $h(t)$ is monotonously increasing for $t > -1/2$, and $h \in C^{(1)}(-1/2, \infty)$.

LEMMA 3.5 ([4, p. 260]). A set of positive integral solutions of

$$(3.5) \quad x^2 + y^2 = u^4 - v^2$$

is given by

$$\begin{aligned} x &= p^4 - q^4, \\ y &= 4p^2q^2, \\ u &= p^2 + q^2, \\ v &= 2pq(p^2 - q^2) \end{aligned}$$

(up to exchange of x and y), where $p, q \in \mathbb{Z}$ with $p > q$. Moreover, if we put

$$h(t) := \frac{t^2}{4} - \frac{1}{4t^2},$$

we have $x/y = h(p/q)$ for any solution $x, y (\neq 0)$ of the above equation (3.5). The function $h(t)$ is monotonously increasing for $t > 0$, and $h \in C^{(1)}(0, \infty)$.

4. PROOF OF THEOREM 1.5

By the function $h(t)$ defined in Lemma 3.1, we have

$$\begin{aligned}\xi_s &:= h(e^{1/s}) = \frac{(e^{2/s} - 1)^2 - 4e^{2/s}}{4e^{1/s}(e^{2/s} - 1)} = \frac{(e^{1/s} - e^{-1/s})^2 - 4}{4(e^{1/s} - e^{-1/s})} \\ &= \frac{\sinh^2(1/s) - 1}{2\sinh(1/s)} = \frac{1}{2}(\sinh(1/s) - \operatorname{cosech}(1/s)).\end{aligned}$$

Let P_n, Q_n ($n \geq 3$) be convergents of $e^{1/s}$ given by (2.7), and let

$$\begin{aligned}x_n &= P_n^4 - 6P_n^2Q_n^2 + Q_n^4, \\ y_n &= 4P_nQ_n(P_n^2 - Q_n^2), \\ z_n &= P_n^2 + Q_n^2.\end{aligned}$$

Then it follows from Lemma 3.1 and $P_nQ_n \equiv 1 \pmod{2}$ that

$$x_n^2 + y_n^2 = z_n^4, \quad x_n > 0, \quad y_n > 0, \quad z_n > 0, \quad 4|x_n, \quad 4|y_n, \quad 2|z_n, \quad \text{and}$$

$$x_n/y_n = h(P_n/Q_n).$$

Applying Lemma 1.1, we have

$$(4.1) \quad \left| \xi_s - \frac{x_n}{y_n} \right| = \left| h(e^{1/s}) - h\left(\frac{P_n}{Q_n}\right) \right| \leq C_2 \frac{\log \log Q_n}{Q_n^2 \log Q_n}.$$

Since $1 < P_n/Q_n < 3$ for all integers $n \geq 1$,

$$(4.2) \quad Q_n^2 < P_nQ_n < P_nQ_n(P_n^2 - Q_n^2) = \frac{y_n}{4} < P_n^3Q_n < 27Q_n^4.$$

Particularly, for $Q_n \geq 3$, we get

$$Q_n^7 > \frac{Q_n^3 y_n}{108} \geq \frac{y_n}{4}, \quad \log Q_n > \frac{\log(y_n/4)}{7},$$

and so

$$Q_n^2 > \frac{\sqrt{y_n}}{6\sqrt{3}} = \frac{\sqrt{y_n/4}}{3\sqrt{3}}, \quad \log \log Q_n < \log \log(y_n/4).$$

Hence, from (4.1) we conclude that

$$\left| \xi_s - \frac{x_n}{y_n} \right| < 21\sqrt{3}C_2 \frac{\log \log(y_n/4)}{\sqrt{y_n/4} \log(y_n/4)} \quad (n \geq 1).$$

Setting

$$C_9 := 21\sqrt{3}C_2, \quad x := x_n/4, \quad y := y_n/4, \quad z := z_n/2,$$

we get the upper bound in Theorem 1.5.

Conversely, we apply Lemma 2.1 to the function h defined in Lemma 3.1. There exists a nontrivial closed interval $I_s \subset (1, \infty)$ centered around $e^{1/s}$ such that for any positive integers $p, q (\geq 3)$, $p/q \in I_s$ the inequality

$$(4.3) \quad \left| h(e^{1/s}) - h(p/q) \right| > C \frac{\log \log q}{q^2 \log q}$$

holds. Let positive integers $x, y (\geq 3), z$ be given such that $x^2 + y^2 = z^4$. Since $h((1, \infty)) = \mathbb{R}$, x/y may take every rational number. By Lemma 3.1 there are integral parameters p, q with

$$\begin{aligned} x &= p^4 - 6p^2q^2 + q^4, \\ y &= 4pq(p^2 - q^2), \\ z &= p^2 + q^2, \end{aligned}$$

and $x/y = h(p/q)$. By $h'(t) > 0$ ($t \in I_s$), the inverse function h^{-1} exists on $h(I_s)$, i.e. $h^{-1}(x/y) = p/q$. Now assuming $p/q = h^{-1}(x/y) \in I_s$, we obtain the inequality (4.3), namely

$$\left| \xi_s - \frac{x}{y} \right| > C \frac{\log \log q}{q^2 \log q}.$$

The interval I_s has the form $I_s = [e^{1/s} - \alpha, e^{1/s} + \alpha]$, where $0 < \alpha < e^{1/s} - 1$. Hence, if $p/q \in I_s$, then $p > q(e^{1/s} - \alpha)$, so that we get $p^2 - q^2 > \beta q^2$ with $\beta := (e^{1/s} - \alpha)^2 - 1$. Thus we have $y = 4pq(p^2 - q^2) > 4q^4\beta\sqrt{1+\beta} > 4\beta q^4$ or $q^2 < \sqrt{y}/(2\sqrt{\beta})$. Then, for some positive constant C_{26} depending at most on s , we get

$$\left| \xi_s - \frac{x}{y} \right| > C_{26} \frac{\log \log y}{\sqrt{y} \log y} \quad (y \geq 3).$$

Since h is monotonously increasing on $(0, \infty)$, there exists a constant C_{27} such that the inequality $|\xi_s - x/y| > C_{27} > 0$ holds for $p/q = h^{-1}(x/y) \notin I_s$. \square

5. REMARKS ON THE PROOFS OF THEOREMS 1.6 - 1.9

In this section we sketch the proofs of Theorems 1.6 - 1.9. The arguments are always the same as in the proof of Theorem 1.5 given in section 4. Therefore we only mention the main formulas of the proofs.

PROOF OF THEOREM 1.6. This Theorem is based on Lemma 3.2.

Upper bound: We have

$$\begin{aligned} x_n &= P_n^2 - Q_n^2 - 2P_nQ_n, \\ y_n &= P_n^2 - Q_n^2 + 2P_nQ_n = (P_n + Q_n)^2 - 2Q_n^2, \\ z_n &= P_n^2 + Q_n^2. \end{aligned}$$

By $Q_n < P_n < 3Q_n$ ($n \geq 1$) we get $2Q_n^2 < y_n < 14Q_n^2$, and so

$$Q_n^2 > \frac{y_n}{14}, \quad \log Q_n > \frac{\log(y_n/14)}{2} > C_{28} \log y_n, \quad \log \log Q_n < \log \log y_n.$$

Lower bound: We have

$$\xi_s = h(e^{1/s}) = \frac{\sinh(1/s) - 1}{\sinh(1/s) + 1}$$

and

$$\begin{aligned} x &= p^2 - q^2 - 2pq, \\ y &= p^2 - q^2 + 2pq > 3, \\ z &= p^2 + q^2. \end{aligned}$$

The assumption $x/y < 1$ of the theorem implies that x/y belongs to the range of h . In addition, we have $I_s \subset [1, \infty)$. Thus, $I_s = [e^{1/s} - \alpha, e^{1/s} + \alpha]$, where $0 < \alpha < e^{1/s} - 1$. If $p/q \in I_s$, then $p > q(e^{1/s} - \alpha)$, and $y = (p+q)^2 - 2q^2 > \beta q^2$, where $\beta = (e^{1/s} - \alpha + 1)^2 - 2 > 2$. For $p/q = h^{-1}(x/y) \notin I_s$ and $p/q > \sqrt{2} - 1$, the inequality $|y\xi_s - x| > C > 0$ holds, since h is monotonously increasing on $(\sqrt{2} - 1, \infty)$. In particular, h^{-1} exists for every $\varepsilon > 0$ on $[\sqrt{2} - 1 + \varepsilon, \infty)$. \square

PROOF OF THEOREM 1.7. This Theorem is based on Lemma 3.3.

Upper bound: We have

$$\begin{aligned} x_n &= P_n(P_n^3 + 4Q_n^3), \\ y_n &= Q_n(Q_n^3 - 2P_n^3) < 0, \\ z_n &= P_n^6 - 10P_n^3Q_n^3 - 2Q_n^6. \end{aligned}$$

By $Q_n < P_n < 3Q_n$ ($n \geq 1$) we get $Q_n^4 < |y_n| < 53Q_n^4$, and so

$$Q_n^2 > \sqrt{\frac{|y_n|}{53}}, \quad \log Q_n > \frac{\log(|y_n|/53)}{4} > C_{29} \log |y_n|, \quad \log \log Q_n < \log \log |y_n|.$$

Lower bound: Let $\rho = \sqrt[3]{5 + 3\sqrt{3}}$. By the condition of the theorem

$$\frac{x}{y} \leq -\frac{(\sqrt[3]{5 + 3\sqrt{3}})(3 + \sqrt{3})}{3 + 2\sqrt{3}} = h(\rho),$$

so that $x/y(> 0)$ belongs to the range of h . We have

$$\begin{aligned} \xi_s &= h(e^{1/s}) = e^{1/s} \frac{4 + e^{3/s}}{1 - 2e^{3/s}} \\ &= \frac{\sinh(4/s) + \cosh(4/s) + 4\sinh(1/s) + 4\cosh(1/s)}{1 - 2\sinh(3/s) - 2\cosh(3/s)} < 0, \end{aligned}$$

and

$$\begin{aligned} x &= p(p^3 + 4q^3), \\ y &= q(q^3 - 2p^3) < 0, \\ z &= p^6 - 10p^3q^3 - 2q^6. \end{aligned}$$

In addition, $I_s \subset [\rho, \infty)$ ($s = 1$), $I_s \subset [1, \rho]$ ($s > 1$). Thus, $I_s = [e^{1/s} - \alpha, e^{1/s} + \alpha]$, where $0 < \alpha < e - \rho$ if $s = 1$, $0 < \alpha < e^{1/s} - 1$ if

$s > 1$. If $p/q \in I_s$, then $p > q(e^{1/s} - \alpha)$, and $|y| = q(2p^3 - q^3) > \beta q^4$, where $\beta = 2(e^{1/s} - \alpha)^3 - 1 > 1$. If $p/q \notin I_s$, we again distinguish the cases $s = 1$ and $s > 1$. If $s = 1$, we have the additional condition of the theorem that $p/q > \rho$. Since h is monotonously decreasing on (ρ, ∞) , there exists a constant C' such that $|\xi_s - x/y| > C' > 0$. If $s > 1$, we have $\sqrt[3]{1/2} < p/q < \rho$. Since h is monotonously increasing on $(\sqrt[3]{1/2}, \rho)$, there exists a constant C'' such that $|\xi_s - x/y| > C'' > 0$. \square

PROOF OF THEOREM 1.8. This Theorem is based on Lemma 3.4.

Upper bound: We have

$$\begin{aligned} x_n &= P_n^2 - Q_n^2, \\ y_n &= 2P_n Q_n + Q_n^2, \\ z_n &= P_n^2 + P_n Q_n + Q_n^2. \end{aligned}$$

Since $Q_n < P_n < 3Q_n$ ($n \geq 1$), we have $3Q_n^2 < y_n < 7Q_n^2$, and so

$$Q_n^2 > \frac{y_n}{7}, \quad \log Q_n > \frac{\log(y_n/7)}{2} > C_{30} \log y_n, \quad \log \log Q_n < \log \log y_n.$$

Lower bound: We have

$$\xi_s = h(e^{1/s}) = \frac{2 \sinh(1/s)}{2 + \cosh(1/s) - \sinh(1/s)}$$

and

$$\begin{aligned} x &= p^2 - q^2, \\ y &= 2pq + q^2, \\ z &= p^2 + pq + q^2. \end{aligned}$$

In addition, $I_s \subset [0, \infty)$. Hence, $I_s = [e^{1/s} - \alpha, e^{1/s} + \alpha]$, where $0 < \alpha < e^{1/s}$. If $p/q \in I_s$, then $p > q(e^{1/s} - \alpha)$, and $y = 2pq + q^2 > \beta q^2$, where $\beta = 2(e^{1/s} - \alpha) + 1 > 1$. Since h is monotonously increasing on $[0, \infty)$, there exists a constant C'_{30} such that the inequality $|\xi_s - x/y| > C'_{30} > 0$ holds for $p/q = h^{-1}(x/y) \notin I_s$. In particular, h^{-1} exists on $[0, \infty)$. \square

PROOF OF THEOREM 1.9. This Theorem is based on Lemma 3.5.

Upper bound: We have

$$\begin{aligned} x_n &= P_n^4 - Q_n^4, \\ y_n &= 4P_n^2 Q_n^2, \\ u_n &= P_n^2 + Q_n^2, \\ v_n &= 2(P_n^2 - Q_n^2)P_n Q_n. \end{aligned}$$

Since $Q_n < P_n < 3Q_n$ ($n \geq 1$), we have $4Q_n^4 < y_n < 36Q_n^4$, and so

$$Q_n^2 > \frac{\sqrt{y_n}}{6}, \quad \log Q_n > \frac{\log(y_n/36)}{4} > C_{31} \log y_n, \quad \log \log Q_n < \log \log y_n.$$

Lower bound: We have

$$\xi_s = h(e^{1/s}) = \frac{1}{2} \sinh\left(\frac{2}{s}\right)$$

and

$$\begin{aligned} x &= p^4 - q^4, \\ y &= 4p^2q^2, \\ u &= p^2 + q^2, \\ v &= 2(p^2 - q^2)pq. \end{aligned}$$

In addition, $I_s \subset (0, \infty)$, and so $I_s = [e^{1/s} - \alpha, e^{1/s} + \alpha]$, where $0 < \alpha < e^{1/s}$. If $p/q \in I_s$, then $p > q(e^{1/s} - \alpha)$, and $y = 4p^2q^2 > \beta q^4$, where $\beta := 4(e^{1/s} - \alpha)^2 > 0$. Since h is monotonously increasing on $(0, \infty)$, there exists a constant C'_{31} such that the inequality $|\xi_s - x/y| > C'_{31} > 0$ holds for $p/q = h^{-1}(x/y) \notin I_s$. In particular, h^{-1} exists for every $\varepsilon > 0$ on $[\varepsilon, \infty)$. \square

6. GENERALIZATION TO TRIGONOMETRIC FUNCTIONS

Let s be a positive integer. Then, the following continued fraction expansions are known:

$$\begin{aligned} \tan(1) &= [1; \overline{2k-1, 1}]_{k=1}^{\infty}, \\ \tan(1/s) &= [0; s-1, \overline{1, s-2+2ks}]_{k=1}^{\infty} \quad (s > 1). \end{aligned}$$

Let p_n/q_n be the n -th convergent of $\tan(1/s)$. Then, we have for $P_\nu := p_{2\nu}$ and $Q_\nu := q_{2\nu}$ that

$$\begin{aligned} \left| \tan(1) - \frac{P_\nu}{Q_\nu} \right| &< \frac{1}{(2\nu+1)Q_\nu^2} \quad (\nu \geq 0), \\ \left| \tan\left(\frac{1}{s}\right) - \frac{P_\nu}{Q_\nu} \right| &< \frac{1}{(s(2\nu+1)-2)Q_\nu^2} \quad (s > 1, \nu \geq 1). \end{aligned}$$

Applying the method given by [8, Corollary 1], we find three-term linear recurrence formulae for P_n and Q_n : For $s = 1$ we get with $P_0 = 1$, $P_1 = 3$, $Q_0 = 1$, $Q_1 = 2$ that

$$P_n = (2n+1)P_{n-1} - P_{n-2}, \quad Q_n = (2n+1)Q_{n-1} - Q_{n-2} \quad (n \geq 2),$$

whereas for $s > 1$ the recurrences start with $P_0 = 0$, $P_1 = 1$, $Q_0 = 1$, $Q_1 = s$:

$$P_n = s(2n-1)P_{n-1} - P_{n-2}, \quad Q_n = s(2n-1)Q_{n-1} - Q_{n-2} \quad (n \geq 2).$$

These recurrence formulae correspond to the well-known non-regular continued fraction

$$\tan(1/s) = \frac{1}{s - \frac{1}{3s - \frac{1}{5s - \ddots}}},$$

see [13, ch. 8, (27)]. By these facts, in connection with the results for $\exp(1/s)$ in (1.1) and (1.2), it follows that Lemma 1.1 holds for $h(e^{1/s})$ just as for $h(\tan(1/s))$. As a consequence, we may replace the hyperbolic functions in our preceding results by certain trigonometric functions.

THEOREM 6.1. *Let s be a positive integer. Then, the theorems listed in the following table hold for the numbers $h(e^{1/s})$ and $h(\tan(1/s))$ both.*

Theorem No.	$\xi_s = h(e^{1/s})$	$\xi_s = h(\tan(1/s))$
Proposition 1.2	$\sinh(1/s)$	$-\cot(2/s)$
Proposition 1.3	$\cosh(1/s)$	$\operatorname{cosec}(2/s)$
Proposition 1.4	$\tanh(1/s)$	$-\cos(2/s)$
Theorem 1.5	$(\sinh(1/s) - \operatorname{cosech}(1/s))/2$	$-\cot(4/s)$
Theorem 1.6	$\frac{\sinh(1/s) - 1}{\sinh(1/s) + 1}$	$\frac{1 + \tan(2/s)}{1 - \tan(2/s)}$
Theorem 1.9	$(\sinh(2/s))/2$	$-\operatorname{cosec}(2/s) \cot(2/s)$

A real irrational number ξ is said to be a *Liouville number*, if there is a sequence of rationals $(a_n/b_n)_{n>0}$ with $1 < b_1 < b_2 < \dots$ and

$$(6.1) \quad \left| \xi - \frac{a_n}{b_n} \right| < \frac{1}{b_n^n} \quad (n > 0).$$

REMARK 6.2. If ξ is a Liouville number and $\kappa(n)$ any strictly increasing sequence of positive integers satisfying $\kappa(n) \geq n$, then there is a sequence of rationals $(A_n/B_n)_{n>0}$ with $1 < B_1 < B_2 < \dots$ and

$$\left| \xi - \frac{A_n}{B_n} \right| < \frac{1}{B_n^{\kappa(n)}} \quad (n > 0).$$

This follows by setting $A_n := a_{\kappa(n)}$, $B_n := b_{\kappa(n)}$ ($n > 0$).

REMARK 6.3. When the inequality (6.1) holds for all subscripts $n > n_0$ only, ξ is a Liouville number defined by the shifted sequence of rationals a'_n/b'_n with $a'_n := a_{n+n_0}$, $b'_n := b_{n+n_0}$ ($n > 0$).

THEOREM 6.4. *Let α be a real number such that $\tan(\pi\alpha/2)$ is a Liouville number. Then there is a sequence of rationals $(p_n/q_n)_{n>0}$ with $1 < q_1 < q_2 < \dots$ and a sequence of positive integers $(r_n)_{n>0}$ satisfying*

$$\left| \sin(\pi\alpha) - \frac{p_n}{q_n} \right| < \frac{1}{q_n^2}, \quad q_n^2 = p_n^2 + r_n^2, \quad p_n \equiv 0 \pmod{2}.$$

In particular, $\sin(\pi\alpha)$ is a Liouville number.

PROOF. Without loss of generality we may assume that $0 < \alpha < 1$. From the hypothesis of the theorem it follows that there is a sequence of rationals $(a_n/b_n)_{n>0}$ with $1 < b_1 < b_2 < \dots$ satisfying

$$\left| \tan\left(\frac{\pi\alpha}{2}\right) - \frac{a_n}{b_n} \right| < \frac{1}{b_n^{3n+1}}.$$

For $0 < \alpha < 1$ the number $\tan(\pi\alpha/2)$ is positive. In what follows we separate our arguments according to the cases $\tan(\pi\alpha/2) < 1$ and $\tan(\pi\alpha/2) > 1$.

CASE 1. $\tan(\pi\alpha/2) < 1$

Then, for all sufficiently large subscripts $n > n_1$, we have $0 < a_n/b_n < 1$.

CASE 2. $\tan(\pi\alpha/2) > 1$

Here, we have for large subscripts $n > n_2$ that $1 < a_n/b_n < 1 + \tan(\pi\alpha/2)$. Set

$$f(x) := \frac{2x}{1+x^2}.$$

Then,

$$f'(x) = \frac{2(1-x^2)}{(1+x^2)^2}.$$

For $n > 0$ there is a real number η depending possibly on α and n such that

$$(6.2) \quad \left| f\left(\tan\left(\frac{\pi\alpha}{2}\right)\right) - f\left(\frac{a_n}{b_n}\right) \right| = |f'(\eta)| \left| \tan\left(\frac{\pi\alpha}{2}\right) - \frac{a_n}{b_n} \right|,$$

where either $a_n/b_n < \eta < \tan(\pi\alpha/2)$ or $\tan(\pi\alpha/2) < \eta < a_n/b_n$ holds. By our construction, the situation described in Case 1 yields $0 < \eta < 1$ for $n > n_1$, whereas in Case 2 we have $1 < \eta < 1 + \tan(\pi\alpha/2)$ for $n > n_2$. Hence

$$\begin{aligned} \frac{1}{2} |f'(\eta)| &= \frac{|1-\eta^2|}{(1+\eta^2)^2} \leq |1-\eta^2| \\ &= \begin{cases} 1-\eta^2 & \leq 1 & \text{(Case 1)} \\ \eta^2-1 & \leq \tan(\pi\alpha/2)(2+\tan(\pi\alpha/2)) & \text{(Case 2)} \end{cases}, \end{aligned}$$

or $|f'(\eta)| \leq C_\alpha := 2 \max\{1; \tan(\pi\alpha/2)(2+\tan(\pi\alpha/2))\}$. In the first case we have $0 < a_n < b_n$, and in the second case $0 < a_n < b_n(1+\tan(\pi\alpha/2))$, which is summarized by $0 < a_n < C_\alpha b_n$. Now, the right-hand side of (6.2) can be estimated as follows: Let $b_n > 1 + C_\alpha^2 > C_\alpha$, and $n > n_1$ or $n > n_2$, respectively. Then,

$$\begin{aligned} |f'(\eta)| \left| \tan\left(\frac{\pi\alpha}{2}\right) - \frac{a_n}{b_n} \right| &\leq \frac{C_\alpha}{b_n^{3n+1}} < \frac{1}{b_n^{3n}} < \frac{1}{b_n^{2n}(1+C_\alpha^2)^n} \\ &= \frac{1}{(b_n^2 + C_\alpha^2 b_n^2)^n} < \frac{1}{(b_n^2 + a_n^2)^n}. \end{aligned}$$

The left-hand side of (6.2) equals to

$$\left| \frac{2 \tan(\pi\alpha/2)}{1 + \tan^2(\pi\alpha/2)} - \frac{2a_n/b_n}{1 + a_n^2/b_n^2} \right| = \left| \sin(\pi\alpha) - \frac{2a_nb_n}{a_n^2 + b_n^2} \right|.$$

With $p_n = 2a_nb_n$, $q_n := a_n^2 + b_n^2$, and $r_n := |a_n^2 - b_n^2|$, we have proven for sufficiently large n that

$$\left| \sin(\pi\alpha) - \frac{p_n}{q_n} \right| < \frac{1}{q_n^n} \quad \text{with} \quad p_n^2 + r_n^2 = q_n^2,$$

and $(q_n)_{n>0}$ is a strictly increasing sequence of positive integers. To prove that $\sin(\pi\alpha)$ is a Liouville number, it suffices to show that it is not rational. But, assuming the contrary, it follows by

$$\sin(\pi\alpha) \tan^2\left(\frac{\pi\alpha}{2}\right) - 2 \tan\left(\frac{\pi\alpha}{2}\right) + \sin(\pi\alpha) = 0$$

that $\tan(\pi\alpha/2)$ is an algebraic number. But every Liouville number is transcendental, a contradiction. This completes the proof of the theorem. \square

Observing the identity

$$\cos(\pi\alpha) = \frac{1 - \tan^2(\pi\alpha/2)}{1 + \tan^2(\pi\alpha/2)},$$

the following result can be proven in a similar manner by applying the mean value theorem to the function $f(x) = (1 - x^2)/(1 + x^2)$.

THEOREM 6.5. *Let α be a real number such that $\tan(\pi\alpha/2)$ is a Liouville number. Then there is a sequence of rationals $(p_n/q_n)_{n>0}$ with $1 < q_1 < q_2 < \dots$ and a sequence of positive even integers $(r_n)_{n>0}$ satisfying*

$$\left| \cos(\pi\alpha) - \frac{p_n}{q_n} \right| < \frac{1}{q_n^n}, \quad q_n^2 = p_n^2 + r_n^2.$$

In particular, $\cos(\pi\alpha)$ is a Liouville number.

REFERENCES

- [1] E. B. Burger and A. M. Pillai, *On Diophantine approximation along algebraic curves*, Proc. Amer. Math. Soc. **136** (2008), 11-19.
- [2] H. Cohen, *Number Theory, vol.I: Tools and Diophantine Equations*, Springer, 2007.
- [3] H. Cohen, *Number Theory, vol.II: Analytic and Modern Tools*, Springer, 2007.
- [4] L. E. Dickson, *History of the theory of numbers, vol.2*, Dover Publications, 2005.
- [5] C. Elsner, *On arithmetic properties of the convergents of Euler's number*, Colloq. Math. **79** (1999), 133-145.
- [6] C. Elsner, *On rational approximations by Pythagorean numbers*, Fibonacci Quart. **41** (2003), 98-104.
- [7] C. Elsner, T. Komatsu, and I. Shiokawa, *Approximation of values of hypergeometric functions by restricted rationals*, J. Théorie des Nombres de Bordeaux. **19** (2007), 387-399.
- [8] C. Elsner and T. Komatsu, *A recurrence formula for leaping convergents of non-regular continued fractions*, Linear Algebra Appl. **428** (2008), 824-833.

- [9] C. Georgikopoulos, *Rational integral solutions of the equations $x^3 + 4y^3 = z^2$ and $x^3 + 2y^3 = z^2$* , Bull. Soc. Math. Grèce **24** (1948), 13-19.
- [10] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, fifth edition, Clarendon Press, Oxford, 1979.
- [11] T. Komatsu, *Arithmetical properties of the leaping convergents of $e^{1/s}$* , Tokyo J. Math. **27** (2004), 1-12.
- [12] L. J. Mordell, *Diophantine equations*, Academic Press, London and New York, 1969.
- [13] O. Perron, *Die Lehre von den Kettenbrüchen*, Chelsea, New York, 1950.

C. Elsner
FHDW, Fachhochschule für die Wirtschaft
University of Applied Sciences
Freundallee 15, D-30173 Hannover
Germany
E-mail: carsten.elsner@fhdw.de

T. Komatsu
Graduate School of Science and Technology
Hirosaki University
Hirosaki, 036-8561
Japan
E-mail: komatsu@cc.hirosaki-u.ac.jp

I. Shiokawa
Department of Mathematics
Keio University
Yokohama, 223-8522
Japan
E-mail: shiokawa@math.keio.ac.jp

Received: 10.9.2008.

Revised: 13.1.2009.