

## ON SOME FUNCTIONAL EQUATIONS RELATED TO DERIVATIONS AND BICIRCULAR PROJECTIONS IN RINGS

MAJA FOŠNER, BENJAMIN MARCEN, NEJC ŠIROVNIK AND JOSO VUKMAN  
University of Maribor and University of Primorska, Slovenia

ABSTRACT. In this paper we prove the following result. Let  $n \geq 1$  be some fixed integer and let  $R$  be a prime ring with  $2n < \text{char}(R) \neq 2$ . Suppose there exist additive mappings  $S, T : R \rightarrow R$  satisfying the relations

$$\begin{aligned} S(x^{2n}) &= S(x)x^{2n-1} + xT(x)x^{2n-2} + x^2S(x)x^{2n-3} + \cdots + x^{2n-1}T(x), \\ T(x^{2n}) &= T(x)x^{2n-1} + xS(x)x^{2n-2} + x^2T(x)x^{2n-3} + \cdots + x^{2n-1}S(x) \end{aligned}$$

for all  $x \in R$ . In this case  $S$  and  $T$  are of the form  $2S(x) = D(x) + \zeta(x)$ ,  $2T(x) = D(x) - \zeta(x)$  for all  $x \in R$ , where  $D : R \rightarrow R$  is a derivation and  $\zeta$  is an additive mapping, which maps  $R$  into its extended centroid. Besides,  $\zeta(x^{2n}) = 0$  for all  $x \in R$ . Functional equations related to bicircular projections are also investigated.

This research is a continuation of a recent work of M. Fošner and Vukman ([14]). Throughout the paper,  $R$  will represent an associative ring with center  $Z(R)$ . Given an integer  $n > 1$ , a ring  $R$  is said to be  $n$ -torsion free if for  $x \in R$ ,  $nx = 0$  implies  $x = 0$ . As usual the commutator  $xy - yx$  will be denoted by  $[x, y]$ . An additive mapping  $x \mapsto x^*$  on a ring  $R$  is called an involution if  $(xy)^* = y^*x^*$  and  $x^{**} = x$  hold for all pairs  $x, y \in R$ . A ring equipped with an involution is called a ring with involution or  $*$ -ring. Recall that a ring  $R$  is prime if for  $a, b \in R$ ,  $aRb = (0)$  implies that either  $a = 0$  or  $b = 0$  and is semiprime in case  $aRa = (0)$  implies  $a = 0$ . We denote by  $\text{char}(R)$  the characteristic of a prime ring  $R$ . An additive mapping  $D : R \rightarrow R$ , where  $R$  is an arbitrary ring, is called a derivation if  $D(xy) = D(x)y + xD(y)$  holds for all pairs  $x, y \in R$  and is called a Jordan derivation in case  $D(x^2) = D(x)x + xD(x)$

---

2010 *Mathematics Subject Classification.* 16R60, 16W10, 39B05.

*Key words and phrases.* Derivation, functional identity, bicircular projection.

is fulfilled for all  $x \in R$ . A derivation  $D$  is inner in case there exists  $a \in R$  such that  $D(x) = [a, x]$  holds for all  $x \in R$ .

Every derivation is a Jordan derivation. The converse is in general not true. A classical result of Herstein ([15]) asserts that any Jordan derivation on a 2-torsion free prime ring is a derivation. Cusack ([8]) generalized Herstein theorem to 2-torsion free semiprime rings.

We denote by  $Q_{mr}$ ,  $Q_S$  and  $C$  the maximal right ring of quotients, symmetric Martindale ring of quotients and extended centroid of a semiprime ring  $R$ , respectively. For the explanation of  $Q_{mr}$ ,  $Q_S$  and  $C$  we refer the reader to [1]. Given some  $X \subset R$  we denote  $C(X) = \{r \in R \mid [r, X] = 0\}$ .

Let  $X$  be a complex Banach space and let  $\mathcal{L}(X)$  be the algebra of all bounded linear operators on  $X$ . A projection  $P \in \mathcal{L}(X)$  is bicircular in case all mappings of the form  $e^{i\alpha}P + e^{i\beta}(I - P)$ , where  $I$  denotes the identity operator, are isometric for all pairs of real numbers  $\alpha, \beta$ .

Beidar, Brešar, Chebotar and Martindale ([3]) have proved the following result, which fairly generalizes Herstein theorem.

**THEOREM 1.** *Let  $n > 1$  be some fixed integer and let  $R$  be a prime ring with  $\text{char}(R) \neq 2$ . Suppose there exists an additive mapping  $D : R \rightarrow R$  satisfying the relation*

$$D(x^n) = \sum_{i=1}^n x^{i-1} D(x) x^{n-i}$$

for all  $x \in R$ . In this case  $D$  is a derivation.

Recently, M. Fošner and Vukman ([14]) proved the following result.

**THEOREM 2.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a prime ring with  $2n \leq \text{char}(R) \neq 2$ . Suppose there exists an additive mapping  $T : R \rightarrow R$  satisfying the relation*

$$(1) \quad T(x^{2n+1}) = \sum_{i=1}^{2n+1} (-1)^{i+1} x^{i-1} T(x) x^{2n+1-i}$$

for all  $x \in R$ . In this case  $T$  is of the form  $T(x) = qx + xq$  for all  $x \in R$  and some fixed  $q \in Q_S$ .

It seems natural to ask what can be proved in case we have an even number in the relation (1). More precisely, we are talking about the solution of the functional equation

$$T(x^{2n}) = \sum_{i=1}^{2n} (-1)^{i+1} x^{i-1} T(x) x^{2n-i}.$$

It is our aim in this paper to prove the following result.

**THEOREM 3.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a prime ring with  $2n < \text{char}(R) \neq 2$ . Suppose there exists an additive mapping  $T : R \rightarrow R$  satisfying*

$$(2) \quad T(x^{2n}) = \sum_{i=1}^{2n} (-1)^{i-1} x^{i-1} T(x) x^{2n-i}$$

for all  $x \in R$ . In this case  $T$  maps  $R$  into  $C$  and  $T(x^{2n}) = 0$  for all  $x \in R$ .

In the proof of Theorem 3 we use as the main tool the theory of functional identities (Beidar - Brešar - Chebotar theory). The theory of functional identities considers set-theoretic maps on rings that satisfy some identical relations. When treating such relations, one usually concludes that the form of the maps involved can be described, unless the ring is very special. For the full treatment on this theory, we refer the reader to [7].

For the proof of Theorem 3 we need Theorem 4, which might be of independent interest. Let  $R$  be an algebra over a commutative ring  $\xi$  and let

$$p(x_1, x_2, \dots, x_{2n}) = \sum_{\pi \in S_{2n}} x_{\pi(1)} x_{\pi(2)} \cdots x_{\pi(2n)}$$

be a fixed multilinear polynomial in noncommuting indeterminates  $x_1, x_2, \dots, x_{2n}$ . Further, let  $\mathcal{L}$  be a subset of  $R$  closed under  $p$ , which means  $p(\bar{x}_{2n}) \in \mathcal{L}$  for all  $x_1, x_2, \dots, x_{2n} \in \mathcal{L}$ , where  $\bar{x}_{2n} = (x_1, x_2, \dots, x_{2n})$ . We shall consider a mapping  $T : \mathcal{L} \rightarrow R$  satisfying

$$(3) \quad T(p(\bar{x}_{2n})) = \sum_{\pi \in S_{2n}} \sum_{i=1}^{2n} (-1)^{i-1} x_{\pi(1)} \cdots x_{\pi(i-1)} T(x_{\pi(i)}) x_{\pi(i+1)} \cdots x_{\pi(2n)}$$

for all  $x_1, x_2, \dots, x_{2n} \in \mathcal{L}$ . Let us mention that the idea of considering the expression  $[p(\bar{x}_{2n}), p(\bar{y}_{2n})]$  in its proof is taken from [2].

**THEOREM 4.** *Let  $\mathcal{L}$  be a  $4n$ -free Lie subring of  $R$  closed under  $p$ . If  $T : \mathcal{L} \rightarrow R$  is an additive mapping satisfying (3), then  $T$  maps  $\mathcal{L}$  into  $C$  and  $T(x^{2n}) = 0$  for all  $x \in \mathcal{L}$ .*

**PROOF.** Let us write  $k = 2n$  for brevity. Note that for any  $a \in R$  and  $\bar{x}_k \in \mathcal{L}^k$ , we have

$$[p(\bar{x}_k), a] = \sum_{i=1}^k p(x_1, x_2, \dots, x_{i-1}, [x_i, a], x_{i+1}, \dots, x_k).$$

Thus

$$T[p(\bar{x}_k), a] = \sum_{i=1}^k T(p(x_1, x_2, \dots, x_{i-1}, [x_i, a], x_{i+1}, \dots, x_k)).$$

Let  $f, g : \mathbb{N} \rightarrow \mathbb{N}$  be mappings with properties  $f(1) = g(k) = 0$  and  $f(i) = 1$  for  $i = 2, 3, \dots, k$  and  $g(i) = 1$  for  $i = 1, 2, \dots, k-1$ . Therefore

$$\begin{aligned} & T[p(\bar{x}_k), a] \\ &= \sum_{\pi \in S_k} \sum_{i=1}^k (-1)^{i-1} \left( f(i)[x_{\pi(1)} \dots x_{\pi(i-1)}, a] T(x_{\pi(i)}) x_{\pi(i+1)} \dots x_{\pi(k)} \right. \\ &\quad \left. + x_{\pi(1)} \dots x_{\pi(i-1)} T[x_{\pi(i)}, a] x_{\pi(i+1)} \dots x_{\pi(k)} \right. \\ &\quad \left. + g(i)x_{\pi(1)} \dots x_{\pi(i-1)} T(x_{\pi(i)}) [x_{\pi(i+1)} \dots x_{\pi(k)}, a] \right). \end{aligned}$$

In particular, we have

$$\begin{aligned} & T[p(\bar{x}_k), p(\bar{y}_k)] \\ (4) \quad &= \sum_{\pi \in S_k} \sum_{i=1}^k (-1)^{i-1} \left( f(i)[x_{\pi(1)} \dots x_{\pi(i-1)}, p(\bar{y}_k)] T(x_{\pi(i)}) x_{\pi(i+1)} \dots x_{\pi(k)} \right. \\ &\quad \left. + x_{\pi(1)} \dots x_{\pi(i-1)} T[x_{\pi(i)}, p(\bar{y}_k)] x_{\pi(i+1)} \dots x_{\pi(k)} \right. \\ &\quad \left. + g(i)x_{\pi(1)} \dots x_{\pi(i-1)} T(x_{\pi(i)}) [x_{\pi(i+1)} \dots x_{\pi(k)}, p(\bar{y}_k)] \right). \end{aligned}$$

For  $i = 1, 2, \dots, k$  let us denote  $\varphi(x_{\pi(i)}) = T[x_{\pi(i)}, p(\bar{y}_k)]$ . We therefore have

$$\begin{aligned} \varphi(x_{\pi(i)}) &= T[x_{\pi(i)}, p(\bar{y}_k)] = -T[p(\bar{y}_k), x_{\pi(i)}] \\ &= \sum_{\sigma \in S_k} \sum_{j=1}^k (-1)^{j-1} \left( f(j)[x_{\pi(i)}, y_{\sigma(1)} \dots y_{\sigma(j-1)}] T(y_{\sigma(j)}) y_{\sigma(j+1)} \dots y_{\sigma(k)} \right. \\ &\quad \left. + y_{\sigma(1)} \dots y_{\sigma(j-1)} T[x_{\pi(i)}, y_{\sigma(j)}] y_{\sigma(j+1)} \dots y_{\sigma(k)} \right. \\ &\quad \left. + g(j)y_{\sigma(1)} \dots y_{\sigma(j-1)} T(y_{\sigma(j)}) [x_{\pi(i)}, y_{\sigma(j+1)} \dots y_{\sigma(k)}] \right). \end{aligned}$$

Therefore (4) can be written as

$$\begin{aligned} & T[p(\bar{x}_k), p(\bar{y}_k)] \\ &= \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} f(i)[x_{\pi(1)} \dots x_{\pi(i-1)}, y_{\sigma(1)} \dots y_{\sigma(k)}] \cdot \\ &\quad \cdot T(x_{\pi(i)}) x_{\pi(i+1)} \dots x_{\pi(k)} \\ (5) \quad &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} x_{\pi(1)} \dots x_{\pi(i-1)} \varphi(x_{\pi(i)}) x_{\pi(i+1)} \dots x_{\pi(k)} \\ &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} g(i)x_{\pi(1)} \dots x_{\pi(i-1)} T(x_{\pi(i)}) \cdot \\ &\quad \cdot [x_{\pi(i+1)} \dots x_{\pi(k)}, y_{\sigma(1)} \dots y_{\sigma(k)}]. \end{aligned}$$

If we replace the roles of denotations  $\pi$  and  $\sigma$ , we obtain from (5) that

$$\begin{aligned}
 & T[p(\bar{x}_k), p(\bar{y}_k)] \\
 &= \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} f(i)[x_{\pi(1)} \cdots x_{\pi(k)}, y_{\sigma(1)} \cdots y_{\sigma(i-1)}] \cdot \\
 & \quad \cdot T(y_{\sigma(i)} y_{\sigma(i+1)} \cdots y_{\sigma(k)}) \\
 (6) \quad &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} y_{\sigma(1)} \cdots y_{\sigma(i-1)} \bar{\varphi}(y_{\sigma(i)} y_{\sigma(i+1)} \cdots y_{\sigma(k)}) \\
 &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} g(i) y_{\sigma(1)} \cdots y_{\sigma(i-1)} T(y_{\sigma(i)}) \cdot \\
 & \quad \cdot [x_{\pi(1)} \cdots x_{\pi(k)}, y_{\sigma(i+1)} \cdots y_{\sigma(k)}],
 \end{aligned}$$

where  $\bar{\varphi}(y_{\sigma(i)}) = T[p(\bar{x}_k), y_{\sigma(i)}]$ . One can easily check that

$$\bar{\varphi}(x_{\pi(i)}) = -\varphi(x_{\pi(i)})$$

for all  $i = 1, 2, \dots, k$ . Comparing (5) and (6) we obtain the identity

$$\begin{aligned}
 0 = & \\
 = & \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} f(i)[x_{\pi(1)} \cdots x_{\pi(i-1)}, y_{\sigma(1)} \cdots y_{\sigma(k)}] \cdot \\
 & \quad \cdot T(x_{\pi(i)} x_{\pi(i+1)} \cdots x_{\pi(k)}) \\
 &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} x_{\pi(1)} \cdots x_{\pi(i-1)} \varphi(x_{\pi(i)} x_{\pi(i+1)} \cdots x_{\pi(k)}) \\
 &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} g(i) x_{\pi(1)} \cdots x_{\pi(i-1)} T(x_{\pi(i)}) \cdot \\
 & \quad \cdot [x_{\pi(i+1)} \cdots x_{\pi(k)}, y_{\sigma(1)} \cdots y_{\sigma(k)}] \\
 &- \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} f(i)[x_{\pi(1)} \cdots x_{\pi(k)}, y_{\sigma(1)} \cdots y_{\sigma(i-1)}] \cdot \\
 & \quad \cdot T(y_{\sigma(i)} y_{\sigma(i+1)} \cdots y_{\sigma(k)}) \\
 &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} y_{\sigma(1)} \cdots y_{\sigma(i-1)} \varphi(y_{\sigma(i)} y_{\sigma(i+1)} \cdots y_{\sigma(k)}) \\
 &- \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \sum_{i=1}^k (-1)^{i-1} g(i) y_{\sigma(1)} \cdots y_{\sigma(i-1)} T(y_{\sigma(i)}) \cdot \\
 & \quad \cdot [x_{\pi(1)} \cdots x_{\pi(k)}, y_{\sigma(i+1)} \cdots y_{\sigma(k)}]
 \end{aligned}$$

for all  $x_1, \dots, x_k, y_1, \dots, y_k \in \mathcal{L}$ . The last relation can be written as

$$\begin{aligned}
0 = & \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \left( T(y_{\sigma(1)})y_{\sigma(2)} \cdots y_{\sigma(k)}x_{\pi(1)} \cdots x_{\pi(k-1)} \right. \\
& + \varphi(x_{\pi(1)})x_{\pi(2)} \cdots x_{\pi(k-1)} \\
& - T(x_{\pi(1)})y_{\sigma(1)} \cdots y_{\sigma(k)}x_{\pi(2)} \cdots x_{\pi(k-1)} \\
& + \sum_{i=2}^{k-1} (-1)^{i-1} x_{\pi(1)} \cdots x_{\pi(i-1)} \varphi(x_{\pi(i)})x_{\pi(i+1)} \cdots x_{\pi(k-1)} \\
& + \sum_{i=2}^{k-1} (-1)^{i-1} [x_{\pi(1)} \cdots x_{\pi(i-1)}, y_{\sigma(1)} \cdots y_{\sigma(k)}] \cdot \\
& \quad \left. T(x_{\pi(i)})x_{\pi(i+1)} \cdots x_{\pi(k-1)} \right) x_{\pi(k)} \\
& + \sum_{\pi \in S_k} \sum_{\sigma \in S_k} \left( T(x_{\pi(1)})x_{\pi(2)} \cdots x_{\pi(k)}y_{\sigma(1)} \cdots y_{\sigma(k-1)} \right. \\
& + \varphi(y_{\sigma(1)})y_{\sigma(2)} \cdots y_{\sigma(k-1)} \\
& - T(y_{\sigma(1)})x_{\pi(1)} \cdots x_{\pi(k)}y_{\sigma(2)} \cdots y_{\sigma(k-1)} \\
& + \sum_{i=2}^{k-1} (-1)^{i-1} y_{\sigma(1)} \cdots y_{\sigma(i-1)} \varphi(y_{\sigma(i)})y_{\sigma(i+1)} \cdots y_{\sigma(k-1)} \\
(7) \quad & + \sum_{i=2}^{k-1} (-1)^{i-1} [y_{\sigma(1)} \cdots y_{\sigma(i-1)}, x_{\pi(1)} \cdots x_{\pi(k)}] \cdot \\
& \quad \left. T(y_{\sigma(i)})y_{\sigma(i+1)} \cdots y_{\sigma(k-1)} \right) y_{\sigma(k)} \\
& + \sum_{\pi \in S_k} \sum_{\sigma \in S_k} x_{\pi(1)} \left( -x_{\pi(2)} \cdots x_{\pi(k-1)}y_{\sigma(1)} \cdots y_{\sigma(k)}T(x_{\pi(k)}) \right. \\
& + x_{\pi(2)} \cdots x_{\pi(k)}y_{\sigma(1)} \cdots y_{\sigma(k-1)}T(y_{\sigma(k)}) \\
& - x_{\pi(2)} \cdots x_{\pi(k-1)}\varphi(x_{\pi(k)}) \\
& + \sum_{i=2}^{k-1} (-1)^{i-1} x_{\pi(2)} \cdots x_{\pi(i-1)}T(x_{\pi(i)}) \cdot \\
& \quad \left. [x_{\pi(i+1)} \cdots x_{\pi(k)}, y_{\sigma(1)} \cdots y_{\sigma(k)}] \right) \\
& + \sum_{\pi \in S_k} \sum_{\sigma \in S_k} y_{\sigma(1)} \left( -y_{\sigma(2)} \cdots y_{\sigma(k-1)}x_{\pi(1)} \cdots x_{\pi(k)}T(y_{\sigma(k)}) \right. \\
& + y_{\sigma(2)} \cdots y_{\sigma(k)}x_{\pi(1)} \cdots x_{\pi(k-1)}T(x_{\pi(k)}) \\
& - y_{\sigma(2)} \cdots y_{\sigma(k-1)}\varphi(y_{\sigma(k)}) \\
& + \sum_{i=2}^{k-1} (-1)^{i-1} y_{\sigma(2)} \cdots y_{\sigma(i-1)}T(y_{\sigma(i)}) \cdot \\
& \quad \left. [y_{\sigma(i+1)} \cdots y_{\sigma(k)}, x_{\pi(1)} \cdots x_{\pi(k)}] \right)
\end{aligned}$$

for all  $x_1, \dots, x_k, y_1, \dots, y_k \in \mathcal{L}$ . Let us define mappings  $E, F : \mathcal{L}^{2k-1} \rightarrow R$  by the rule

$$\begin{aligned} E(u_1, u_2, u_3, \dots, u_{2k-1}) &= T(u_k)u_{k+1} \dots u_{2k-1}u_1 \dots u_{k-1} \\ &+ \varphi(u_1)u_2 \dots u_{k-1} - T(u_1)u_k \dots u_{2k-1}u_2u_3 \dots u_{k-1} \\ &+ \sum_{i=2}^{k-1} (-1)^{i-1}u_1 \dots u_{i-1}\varphi(u_i)u_{i+1} \dots u_{k-1} \\ &+ \sum_{i=2}^{k-1} (-1)^{i-1}[u_1 \dots u_{i-1}, u_k \dots u_{2k-1}]T(u_i)u_{i+1} \dots u_{k-1} \end{aligned}$$

and

$$\begin{aligned} F(u_1, u_2, u_3, \dots, u_{2k-1}) &= -u_1u_2 \dots u_{k-2}u_ku_{k+1} \dots u_{2k-1}T(u_{k-1}) \\ &+ u_1u_2 \dots u_{k-1}u_ku_{k+1} \dots u_{2k-2}T(u_{2k-1}) - u_1u_2 \dots u_{k-2}\varphi(u_{k-1}) \\ &+ \sum_{i=1}^{k-2} (-1)^{i-1}u_1u_2 \dots u_{i-1}T(u_i)[u_{i+1} \dots u_{k-1}, u_k \dots u_{2k-1}] \end{aligned}$$

for all  $\bar{u}_{2k-1} \in \mathcal{L}^{2k-1}$ . Accordingly, (7) can be rewritten as

$$\begin{aligned} 0 &= \sum_{\pi \in S_k} \sum_{\sigma \in S_k} E(x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k-1)}, y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)})x_{\pi(k)} \\ &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} E(y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k-1)}, x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k)})y_{\sigma(k)} \\ &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} x_{\pi(1)}F(x_{\pi(2)}, x_{\pi(3)}, \dots, x_{\pi(k)}, y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)}) \\ &+ \sum_{\pi \in S_k} \sum_{\sigma \in S_k} y_{\sigma(1)}F(y_{\sigma(2)}, y_{\sigma(3)}, \dots, y_{\sigma(k)}, x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k)}) \end{aligned}$$

and hence

$$\begin{aligned} 0 &= \sum_{i=1}^k \left( \sum_{\substack{\pi \in S_k \\ \pi(k)=i}} \sum_{\sigma \in S_k} E(x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k-1)}, y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)}) \right) x_i \\ &+ \sum_{i=k+1}^{2k} \left( \sum_{\pi \in S_k} \sum_{\substack{\sigma \in S_k \\ \sigma(k)=i}} E(y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k-1)}, x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k)}) \right) y_i \\ &+ \sum_{j=1}^k x_j \left( \sum_{\substack{\pi \in S_k \\ \pi(1)=j}} \sum_{\sigma \in S_k} F(x_{\pi(2)}, x_{\pi(3)}, \dots, x_{\pi(k)}, y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)}) \right) \end{aligned}$$

$$+ \sum_{j=k+1}^{2k} y_j \left( \sum_{\pi \in S_k} \sum_{\substack{\sigma \in S_k \\ \sigma(1)=j}} F(y_{\sigma(2)}, y_{\sigma(3)}, \dots, y_{\sigma(k)}, x_{\pi(1)}, x_{\pi(2)}, \dots, x_{\pi(k)}) \right).$$

Let  $s : \mathbb{Z} \rightarrow \mathbb{Z}$  be a mapping defined by  $s(i) = i - k$ . For each  $\sigma \in S_k$  the mapping  $s^{-1}\sigma s : \{k+1, \dots, 2k\} \rightarrow \{k+1, \dots, 2k\}$  will be denoted by  $\bar{\sigma}$ . Writting  $x_{k+i}$  instead of  $y_i$ ,  $i = 1, 2, \dots, k$  in the above identity, we can express this relation as

$$\sum_{i=1}^{2k} E_i(\bar{x}_{2k}^i) x_i + \sum_{j=1}^{2k} x_j F_j(\bar{x}_{2k}^j) = 0.$$

According to the theory of functional identities (see [7, Chapter 3.2]) there exist mappings  $p_{k,i} : \mathcal{L}^{k-2} \rightarrow R$ ,  $i = 1, \dots, k-1$  and  $\lambda_k : \mathcal{L}^{k-1} \rightarrow C(\mathcal{L})$  such that

$$\begin{aligned} & \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} \sum_{\sigma \in S_k} F(x_{\pi(2)}, x_{\pi(3)}, \dots, x_{\pi(k)}, x_{\bar{\sigma}(k+1)}, x_{\bar{\sigma}(k+2)}, \dots, x_{\bar{\sigma}(2k)}) \\ &= \sum_{i=1}^{k-1} p_{k,i}(\bar{x}_{k-1}^i) x_i + \lambda_k(\bar{x}_{k-1}) \end{aligned}$$

for all  $\bar{x}_{k-1} \in \mathcal{L}^{k-1}$ . Recalling the definition of a mapping  $F$  leads to

$$\begin{aligned} & \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} \sum_{\sigma \in S_k} x_{\pi(2)} \left( -x_{\pi(3)} \dots x_{\pi(k-1)} y_{\sigma(1)} \dots y_{\sigma(k)} T(x_{\pi(k)}) \right. \\ & \quad \left. + x_{\pi(3)} \dots x_{\pi(k)} y_{\sigma(1)} \dots y_{\sigma(k-1)} T(y_{\sigma(k)}) \right. \\ & \quad \left. - x_{\pi(3)} \dots x_{\pi(k-1)} \varphi(x_{\pi(k)}) \right) \\ & + \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} \sum_{\sigma \in S_k} \left( \sum_{i=2}^{k-1} (-1)^{i-1} x_{\pi(2)} \dots x_{\pi(i-1)} \cdot \right. \\ & \quad \left. \cdot T(x_{\pi(i)}) x_{\pi(i+1)} \dots x_{\pi(k)} y_{\sigma(1)} \dots y_{\sigma(k-1)} \right) y_{\sigma(k)} \\ & - \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} \sum_{\sigma \in S_k} \left( \sum_{i=2}^{k-1} (-1)^{i-1} x_{\pi(2)} \dots x_{\pi(i-1)} \cdot \right. \\ & \quad \left. \cdot T(x_{\pi(i)}) y_{\sigma(1)} \dots y_{\sigma(k)} x_{\pi(i+1)} \dots x_{\pi(k-1)} \right) x_{\pi(k)} \\ & - \sum_{i=1}^{k-1} p_{k,i}(\bar{x}_{k-1}^i) x_i \in C(\mathcal{L}) \end{aligned}$$



for all  $\bar{x}_{k-1} \in \mathcal{L}^{k-1}$ . Applying the theory of functional identities it follows that

$$\begin{aligned} & \sum_{\substack{\pi \in S_k \\ \pi(1)=1 \\ \pi(2)=2}} \sum_{\sigma \in S_k} -x_{\pi(3)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k)} T(x_{\pi(k)}) \\ & \quad + x_{\pi(3)} \cdots x_{\pi(k)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T(y_{\sigma(k)}) \\ & \quad - x_{\pi(3)} \cdots x_{\pi(k-1)} \varphi(x_{\pi(k)}) \\ & - \sum_{i=1}^{k-2} p_{k,i}(\bar{x}_{k-2}^i) x_i \in C(\mathcal{L}) \end{aligned}$$

for all  $\bar{x}_{k-1} \in \mathcal{L}^{k-1}$ . Recalling the definition of a mapping  $\varphi(x_{\pi(k)})$  leads to

$$\begin{aligned} & \sum_{\substack{\pi \in S_k \\ \pi(1)=1 \\ \pi(2)=2}} \sum_{\sigma \in S_k} x_{\pi(3)} \left( -x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k)} T(x_{\pi(k)}) \right. \\ & \quad + x_{\pi(4)} \cdots x_{\pi(k)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T(y_{\sigma(k)}) \\ & \quad + x_{\pi(4)} \cdots x_{\pi(k)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T(y_{\sigma(k)}) \\ & \quad - x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} x_{\pi(k)} T(y_{\sigma(k)}) \\ & \quad \left. + x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T[x_{\pi(k)}, y_{\sigma(k)}] \right) \\ & + \sum_{\substack{\pi \in S_k \\ \pi(1)=1 \\ \pi(2)=2}} \sum_{\sigma \in S_k} \sum_{j=1}^{k-1} \left( x_{\pi(3)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(j-1)} \right. \\ & \quad \left. \cdot T(y_{\sigma(j)}) y_{\sigma(j+1)} \cdots y_{\sigma(k)} \right) x_{\pi(k)} \\ & + \sum_{\substack{\pi \in S_k \\ \pi(1)=1 \\ \pi(2)=2}} \sum_{\sigma \in S_k} \sum_{j=1}^{k-1} \left( -f(j) x_{\pi(3)} \cdots x_{\pi(k-1)} [x_{\pi(i)}, y_{\sigma(1)} \cdots y_{\sigma(j-1)}] \right. \\ & \quad \cdot T(y_{\sigma(j)}) y_{\sigma(j+1)} \cdots y_{\sigma(k-1)} \\ & \quad - x_{\pi(3)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(j-1)} T[x_{\pi(k)}, y_{\sigma(j)}] y_{\sigma(j+1)} \cdots y_{\sigma(k-1)} \\ & \quad \left. - x_{\pi(3)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(j-1)} T(y_{\sigma(j)}) x_{\pi(k)} y_{\sigma(j+1)} \cdots y_{\sigma(k-1)} \right) y_{\sigma(k)} \\ & - \sum_{i=1}^{k-2} p_{k,i}(\bar{x}_{k-2}^i) x_i \in C(\mathcal{L}). \end{aligned}$$

Applying the theory of functional identities gives

$$\begin{aligned} & \sum_{\substack{\pi \in S_k \\ \pi(1)=1 \\ \pi(2)=2 \\ \pi(3)=3}} \sum_{\sigma \in S_k} -x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k)} T(x_{\pi(k)}) \\ & \quad + x_{\pi(4)} \cdots x_{\pi(k)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T(y_{\sigma(k)}) \\ & \quad + x_{\pi(4)} \cdots x_{\pi(k)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T(y_{\sigma(k)}) \\ & \quad - x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} x_{\pi(k)} T(y_{\sigma(k)}) \\ & \quad + x_{\pi(4)} \cdots x_{\pi(k-1)} y_{\sigma(1)} \cdots y_{\sigma(k-1)} T[x_{\pi(k)}, y_{\sigma(k)}] \\ & - \sum_{i=1}^{k-3} p_{k,i}(\bar{x}_{k-3}^i) x_i \in C(\mathcal{L}). \end{aligned}$$

After finite number of steps we arrive at

$$(8) \quad 2T(x) = px + \mu(x)$$

for all  $x \in \mathcal{L}$ , where  $p \in \mathcal{L}$  and  $\mu : \mathcal{L} \rightarrow C(\mathcal{L})$ . Putting  $x^k$  for  $x$  in the above relation leads to

$$(9) \quad 2T(x^k) = px^k + \mu(x^k).$$

Combining the above relation and (2) we obtain

$$0 = px^k + \mu(x^k) - \sum_{i=1}^k (-1)^{i-1} x^{i-1} (px + \mu(x)) x^{k-i}$$

and few calculations lead to

$$0 = \mu(x^k) - \mu(x)x^{k-1} - \sum_{i=2}^k (-1)^{i-1} x^{i-1} (px + \mu(x)) x^{k-i}.$$

Since  $\mu(x) \in C(\mathcal{L})$  for all  $x \in \mathcal{L}$ , the above relation reduces to

$$(10) \quad 0 = \mu(x^k) - \sum_{i=2}^k (-1)^{i-1} x^{i-1} px^{k+1-i}.$$

After a complete linearization of the last relation we obtain

$$\begin{aligned} 0 = & \mu(x_{\pi(1)} \cdots x_{\pi(k)}) + \sum_{\pi \in S_k} x_{\pi(1)} \left( px_{\pi(2)} \cdots x_{\pi(k)} - x_{\pi(2)} px_{\pi(3)} \cdots x_{\pi(k)} \right. \\ & \left. + \cdots - x_{\pi(2)} \cdots x_{\pi(k-2)} px_{\pi(k-1)} x_{\pi(k)} + x_{\pi(2)} \cdots x_{\pi(k-1)} px_{\pi(k)} \right). \end{aligned}$$

The theory of functional identities implies

$$0 = \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} \left( px_{\pi(2)} \cdots x_{\pi(k-1)} \right) x_{\pi(k)}$$

$$\begin{aligned}
 &+ \sum_{\substack{\pi \in S_k \\ \pi(1)=1}} x_{\pi(2)} \left( -px_{\pi(3)} \cdots x_{\pi(k)} + \cdots - x_{\pi(3)} \cdots x_{\pi(k-2)}px_{\pi(k-1)}x_{\pi(k)} \right. \\
 &\quad \left. + x_{\pi(3)} \cdots x_{\pi(k-1)}px_{\pi(k)} \right).
 \end{aligned}$$

After a finite number of steps we obtain

$$(11) \quad px = xq + \lambda(x)$$

for all  $x \in \mathcal{L}$ , where  $q \in \mathcal{L}$  and  $\lambda : \mathcal{L} \rightarrow C(\mathcal{L})$ . Right multiplication of the relation (11) by  $y \in \mathcal{L}$  gives  $pxy = xqy + \lambda(x)y$ . Putting  $xy$  for  $x$  in the relation (11) leads to  $pxy = xyq + \lambda(xy)$ . Comparing the last two relations we obtain

$$0 = x[q, y] + \lambda(x)y - \lambda(xy),$$

which implies  $[q, y] = 0$  for all  $y \in \mathcal{L}$ . Note that (11) can now be rewritten as  $(p - q)x = \lambda(x)$ . From the last relation we obtain  $p = q$  and  $\lambda(x) = 0$ . Considering that  $p \in C(\mathcal{L})$  in the relation (10) gives

$$0 = px^k + \mu(x^k).$$

Considering the above relation in (9) gives  $T(x^k) = 0$  for all  $x \in \mathcal{L}$ . After a complete linearization of the above relation and using the theory of functional identities, we obtain  $p = 0$ . The relation (8) now implies that  $2T(x) = \mu(x)$ , which means that  $T(x) \in C(\mathcal{L})$ . Thereby the proof is complete.  $\square$

We are now in the position to prove Theorem 3.

**PROOF OF THEOREM 3.** The complete linearization of (2) gives (3). Assume first that  $R$  is not a PI ring. According to Theorem 4 we have  $T(R) \in C$ . Now suppose that  $R$  is a PI ring. It is well-known that in this case  $R$  has a nonzero center (see [21]). Let  $c$  be a nonzero central element and let us write  $k$  for  $2n$  for brevity. Putting  $x_1 = x_2 = \dots = x_k = c$  in (3) leads to

$$T(c^k) = 0.$$

Pick any  $x \in R$  and set  $x_1 = cx^2$  and  $x_2 = x_3 = \dots = x_k = c$  in (3). Hence we obtain

$$\begin{aligned}
 k!T(c^kx^2) &= (k-1)!T(cx^2)c^{k-1} + (k-1)(k-1)!T(c)x^2c^{k-1} \\
 &\quad - (k-1)!T(cx^2)c^{k-1} - (k-1)!x^2T(c)c^{k-1} - (k-2)(k-1)!T(c)x^2c^{k-1} \\
 &\quad + (k-1)!T(cx^2)c^{k-1} + 2(k-1)!x^2T(c)c^{k-1} + (k-3)(k-1)!T(c)x^2c^{k-1} \\
 &\quad - \dots \\
 &\quad + (k-1)!T(cx^2)c^{k-1} + (k-2)(k-1)!x^2T(c)c^{k-1} + (k-1)!T(c)x^2c^{k-1} \\
 &\quad - (k-1)!T(cx^2)c^{k-1} - (k-1)(k-1)!x^2T(c)c^{k-1}.
 \end{aligned}$$

The above relation reduces to

$$(12) \quad 2k!T(c^k x^2) = k(k-1)!c^{k-1} \left( T(c)x^2 - x^2T(c) \right).$$

Putting  $x_1 = x_2 = cx$  and  $x_3 = x_4 = \dots = x_k = c$  in (3) gives, after some calculations,

$$2k!T(c^k x^2) = k(k-2)!c^{k-1} \left( 2T(cx)x - 2xT(cx) + (k-2)(T(c)x^2 - x^2T(c)) \right).$$

Comparing the last two relations leads to

$$(13) \quad 2T(cx)x - 2xT(cx) = T(c)x^2 - x^2T(c).$$

Putting  $x_1 = x$  and  $x_2 = c^{k-1}$  in the complete linearization of the above relation we obtain

$$T(c^k)x - xT(c^k) = T(c)xc^{k-1} - xT(c)c^{k-1}.$$

Since  $T(c^k) = 0$ , it follows from the above relation that

$$(14) \quad T(c)x = xT(c).$$

Considering the above relation in (13) gives

$$(15) \quad T(cx)x = xT(cx).$$

Putting  $x_1 = x$ ,  $x_2 = c^2x$  and  $x_3 = x_4 = \dots = x_k = c$  in (3) and also considering (14) and (15) gives

$$2k!T(c^k x^2) = k(k-2)!c^{k-2} \left( T(x)xc^2 - xT(x)c^2 \right).$$

Since  $T(c^k x^2) = 0$  by relations (12) and (14), the above relation reduces to

$$(16) \quad [T(x), x] = 0$$

for all  $x \in R$ . Considering the above relation in (2) leads to

$$(17) \quad T(x^k) = 0$$

for all  $x \in R$ . The relation (16) implies the existence of such  $\lambda \in R$  and  $\mu : R \rightarrow C$  that

$$(18) \quad T(x) = \lambda x + \mu(x)$$

for all  $x \in R$ . By (17) we also have

$$0 = \lambda x^k + \mu(x^k)$$

for all  $x \in R$ . Therefore  $\lambda x^k \in C$ , whence it follows that

$$(19) \quad [\lambda x^k, y] = 0$$

for all  $x, y \in R$ . Putting  $x_1 = x_2 = \dots = x_k = c$  in the complete linearization of the above relation gives  $[\lambda, y] = 0$  for all  $y \in R$ , which implies that  $\lambda \in Z(R)$ . Now putting  $x_1 = x$  and  $x_2 = x_3 = \dots = x_k = c$  in the complete

linearization of the relation (19) leads to  $[\lambda x, y] = 0$ . Left multiplication of the last relation by  $z \in R$  and considering  $\lambda \in Z(R)$  gives  $\lambda z[x, y] = 0$  for all  $x, y, z \in R$ . Since  $R$  is prime, it follows that  $\lambda = 0$  or  $[x, y] = 0$  for all  $x, y \in R$ . If  $\lambda = 0$  the relation (18) gives  $T(x) = \mu(x)$  and therefore  $T(x) \in C$  for all  $x \in R$ . In case  $[x, y] = 0$  it is obvious that  $[T(x), y] = 0$  for all  $x, y \in R$ , which means that  $T(x) \in Z(R)$ . The proof of the theorem is therefore complete.  $\square$

We proceed with the following result.

**THEOREM 5.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a prime ring with  $2n < \text{char}(R) \neq 2$ . Suppose there exist additive mappings  $S, T : R \rightarrow R$  satisfying the relations*

$$(20) \quad \begin{aligned} S(x^{2n}) &= S(x)x^{2n-1} + xT(x)x^{2n-2} + x^2S(x)x^{2n-3} + \dots + x^{2n-1}T(x), \\ T(x^{2n}) &= T(x)x^{2n-1} + xS(x)x^{2n-2} + x^2T(x)x^{2n-3} + \dots + x^{2n-1}S(x). \end{aligned}$$

*In this case  $S$  and  $T$  are of the form*

$$\begin{aligned} 2S(x) &= D(x) + \zeta(x), \\ 2T(x) &= D(x) - \zeta(x) \end{aligned}$$

*for all  $x \in R$ , where  $D : R \rightarrow R$  is a derivation and  $\zeta : R \rightarrow C$  is an additive mapping such that  $\zeta(x^{2n}) = 0$  for all  $x \in R$ .*

**PROOF.** Combining relations (20) we obtain

$$D(x^{2n}) = D(x)x^{2n-1} + xD(x)x^{2n-2} + x^2D(x)x^{2n-3} + \dots + x^{2n-1}D(x),$$

where  $D$  stands for  $S + T$ . According to the above relation and Theorem 1,  $D$  is a derivation. Subtracting relations (20) we obtain

$$(21) \quad \zeta(x^{2n}) = \zeta(x)x^{2n-1} - x\zeta(x)x^{2n-2} + x^2\zeta(x)x^{2n-3} + \dots - x^{2n-1}\zeta(x),$$

where  $\zeta$  denotes  $S - T$ . From (21) and Theorem 3 it follows that  $\zeta$  maps  $R$  into  $C$  and  $\zeta(x^{2n}) = 0$  for all  $x \in R$ . We therefore have  $S + T = D$  and  $S - T = \zeta$ , whence it follows that  $2S(x) = D(x) + \zeta(x)$ ,  $2T(x) = D(x) - \zeta(x)$  for all  $x \in R$ , which completes the proof of the theorem.  $\square$

Let us point out that in Theorem 3 we have not assumed that the ring has an identity element. In case the ring has an identity element, the proof is considerably simpler and one can prove the result below even in case  $R$  is an arbitrary ring with some torsion restrictions.

**THEOREM 6.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a  $2n!$ -free semiprime ring with the identity element. Suppose there exists an additive mapping  $T : R \rightarrow R$  satisfying*

$$(22) \quad T(x^{2n}) = \sum_{i=1}^{2n} (-1)^{i-1} x^{i-1} T(x) x^{2n-i}$$

*for all  $x \in R$ . In this case  $T(x) = 0$  for all  $x \in R$ .*

PROOF. Let  $e$  be the identity element. Putting  $e$  for  $x$  in the relation (22) gives

$$(23) \quad T(e) = 0.$$

Let  $y$  be any element of  $Z(R)$ . Linearization in the relation (22) leads to

$$\begin{aligned} \sum_{i=0}^{2n} \binom{2n}{i} T(x^{2n-i}y^i) &= T(x+y) \left( \sum_{i=0}^{2n-1} \binom{2n-1}{i} x^{2n-1-i}y^i \right) \\ &\quad - (x+y)T(x+y) \left( \sum_{i=0}^{2n-2} \binom{2n-2}{i} x^{2n-2-i}y^i \right) \\ &\quad + (x^2 + 2xy + y^2)T(x+y) \left( \sum_{i=0}^{n-3} \binom{n-3}{i} x^{n-3-i}y^i \right) \\ &\quad - \dots \\ &\quad - \left( \sum_{i=0}^{n-3} \binom{n-3}{i} x^{n-3-i}y^i \right) T(x+y)(x^2 + 2xy + y^2) \\ &\quad + \left( \sum_{i=0}^{2n-2} \binom{2n-2}{i} x^{2n-2-i}y^i \right) T(x+y)(x+y) \\ &\quad - \left( \sum_{i=0}^{2n-1} \binom{2n-1}{i} x^{2n-1-i}y^i \right) T(x+y). \end{aligned}$$

Using (22) and rearranging the above relation in sense of collecting together terms involving equal number of factors of  $y$ , we obtain

$$\sum_{i=1}^{2n-1} f_i(x, y) = 0,$$

where  $f_i(x, y)$  stands for the expression of terms involving  $i$  factors of  $y$ . Replacing  $x$  by  $x + 2y$ ,  $x + 3y$ ,  $\dots$ ,  $x + (2n - 1)y$  in turn in the relation (22) and expressing the resulting system of  $2n - 1$  homogeneous equations of variables  $f_i(x, y)$ ,  $i = 1, 2, \dots, 2n - 1$ , we see that the coefficient matrix of the system is a van der Monde matrix

$$\begin{bmatrix} 1 & 1 & \dots & 1 \\ 2 & 2^2 & \dots & 2^{2n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 2n-1 & (2n-1)^2 & \dots & (2n-1)^{2n-1} \end{bmatrix}.$$

Since the determinant of this matrix is different from zero, it follows that the system has only a trivial solution. In particular,

$$\begin{aligned}
 f_{2n-1}(x, e) = & -\binom{2n}{2n-1}T(x) \\
 & + T(x) + \binom{2n-1}{2n-2}T(e)x \\
 & - T(x) - xT(e) - \binom{2n-2}{2n-3}T(e)x \\
 & + T(x) + 2xT(e) + \binom{2n-3}{2n-4}T(e)x \\
 & - \dots \\
 & - T(x) - \binom{2n-3}{2n-4}xT(e) - 2T(e)x \\
 & + T(x) + \binom{2n-2}{2n-3}xT(e) + T(e)x \\
 & - T(x) - \binom{2n-1}{2n-2}xT(e).
 \end{aligned}$$

The relation (23) reduces the above relation to

$$2nT(x) = 0$$

and since  $R$  is  $2n!$ -torsion free, we obtain  $T(x) = 0$  for all  $x \in R$ . The proof of the theorem is therefore complete.  $\square$

Kosi-Ulbl and Vukman ([19]) have proved the following result.

**THEOREM 7.** *Let  $n > 1$  be some fixed integer and let  $R$  be a  $n!$ -torsion free semiprime ring with the identity element. Suppose there exists an additive mapping  $D : R \rightarrow R$  satisfying the relation*

$$D(x^n) = \sum_{i=1}^n x^{i-1}D(x)x^{n-i}$$

for all  $x \in R$ . In this case  $D$  is a derivation.

Applying Theorem 6 and Theorem 7 we obtain the result below.

**THEOREM 8.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a  $2n!$ -torsion free semiprime ring with the identity element. Suppose there exist additive mappings  $S, T : R \rightarrow R$  satisfying the relations*

$$\begin{aligned}
 S(x^{2n}) &= S(x)x^{2n-1} + xT(x)x^{2n-2} + x^2S(x)x^{2n-3} + \dots + x^{2n-1}T(x), \\
 T(x^{2n}) &= T(x)x^{2n-1} + xS(x)x^{2n-2} + x^2T(x)x^{2n-3} + \dots + x^{2n-1}S(x)
 \end{aligned}$$

for all  $x \in R$ . In this case  $S$  and  $T$  are derivations and  $S = T$ .

Stachó and Zalar ([22,23]) investigated bicircular projections on  $C^*$ -algebra  $\mathcal{L}(H)$ , where  $H$  is a complex Hilbert space. According to [22, Proposition 3.4] every bicircular projection  $P : \mathcal{L}(H) \rightarrow \mathcal{L}(H)$  satisfies the functional equation

$$(24) \quad P(ABA) = P(A)BA - AP(B^*)^*A + ABP(A)$$

for all pairs  $A, B \in \mathcal{L}(H)$ , where  $B^*$  stands for the adjoint operator of  $B \in \mathcal{L}(H)$ . Fošner and Ilišević ([11]) investigated the above functional equation on 2-torsion free semiprime  $*$ -ring. They expressed the solution of the equation (24) in terms of derivation and so-called double centralizers. Bicircular projections and related functional equations have been extensively investigated during the last few years (see [5, 6, 9–14, 16–18, 20, 24]). M. Fošner and Vukman ([13]) investigated the following system of functional equations on prime  $*$ -rings with  $\text{char}(R) \neq 2$ .

$$\begin{aligned} P(x^3) &= P(x)x^2 + xQ(x^*)^*x + x^2P(x), \\ Q(x^3) &= Q(x)x^2 + xP(x^*)^*x + x^2Q(x). \end{aligned}$$

Recently, in [14] they considered the following much more general situation

$$\begin{aligned} P(x^{2n+1}) &= P(x)x^{2n} + xQ(x^*)^*x^{2n-1} + x^2P(x)x^{2n} + \cdots + x^{2n}P(x), \\ Q(x^{2n+1}) &= Q(x)x^{2n} + xP(x^*)^*x^{2n-1} + x^2Q(x)x^{2n} + \cdots + x^{2n}Q(x). \end{aligned}$$

In this paper we prove the following theorem.

**THEOREM 9.** *Let  $n \geq 1$  be some fixed integer and let  $R$  be a prime  $*$ -ring with  $2n < \text{char}(R) \neq 2$ . Suppose there exist additive mappings  $P, Q : R \rightarrow R$  satisfying the relations*

$$(25) \quad P(x^{2n}) = P(x)x^{2n-1} + xQ(x^*)^*x^{2n-2} + \cdots + x^{2n-1}Q(x^*)^*,$$

$$(26) \quad Q(x^{2n}) = Q(x)x^{2n-1} + xP(x^*)^*x^{2n-2} + \cdots + x^{2n-1}P(x^*)^*$$

for all  $x \in R$ . In this case  $P$  and  $Q$  are of the form

$$\begin{aligned} 4P(x) &= D(x) + G(x) + \zeta(x) + \theta(x), \\ 4Q(x) &= D(x) - G(x) + \zeta(x) - \theta(x) \end{aligned}$$

for all  $x \in R$ , where  $D, G : R \rightarrow R$  are derivations with properties  $D(x^*)^* = D(x)$ ,  $G(x^*)^* = -G(x)$  and  $\zeta, \theta : R \rightarrow C$  are additive mappings with properties  $\zeta(x^*)^* = -\zeta(x)$ ,  $\theta(x^*)^* = \theta(x)$ ,  $\zeta(x^{2n}) = \theta(x^{2n}) = 0$  for all  $x \in R$ .

**PROOF.** The proof goes through in three steps.

**FIRST STEP.** Let us first assume that  $Q = P$ . In this case, we have the relation

$$(27) \quad P(x^{2n}) = P(x)x^{2n-1} + xP(x^*)^*x^{2n-2} + \cdots + x^{2n-1}P(x^*)^*$$

for all  $x \in R$ . It is our aim to prove that  $P$  is of the form

$$2P(x) = D(x) + \zeta(x)$$

for all  $x \in R$ , where  $D : R \rightarrow R$  is a derivation and  $\zeta : R \rightarrow C$  is an additive mapping such that  $\zeta(x^{2n}) = 0$  for all  $x \in R$ . Besides,  $D(x^*)^* = D(x)$  and  $\zeta(x^*)^* = -\zeta(x)$  for all  $x \in R$ . Let us introduce mappings  $D, \zeta : R \rightarrow R$  by

$$(28) \quad D(x) = P(x) + P(x^*)^*,$$

$$(29) \quad \zeta(x) = P(x) - P(x^*)^*$$



for all  $x \in R$ . We therefore have  $D(x^*)^* = (P(x^*) + P(x^*)^*)^* = P(x) + P(x^*)^* = D(x)$ . Hence,

$$D(x^*)^* = D(x)$$

for all  $x \in R$  and similarly we obtain

$$\zeta(x^*)^* = -\zeta(x)$$

for all  $x \in R$ . From (27) one can easily obtain that

$$(30) \quad D(x^{2n}) = \sum_{i=1}^{2n} x^{i-1} D(x) x^{2n-i}$$

and

$$(31) \quad \zeta(x^{2n}) = \sum_{i=1}^{2n} (-1)^{i+1} x^{i-1} \zeta(x) x^{2n-i}$$

for all  $x \in R$ . Now it follows from (30) and Theorem 1 that  $D$  is a derivation. On the other hand, one can conclude from (31) and Theorem 3 that  $\zeta$  maps  $R$  into  $C$  and

$$\zeta(x^{2n}) = 0$$

for all  $x \in R$ . Combining (28) and (29) gives

$$2P(x) = D(x) + \zeta(x)$$

for all  $x \in R$ , which completes the proof of the first step.

SECOND STEP. Let us now assume that  $Q = -P$ , which according to (25) and (26) gives

$$(32) \quad Q(x^{2n}) = Q(x)x^{2n-1} - xQ(x^*)^*x^{2n-2} + \dots - x^{2n-1}Q(x^*)^*$$

for all  $x \in R$ . In this case,  $Q$  is of the form

$$2Q(x) = G(x) + \theta(x)$$

for all  $x \in R$ , where  $G : R \rightarrow R$  is a derivation and  $\theta : R \rightarrow C$  is an additive mapping such that  $\theta(x^{2n}) = 0$  for all  $x \in R$ . Besides,  $G(x^*)^* = -G(x)$  and  $\theta(x^*)^* = \theta(x)$  for all  $x \in R$ . The proof of the second step goes through by using the same arguments as in the first step and will therefore be omitted.

THIRD STEP. We are now in the position to prove the theorem in its full generality. We have the relations

$$\begin{aligned} P(x^{2n}) &= P(x)x^{2n-1} + xQ(x^*)^*x^{2n-2} + \dots + x^{2n-1}Q(x^*)^*, \\ Q(x^{2n}) &= Q(x)x^{2n-1} + xP(x^*)^*x^{2n-2} + \dots + x^{2n-1}P(x^*)^* \end{aligned}$$

for all  $x \in R$ . Adding (subtracting) the above relations gives, respectively,

$$\begin{aligned} F(x^{2n}) &= F(x)x^{2n-1} + xF(x^*)^*x^{2n-2} + \dots + x^{2n-1}F(x^*)^*, \\ H(x^{2n}) &= H(x)x^{2n-1} - xH(x^*)^*x^{2n-2} + \dots - x^{2n-1}H(x^*)^* \end{aligned}$$

for all  $x \in R$ , where  $F$  denotes  $P+Q$  and  $H$  stands for  $P-Q$ . Now according to the results regarding (27) and (32) in first and second step, we obtain from the above relations that

$$\begin{aligned} 2P(x) + 2Q(x) &= D(x) + \zeta(x), \\ 2P(x) - 2Q(x) &= G(x) + \theta(x) \end{aligned}$$

for all  $x \in R$ , where  $D, G : R \rightarrow R$  are derivations with properties  $D(x^*)^* = D(x)$ ,  $G(x^*)^* = -G(x)$  and  $\zeta, \theta : R \rightarrow C$  are additive mappings with properties  $\zeta(x^*)^* = -\zeta(x)$ ,  $\theta(x^*)^* = \theta(x)$ ,  $\zeta(x^{2n}) = \theta(x^{2n}) = 0$  for all  $x \in R$ . The last two relations imply

$$\begin{aligned} 4P(x) &= D(x) + G(x) + \zeta(x) + \theta(x), \\ 4Q(x) &= D(x) - G(x) + \zeta(x) - \theta(x) \end{aligned}$$

for all  $x \in R$ , which completes the proof of the theorem.  $\square$

#### ACKNOWLEDGEMENTS.

The authors would like to thank the referee for providing valuable comments and suggestions.

#### REFERENCES

- [1] K. I. Beidar, W. S. Martindale 3rd and A. V. Mikhalev, *Rings with generalized identities*, Marcel Dekker, Inc., New York, (1996).
- [2] K. I. Beidaar and Y.-Fong, *On additive isomorphisms of prime rings preserving polynomials*, J. Algebra **217** (1999), 650–667.
- [3] K. I. Beidar, M. Brešar, M. A. Chebotar and W. S. Martindale 3rd, *On Herstein's Lie map Conjectures II*, J. Algebra **238** (2001), 239–264.
- [4] K. I. Beidar, A. V. Mikhalev and M. A. Chebotar, *Functional identities in rings and their applications*, Russian Math. Surveys **59** (2004), 403–428.
- [5] F. Botelho and J. Jamison, *Generalized bicircular projections on minimal ideals of operators*, Proc. Amer. Math. Soc. **136** (2008), 1397–1402.
- [6] F. Botelho and J. Jamison, *Generalized bi-circular projections on spaces of analytic functions*, Acta Sci. Math. (Szeged) **75** (2009), 527–546.
- [7] M. Brešar, M. A. Chebotar and W. S. Martindale 3rd, *Functional identities*, Birkhäuser Verlag, Basel-Boston-Berlin (2007).
- [8] J. Cusack, *Jordan derivations on rings*, Proc. Amer. Math. Soc. **53** (1975), 321–324.
- [9] A. Fošner and J. Vukman, *Some functional equations on standard operator algebras*, Acta Math. Hungar. **118** (2008), 299–306.
- [10] A. Fošner and D. Ilišević, *Generalized bicircular projections via rank preserving maps on the spaces of symmetric and antisymmetric operators*, Oper. Matrices **5** (2011), 239–260.
- [11] M. Fošner and D. Ilišević, *On a class of projections on \*-rings*, Comm. Algebra **33** (2005), 3293–3310.
- [12] M. Fošner, D. Ilišević and C.-K. Li, *G-invariant norms and bicircular projections*, Linear Algebra Appl. **420** (2007), 596–608.
- [13] M. Fošner and J. Vukman, *On some equations in prime rings*, Monatsh. Math. **152** (2007), 135–150.
- [14] M. Fošner and J. Vukman, *On some functional equations in rings*, Commun. Algebra **39** (2011), 2647–2658.

- [15] I. N. Herstein, *Jordan derivations of prime rings*, Proc. Amer. Math. Soc. **8** (1957), 1104–1110.
- [16] D. Ilišević, *Generalized bicircular projections via the operator equation  $\alpha X^2 + AY^2 + \beta XAY + A = 0$* , Linear Algebra Appl. **429** (2008), 2025–2029.
- [17] D. Ilišević, *Generalized bicircular projections on  $JB^*$ -triples*, Linear Algebra Appl. **432** (2010), 1267–1276.
- [18] J. Jamison, *Generalized bicircular projections on spaces of operator an  $JB^*$ -triples*, Rocky Mountain. J. Math. **41** (2011), 1241–1245.
- [19] I. Kosi-Ulbl and J. Vukman, *A note on derivations in semiprime rings*, Int. J. Math. & Math. Sci. **20** (2005), 3347–3350.
- [20] I. Kosi-Ulbl and J. Vukman, *On some functional equations on standard operator algebras*, Glas. Mat. Ser. III **44(64)** (2009), 447–455.
- [21] L. H. Rowen, *Some results on the center of a ring with polynomial identity*, Bull. Amer. Math. Soc. **79** (1973), 219–223.
- [22] L. L. Stachó and B. Zalar, *Bicircular projections on some matrix and operator spaces*, Linear Algebra Appl. **384** (2004), 9–20.
- [23] L. L. Stachó and B. Zalar, *Bicircular projections and characterization of Hilbert spaces*, Proc. Amer. Math. Soc. **132** (2004), 3019–3025.
- [24] J. Vukman, *On functional equations related to bicircular projections*, Glas. Mat. Ser. III **41(61)** (2006), 51–55.

M. Fošner  
Faculty of Logistics  
University of Maribor  
Mariborska cesta 7, 3000 Celje  
Slovenia  
*E-mail:* [maja.fosner@fl.uni-mb.si](mailto:maja.fosner@fl.uni-mb.si)

B. Marcen  
Faculty of Logistics  
University of Maribor  
Mariborska cesta 7, 3000 Celje  
Slovenia  
*E-mail:* [benjamin.marcen@fl.uni-mb.si](mailto:benjamin.marcen@fl.uni-mb.si)

N. Širovnik  
Faculty of Natural Sciences and Mathematics  
University of Maribor  
Koroška cesta 160, 2000 Maribor  
Slovenia  
*E-mail:* [nejc.sirovnik@uni-mb.si](mailto:nejc.sirovnik@uni-mb.si)

J. Vukman  
Faculty of Mathematics, Natural Sciences and Information Technologies  
University of Primorska  
Glagoljaška 8, 6000 Koper  
and  
Institute of Mathematics, Physics and Mechanics  
Jadranska 19, 1000 Ljubljana  
Slovenia  
*E-mail:* [joso.vukman@uni-mb.si](mailto:joso.vukman@uni-mb.si), [joso.vukman@gmail.com](mailto:joso.vukman@gmail.com)

*Received:* 28.1.2014.

*Revised:* 11.4.2014.