

SELF-DUALITY IN THE CASE OF $SO(2n, F)$

DUBRAVKA BAN

University of Split, Croatia and Purdue University, USA

ABSTRACT. The parabolically induced representations of special even-orthogonal groups over p -adic field are considered. The main result is a theorem on self-duality, which gives a condition on initial representations, if induced representation has a square integrable subquotient.

1. INTRODUCTION

The problem of construction of noncuspidal irreducible square integrable representations of classical p -adic groups was studied by M. Tadić in [T3]. He showed ([T3], Lemma 4.1) that among irreducible cuspidal representations of general linear groups only the self-dual play a role in the construction of irreducible noncuspidal square integrable representations of symplectic and odd-orthogonal groups.

In this paper, we show the same property for groups $SO(2n, F)$ (Theorem 6.1). In the second section, we review some notation and results from the representation theory of general linear groups. In the third section, we describe standard parabolics of $SO(2n, F)$. Some properties of induced representations of $SO(2n, F)$ are given in the fourth section. The fifth section exposes the Casselman square integrability criterion for $SO(2n, F)$. In the sixth section, a theorem on self-duality is stated and proved.

By closing the introduction, I would like to thank Marko Tadić, who initiated this paper and helped its realization. I also thank Goran Muić for his helpful comments regarding this paper.

1991 *Mathematics Subject Classification.* 20G05, 22E50.

Key words and phrases. self-duality, parabolic induction, representations, even-orthogonal groups.

2. PRELIMINARIES

Fix a locally compact nonarchimedean field F of characteristic different from 2. Let G be a group of F -points of a connected reductive F -split group. Suppose that G is reductive and split.

Fix a minimal parabolic subgroup $P_0 \subset G$ and a maximal split torus $A_0 \subset P_0$.

Let P be a parabolic subgroup, containing P_0 . We call such a group a standard parabolic subgroup. Let U be the unipotent radical of P . Then, by [BZ], there exists a unique Levi subgroup M in P containing A_0 .

Let P be a standard parabolic subgroup of G , with Levi decomposition $P = MU$. For a smooth representation σ of M , we denote by $i_{G,M}(\sigma)$ the parabolically induced representation of G by σ from P , and for a smooth representation π of G , we denote by $r_{M,G}(\pi)$ the normalised Jacquet module of π with respect to P .

For a smooth finite length representation π we denote by $s.s.(\pi)$ the semi-simplified representation of π . The equivalence $s.s.(\pi_1) \cong s.s.(\pi_2)$ means that π_1 and π_2 have the same irreducible composition factors with the same multiplicities, and we write $\pi_1 = \pi_2$. We write $\pi_1 \cong \pi_2$ if we mean that π_1 and π_2 are actually equivalent.

Now we shall recall some results from [BZ] and [Z] of the representation theory of general linear groups.

For the group $GL(n, F)$, we fix the minimal parabolic subgroup which consists of all upper triangular matrices in $GL(n, F)$. The standard parabolic subgroups of $GL(n, F)$ can be parametrized by ordered partitions of n : for $\alpha = (n_1, \dots, n_k)$ there exists a standard parabolic subgroup (denote it in this section by P_α) of $GL(n, F)$ whose Levi factor M_α is naturally isomorphic to $GL(n_1, F) \times \dots \times GL(n_k, F)$.

Let π_1, π_2 be admissible representation of $GL(n_1, F), GL(n_2, F)$ resp., $n_1 + n_2 = n$. Define

$$\pi_1 \times \pi_2 = i_{GL(n,F), M(n_1+n_2)}(\pi_1 \otimes \pi_2).$$

Denote $\nu = |\det|$. We have the following criterion for irreducibility ([Z], Proposition 1.11):

PROPOSITION 2.1. *Let $\pi_i, i = 1, 2$, be irreducible cuspidal representation of $GL(n_i, F)$.*

1. *If $\pi_1 \not\cong \nu\pi_2$ and $\pi_2 \not\cong \nu\pi_1$ (in particular if $n_1 \neq n_2$), then $\pi_1 \times \pi_2$ is irreducible.*
2. *Suppose that $n_1 = n_2$ and either $\pi_1 \cong \nu\pi_2$ or $\pi_2 \cong \nu\pi_1$. Then the representation $\pi_1 \times \pi_2$ has length 2.*

3. PARABOLIC INDUCTION FOR $SO(2n, F)$

The special orthogonal group $SO(2n, F)$, $n \geq 1$, is the group

$$SO(2n, F) = \{X \in SL(2n, F) \mid {}^\tau X X = I_{2n}\}.$$

Here ${}^\tau X$ denotes the transposed matrix of X with respect to the second diagonal. For $n = 1$ we get

$$SO(2, F) = \left\{ \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} \mid \lambda \in F^\times \right\} \cong F^\times.$$

$SO(0, F)$ is defined to be the trivial group.

Denote by A_0 the maximal split torus in $SO(2n, F)$ which consists of all diagonal matrices in $SO(2n, F)$. Hence,

$$A_0 = \{diag(x_1, \dots, x_n, x_n^{-1}, \dots, x_1^{-1}) \mid x_i \in F^\times\} \cong (F^\times)^n.$$

Fix the minimal parabolic subgroup P_0 which consists of all upper triangular matrices in $SO(2n, F)$.

The root system is of type D_n ; the simple roots are

$$\begin{aligned} \alpha_i &= e_i - e_{i+1}, \quad \text{for } 1 \leq i \leq n-1, \\ \alpha_n &= e_{n-1} + e_n. \end{aligned}$$

The set of simple roots is denoted by Δ .

Let

$$s = \begin{bmatrix} I & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & I \end{bmatrix} \in O(2n, F).$$

We use the same letter s to denote the automorphism of $SO(2n, F)$ defined by $s(g) = sgs^{-1}$.

Let $\theta = \Delta \setminus \{\alpha_i\}$, $i \in \{1, \dots, n\}$, and let $P_\theta = M_\theta U_\theta$ be the maximal parabolic subgroup determined by θ .

If $i \neq n-1$, then

$$M_\theta = \{diag(g, h, {}^\tau g^{-1}) \mid g \in GL(i, F), h \in SO(2(n-i), F)\}.$$

In this case, we denote M_θ by $M_{(i)}$, and we have

$$M_{(i)} \cong GL(i, F) \times SO(2(n-i), F).$$

If $i = n-1$, then

$$M_\theta = s(M_{(n)}).$$

Now let $\theta = \Delta \setminus \{\alpha_{n-1}, \alpha_n\}$. Then

$$M_\theta = \{diag(g, h, {}^\tau g^{-1}) \mid g \in GL(n-1, F), h \in SO(2, F) \cong GL(1, F)\},$$

so

$$\begin{aligned} M_\theta &\cong GL(n-1, F) \times SO(2, F), \\ M_\theta &\cong GL(n-1, F) \times GL(1, F), \end{aligned}$$

and we denote M_θ by $M_{(n-1)}$ or by $M_{(n-1,1)}$.

We shall now describe the set of standard parabolic subgroups of $SO(2n, F)$. Let $\alpha = (n_1, \dots, n_k)$ be an ordered partition of non-negative integer $m \leq n$. Then there exists a standard parabolic subgroup, denote it by $P_\alpha = M_\alpha U_\alpha$, such that

$$M_\alpha = \{ \text{diag}(g_1, \dots, g_k, h, {}^\tau g_k^{-1}, \dots, {}^\tau g_1^{-1}) \mid g_i \in GL(n_i, F), \\ h \in SO(2(n-m), F) \}.$$

Hence,

$$M_\alpha \cong GL(n_1, F) \times GL(n_2, F) \times \dots \times GL(n_k, F) \times SO(2(n-m), F).$$

Mention that if $n_1 + \dots + n_k = n-1$, then

$$M = \{ \text{diag}(g_1, \dots, g_k, h, {}^\tau g_k^{-1}, \dots, {}^\tau g_1^{-1}) \mid g_i \in GL(n_i, F), \\ h \in SO(2, F) \cong GL(1, F) \},$$

so we may consider

$$M \cong GL(n_1, F) \times GL(n_2, F) \times \dots \times GL(n_k, F) \times SO(2, F),$$

or

$$M \cong GL(n_1, F) \times GL(n_2, F) \times \dots \times GL(n_k, F) \times GL(1, F).$$

Hence, we can assign

$$M \mapsto \alpha = (n_1, \dots, n_k),$$

or

$$M \mapsto \alpha' = (n_1, \dots, n_k, 1).$$

Besides the subgroups of type $P_\alpha = M_\alpha U_\alpha$, there is also another type of standard parabolic subgroups. They can be described as

$$M = s(M'),$$

where $M' = M_\alpha$, for some $\alpha = (n_1, \dots, n_k)$, $n_1 + \dots + n_k = n$.

Now, take smooth finite length representations π of $GL(n, F)$ and σ of $SO(2m, F)$. Let $P_{(n)} = M_{(n)} U_{(n)}$ be a standard parabolic subgroup of $G = SO(2(m+n), F)$. Hence, $M_{(n)} \cong GL(n, F) \times SO(2m, F)$, so $\pi \otimes \sigma$ can be taken as a representation of $M_{(n)}$. Define

$$\pi \rtimes \sigma = i_{M_{(n)}, G}(\pi \otimes \sigma).$$

Note that in the case $G = SO(2, F)$, the induction does nothing, since

$$M_{(0)} = M_{(1)} = SO(2, F) \cong GL(1, F),$$

and for a smooth representation π of $GL(1, F)$, we have

$$\pi \rtimes 1 = \pi.$$

It follows from [BZ], Prop.2.3, that for smooth representations π_1 of $GL(n_1, F)$, π_2 of $GL(n_2, F)$ and σ of $SO(2m, F)$ we have

$$\pi_1 \rtimes (\pi_2 \rtimes \sigma) \cong (\pi_1 \times \pi_2) \rtimes \sigma.$$

Let σ be a finite length smooth representation of $SO(2n, F)$. Let $\alpha = (n_1, \dots, n_k)$ be an ordered partition of a non-negative integer $m \leq n$. Define

$$s_\alpha(\sigma) = r_{M_\alpha, SO(2n, F)}(\sigma).$$

4. SOME PROPERTIES OF PARABOLICALLY INDUCED REPRESENTATIONS OF $SO(2n, F)$

Let $G = SO(2n, F)$. For $m < n$, let $P = P_{(m)}$ be the standard parabolic subgroup with Levi factor $M \cong GL(m, F) \times SO(2(n - m), F)$. Then

$$s(P) = P, \quad s(M) = M, \quad s(U) = U.$$

The following lemma can be proved directly:

LEMMA 4.1. *For a smooth finite length representation π of $GL(m, F)$ and a smooth finite length representation σ of $SO(2(n - m), F)$, $m < n$, we have*

$$s(\pi \rtimes \sigma) \cong \pi \rtimes s(\sigma).$$

PROPOSITION 4.2. *Let π be a smooth finite length representation of $GL(m, F)$ and σ be a smooth finite length representation of $SO(2(n - m), F)$. Then*

$$\tilde{\pi} \rtimes \sigma = s^m(\pi \rtimes \sigma).$$

Particularly,

1. *If m is even, then*

$$\tilde{\pi} \rtimes \sigma = \pi \rtimes \sigma;$$

2. *If $m < n$ is odd, then*

$$\tilde{\pi} \rtimes \sigma = \pi \rtimes s(\sigma);$$

3. *If $m = n$ is odd, then*

$$\tilde{\pi} \rtimes 1 = s(\pi \rtimes 1).$$

(Here $\tilde{\pi}$ denotes contragredient representation of π .)

Define

$$\beta_i = (\underbrace{1, \dots, 1}_{i \text{ times}}, 0, \dots, 0) \in \mathbb{R}^n, \quad i \leq n - 2,$$

$$\beta_{n-1} = (1, \dots, 1, -1) \in \mathbb{R}^n,$$

$$\beta_n = (1, \dots, 1, 1) \in \mathbb{R}^n.$$

Let π be an irreducible smooth representation of $G = SO(2n, F)$. Let $P = MU$ be a standard parabolic subgroup, minimal among all standard parabolic subgroups which satisfy

$$\tau_{M,G}(\pi) \neq 0.$$

Let ρ be an irreducible subquotient of $\tau_{M,G}(\pi)$.

If $P = P_\alpha$, where $\alpha = (n_1, \dots, n_k)$ is a partition of $m \leq n$, then

$$\rho = \rho_1 \otimes \dots \otimes \rho_k \otimes \sigma,$$

where ρ_i are irreducible cuspidal representations of $GL(n_i, F)$, and σ is an irreducible cuspidal representation of $SO(2(n - m), F)$. If P is not of that type, then

$$\rho = s(\rho_1 \otimes \dots \otimes \rho_{k-1} \otimes \rho_k \otimes 1) = \rho_1 \otimes \dots \otimes \rho_{k-1} \otimes s(\rho_k \otimes 1),$$

where ρ_i are irreducible cuspidal representations of $GL(n_i, F)$.

We have $\rho_i = \nu^{e(\rho_i)} \rho_i^u$, where $e(\rho_i) \in \mathbb{R}$ and ρ_i^u is unitarizable. Define

$$e_*(\rho) = (\underbrace{e(\rho_1), \dots, e(\rho_1)}_{n_1 \text{ times}}, \dots, \underbrace{e(\rho_k), \dots, e(\rho_k)}_{n_k \text{ times}}, \underbrace{0, \dots, 0}_{n-m \text{ times}}).$$

(This definition concerns $\rho = \rho_1 \otimes \dots \otimes \rho_k \otimes \sigma$ as well as $\rho = s(\rho_1 \otimes \dots \otimes \rho_k \otimes 1)$.)

If π is square integrable, then

$$\begin{aligned} (e_*(\rho), \beta_{n_1}) &> 0, \\ (e_*(\rho), \beta_{n_1+n_2}) &> 0, \\ &\vdots \\ (e_*(\rho), \beta_{m-n_k}) &> 0, \\ (e_*(\rho), \beta_m) &> 0. \end{aligned}$$

(Here $(,)$ denotes the standard inner product on \mathbb{R}^n .)

Conversely, if all above inequalities hold for any α and σ as above, then π is square integrable.

The criteria implies

$$\pi \text{ is square integrable} \Leftrightarrow s(\pi) \text{ is square integrable,}$$

but this equivalence can also be proved easily directly from the definition of square integrability.

6. A THEOREM ON SELF-DUALITY

THEOREM 6.1. *Suppose that $\rho_1, \rho_2, \dots, \rho_k$ are irreducible cuspidal representations of $GL(n_1, F), \dots, GL(n_k, F)$, resp., and σ is an irreducible cuspidal representation of $SO(2l, F)$, $l \neq 1$. If $\rho_1 \times \dots \times \rho_k \rtimes \sigma$ contains a square integrable subquotient, then $\rho_i^u \cong (\rho_i^u)^\sim$, for any $i = 1, 2, \dots, k$.*

PROOF. The proof parallels that used in chapter 4 of [T3].

Set $n_1 + \dots + n_k = m$, $m + l = n$. Denote

$$\begin{aligned} G &= SO(2n, F), \\ M &= M_{(n_1, \dots, n_k)}, \\ \rho &= \rho_1 \otimes \dots \otimes \rho_k \otimes \sigma. \end{aligned}$$

Then

$$\rho_1 \times \dots \times \rho_k \rtimes \sigma = i_{G, M}(\rho).$$

Let π be an irreducible square integrable subquotient of $i_{G, M}(\rho)$. First we shall prove the lemma under the assumption that π is a subrepresentation of $i_{G, M}(\rho)$, or, equivalently, that ρ is a quotient of $r_{M, G}(\pi)$.

Fix any $i_0 \in \{1, \dots, k\}$. Set

$$\begin{aligned} Y_{i_0}^0 &= \{i \in \{1, \dots, k\} \mid \exists \alpha \in \mathbb{Z} \text{ such that } \rho_i \cong \nu^\alpha \rho_{i_0}\}, \\ Y_{i_0}^1 &= \{i \in \{1, \dots, k\} \mid \exists \alpha \in \mathbb{Z} \text{ such that } \tilde{\rho}_i \cong \nu^\alpha \rho_{i_0}\}, \\ Y_{i_0} &= Y_{i_0}^0 \cup Y_{i_0}^1, \\ Y_{i_0}^c &= \{1, \dots, k\} \setminus Y_{i_0}. \end{aligned}$$

Suppose that $\rho_{i_0}^u \not\cong (\rho_{i_0}^u)^\sim$. It follows from Proposition 2.1 that for any $j_0, j'_0 \in Y_{i_0}^0, j_1, j'_1 \in Y_{i_0}^1$ and $j_c \in Y_{i_0}^c$ we have

$$\begin{aligned} \rho_{j_0} \times \tilde{\rho}_{j'_0} &\cong \tilde{\rho}_{j'_0} \times \rho_{j_0}, & \rho_{j_1} \times \tilde{\rho}_{j'_1} &\cong \tilde{\rho}_{j'_1} \times \rho_{j_1}, \\ \rho_{j_0} \times \rho_{j_1} &\cong \rho_{j_1} \times \rho_{j_0}, & \tilde{\rho}_{j_0} \times \tilde{\rho}_{j_1} &\cong \tilde{\rho}_{j_1} \times \tilde{\rho}_{j_0}, \\ \rho_{j_0} \times \rho_{j_c} &\cong \rho_{j_c} \times \rho_{j_0}, & \tilde{\rho}_{j_0} \times \rho_{j_c} &\cong \rho_{j_c} \times \tilde{\rho}_{j_0}, \\ \rho_{j_1} \times \rho_{j_c} &\cong \rho_{j_c} \times \rho_{j_1}, & \tilde{\rho}_{j_1} \times \rho_{j_c} &\cong \rho_{j_c} \times \tilde{\rho}_{j_1}. \end{aligned}$$

If $n - m > 1$, then, by Lemma 4.3, $\rho_{j_0} \rtimes \sigma$ and $\rho_{j_1} \rtimes \sigma$ are irreducible. Now we get from Proposition 4.2

$$\begin{aligned} \rho_{j_0} \rtimes \sigma &\cong \tilde{\rho}_{j_0} \rtimes s^{n_{j_0}}(\sigma), \\ \rho_{j_1} \rtimes \sigma &\cong \tilde{\rho}_{j_1} \rtimes s^{n_{j_1}}(\sigma), \end{aligned}$$

for $n - m > 1$, and

$$\begin{aligned} \rho_{j_0} \rtimes 1 &\cong s^{n_{j_0}}(\tilde{\rho}_{j_0} \rtimes 1), \\ \rho_{j_1} \rtimes 1 &\cong s^{n_{j_1}}(\tilde{\rho}_{j_1} \rtimes 1), \end{aligned}$$

for $n = m$. Write

$$\begin{aligned} Y_{i_0}^0 &= \{a_1, \dots, a_{k_0}\}, & a_i < a_j \text{ for } i < j, \\ Y_{i_0}^1 &= \{b_1, \dots, b_{k_1}\}, & b_i < b_j \text{ for } i < j, \\ Y_{i_0}^c &= \{d_1, \dots, d_{k_c}\}, & d_i < d_j \text{ for } i < j. \end{aligned}$$

If $n - m \geq 2$, then we can repeat the proof from [T3], since we have just slightly different relations, and square integrability criteria are the same.

Let $m = n$. Set $\alpha = n_{\beta_{k_1}}$. Then

$$\begin{aligned} &\rho_1 \times \cdots \times \rho_k \rtimes 1 \cong \\ &\cong \rho_{a_1} \times \cdots \times \rho_{a_{k_0}} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \times \rho_{b_1} \times \cdots \times \rho_{b_{k_1}} \rtimes 1 \\ &\cong \rho_{a_1} \times \cdots \times \rho_{a_{k_0}} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \times \rho_{b_1} \times \cdots \times \rho_{b_{k_1-1}} \times s^\alpha(\bar{\rho}_{b_{k_1}} \rtimes 1) \\ &\cong s^\alpha(\rho_{a_1} \times \cdots \times \rho_{a_{k_0}} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \times \rho_{b_1} \times \cdots \times \rho_{b_{k_1-1}} \times \bar{\rho}_{b_{k_1}} \rtimes 1) \\ &\cong s^\alpha(\rho_{a_1} \times \cdots \times \rho_{a_{k_0}} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \times \bar{\rho}_{b_{k_1}} \times \rho_{b_1} \times \cdots \times \rho_{b_{k_1-1}} \rtimes 1). \end{aligned}$$

We proceed in the same way, and finally we get

$$\begin{aligned} &\rho_1 \times \cdots \times \rho_k \rtimes 1 \cong s^\gamma(\rho_{a_1} \times \cdots \\ &\quad \times \rho_{a_{k_0}} \times \bar{\rho}_{b_{k_1}} \times \cdots \times \bar{\rho}_{b_1} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \rtimes 1), \end{aligned}$$

where $\gamma = 0$ or 1 .

In the same manner, we obtain

$$\begin{aligned} &\rho_1 \times \cdots \times \rho_k \rtimes 1 \cong s^\delta(\rho_{b_1} \times \cdots \times \rho_{b_{k_1}} \times \bar{\rho}_{a_{k_0}} \times \cdots \\ &\quad \times \bar{\rho}_{a_1} \times \rho_{d_1} \times \cdots \times \rho_{d_{k_c}} \rtimes 1), \end{aligned}$$

where $\delta = 0$ or 1 .

By the Frobenius reciprocity, the representations

$$\begin{aligned} \rho' &= s^\gamma(\rho_{a_1} \otimes \cdots \otimes \rho_{a_{k_0}} \otimes \bar{\rho}_{b_{k_1}} \otimes \cdots \otimes \bar{\rho}_{b_1} \otimes \rho_{d_1} \otimes \cdots \otimes \rho_{d_{k_c}} \otimes 1), \\ \rho'' &= s^\delta(\rho_{b_1} \otimes \cdots \otimes \rho_{b_{k_1}} \otimes \bar{\rho}_{a_{k_0}} \otimes \cdots \otimes \bar{\rho}_{a_1} \otimes \rho_{d_1} \otimes \cdots \otimes \rho_{d_{k_c}} \otimes 1) \end{aligned}$$

are the quotients of corresponding Jacquet modules. Now $\rho_{a_1} \times \cdots \times \rho_{a_{k_0}} \times \rho_{b_1} \times \cdots \times \rho_{b_{k_1}}$ is representation of $GL(u, F)$, for some $u \leq n$. If $u \neq n - 1$ then $(\beta_u, e_*(\rho')) = -(\beta_u, e_*(\rho''))$. If $u = n - 1$, then

$$(\beta_{n-1}, e_*(\rho')) + (\beta_n, e_*(\rho')) = -(\beta_{n-1}, e_*(\rho'')) - (\beta_n, e_*(\rho'')).$$

Anyway, this contradicts the assumption that π is square integrable.

Generally, let π be an irreducible subquotient of $i_{G,M}(\rho)$. By [C], Corollary 7.2.2, there exists $w \in W = N_G(M)/M$ such that π is a subrepresentation of $i_{G,M}(w(\rho))$. Let $\rho = \rho_1 \otimes \cdots \otimes \rho_k \otimes \sigma$ and $w(\rho) = \delta_1 \otimes \cdots \otimes \delta_k \otimes \tau$. We apply the first part of the proof on $w(\rho)$, and we get $\delta_i^u \cong (\delta_i^u)^\sim$, $i = 1, 2, \dots, k$. By [G], the sequence $\delta_1, \dots, \delta_k$ is, up to a permutation and taking a contragredient, the sequence ρ_1, \dots, ρ_k . □

THEOREM 6.2. *Suppose that $\rho_1, \rho_2, \dots, \rho_k$ are irreducible cuspidal representations of $GL(n_1, F), \dots, GL(n_k, F)$, resp., and σ is an irreducible cuspidal representation of $SO(2l, F)$, $l \neq 1$, such that $\rho_1 \times \dots \times \rho_k \rtimes \sigma$ contains a square integrable subquotient. Further, assume that for each unitary representation ρ , the number α , discussed in Lemma 4.3., satisfies $2\alpha \in \mathbb{Z}$. Then $2e(\rho_i) \in \mathbb{Z}$, for any $i = 1, 2, \dots, k$.*

PROOF. The proof is analogous to that of Theorem 6.1. \square

REFERENCES

- [1] I.N.Bernstein and A.V.Zelevinsky, Induced representations of reductive p-adic groups, I, Ann. Sci. École Norm. Sup. **10** (1977), 441-472.
- [2] J.Bernstein, P.Deligne, and D.Kazhdan, Trace Paley-Wiener theorem for reductive p-adic groups, J.Analyse Math.**42** (1986), 180-192.
- [3] W.Casselman, Introduction to the theory of admissible representations of p-adic reductive groups, preprint
- [4] D.K.Faddeev, On multiplication of representations of classical groups over finite field with representations of the full linear group, Vest. Leningrad. Univ. **13** (1976), 35-40.[in Russian]
- [5] D.Goldberg, Reducibility of induced representations for $Sp(2n)$ and $SO(n)$, Amer. J. Math. vol(116), 1994, 1101-1151.
- [6] M.E.Sweedler, "Hopf Algebras", Benjamin, New York, 1969.
- [7] M.Tadić, Structure arising from induction and Jacquet modules of representations of classical p-adic groups, J. of Algebra **177** (1995), 1-33.
- [8] M.Tadić, Representations of p-adic symplectic groups, Compositio Math. **90** (1994) 123-181
- [9] M.Tadić, On regular square integrable representations of p-adic groups, American Journal of Mathematics **120** (1998), 159-210.
- [10] A.V.Zelevinsky, Induced representations of reductive p-adic groups, II, On irreducible representations of $GL(n)$, Ann. Sci. École Norm. Sup. **13** (1980), 165-210.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SPLIT, TESLINA 12, 21 000 SPLIT, CROATIA

Current address: Dept. of Math., Purdue Univ., West Lafayette, IN 47907, USA

E-mail address: dban@mapmf.pmfst.hr; dban@math.purdue.edu

Received: 21.05.98.

Revised: 09.11.98