

# Euklid's plane through Symmetry

Wolfgang Soergel

Mathematisches Institut  
Universität Freiburg

October 2022

- ▶ **Incidence geometry:** Pair  $(X, G)$  with  $X$  a set of “points”,  $G \subset \mathcal{P}(X)$  a set of “lines”, each line has at least two points, through every two distinct points there goes exactly one line.
- ▶ **Betweenness:** Subset  $Z \subset X^3$  of collinear tripels giving two opposite orders on every line, such that a line never meets only one segment of a triangle.
- ▶ **Congruence group:** A subgroup  $K \subset \text{Aut}(X, G, Z)$  such that for any two halflines  $A, B \subset X$  there exist exactly two  $k, h \in K$  with  $k(A) = B = h(A)$ .
- ▶ **Supremum property:** With respect to a  $Z$ -order every nonempty bounded above subset on a line has a supremum.
- ▶ **Parallel Axiom:**  $\forall g \in G, p \in X \setminus g$  there exists uniquely  $h \in G$  with  $p \in h$  and  $h \cap g = \emptyset$ .

- ▶ **Theorem:** There is up to isomorphism a unique quadrupel  $(X, G, Z, K)$  of an incidence geometry with betweenness relation and congruence group that satisfies supremum and parallel axioms and has at least one line. [Soergel: Elementargeometrie]
- ▶ A **Congruence group**  $K \subset \text{Aut}(X, G, Z)$  is a subgroup such that for any two halflines  $A, B \subset X$  exist exactly two  $k, h \in K$  with  $k(A) = B = h(A)$ .
- ▶ If we ask instead congruences to act free and transitive on the set of halflines, there are is a “bad” model for every nontrivial group homomorphism  $\text{SO}(2) \rightarrow \mathbb{R}_{>0}$ .
- ▶ If we ask the parallel axiom to be false, there should be also a unique model. I would like an easy proof.

Now I am going make lots of claims and if you don't believe one of them, you are welcome to speak up and I will try to explain the proof on the blackboard.

Let  $(X, G, Z)$  be an incidence geometry with betweenness.

- ▶ A line meeting no vertex of a triangle meets exactly to segments or none.
- ▶ The complement of a line is the disjoint union of at most two equivalence classes under the relation “joinable by a segment”.

Let  $(X, G, Z, K)$  be an incidence geometry with betweenness and congruences.

- ▶ Every halfline is infinite.
- ▶ For every line  $g$  there is a unique nontrivial congruence  $s_g$  fixing it pointwise, the **reflection along  $g$** .
- ▶ Every segment is infinite.
- ▶ For every line  $g$  there are exactly two halfspaces. They are exchanged by the reflection  $s_g$ .

Let  $(X, G, Z, K)$  be an incidence geometry with betweenness and congruences.

- ▶ Let  $h \perp g$  mean  $g \neq h = s_g(h)$ .
- ▶ For every line  $g$  and every point  $x$  there is a unique perpendicular  $h \perp g$  with  $x \in h$ .
- ▶  $h \perp g \Leftrightarrow s_h s_g = s_g s_h$
- ▶ Two perpendiculars to a line  $g$  are disjoint.
- ▶ Under the parallel axiom perpendiculars to a line are perpendicular to its parallels.

Let  $(X, G, Z, K)$  be incidence geometry with betweenness and congruences and let  $g \in G$  a line.

- ▶ Denote by  $K|_g \subset K$  the stabilizer of a line and its halfspaces.
- ▶ Denote by  $\vec{g} \subset K|_g$  the subgroup stabilizing both  $Z$ -orders on the line  $g$ .
- ▶  $\vec{g}$  acts free and transitive on  $g$ .



Let  $(X, G, Z, K)$  be an incidence geometry with betweenness and congruences and supremum axiom and let  $g \in G$  be a line.

- ▶ Given  $v \in \vec{g} \setminus e_K$  we have  $v(x) > x$  for all  $x$  and some  $Z$ -order on  $g$ .
- ▶ All elements of  $K|_g \setminus \vec{g}$  are involutions.
- ▶ Any two different points  $x \neq y$  can be exchanged by a unique reflection. It stabilizes the  $\overline{xy}$ -halfplanes.

Let  $(X, G, Z, K)$  be an incidence geometry with betweenness and congruences and supremum axiom and let  $g \in G$  be a line.

- ▶ All elements of  $K|_g \setminus \vec{g}$  are reflections. These elements generate  $K|_g$ .
- ▶ Every element  $v \in \vec{g}$  has a square root. Conjugating  $v$  by an element of  $K|_g \setminus \vec{g}$  we get its inverse.
- ▶  $\vec{g}$  is commutative.

Let  $(X, G, Z, K)$  be an incidence geometry with betweenness and congruences and supremum and parallels axiom.

- ▶  $g \parallel h \Rightarrow \vec{g} = \vec{h}$
- ▶ Assume there exists a line. Then all translations form a commutative subgroup  $\vec{X} := \bigcup_{g \in G} \vec{g} \subset K$  acting free and transitive on  $X$ .

- ▶ Let  $A$  be an ordered group such that no nontrivial cyclic subgroup has an upper bound in  $A$  and every element has a root. Then for every  $a > e_A$  there exists a unique order preserving group homomorphism  $A \rightarrow (\mathbb{R}, +)$  with  $a \mapsto 1$  and it is injective.

Let  $(X, G, Z, K)$  be incidence geometry with betweenness and congruences and supremum axiom.

- ▶ For every line  $g$ , there is a unique structure on  $\vec{g}$  as a real vector space compatible with  $Z$ .
- ▶ If the parallel axiom holds and there is a least one line, there is a unique structure of real vector space on  $\vec{X}$  such that  $\vec{g}$  is a subspace for all  $g \in G$ .  
Furthermore the space  $\vec{X}$  has dimension two, so that  $X$  acquires the structure of a twodimensional real affine space.

Let  $(X, G, Z, K)$  be incidence geometry with betweenness and congruences and supremum and parallel axioms and a line.

- ▶ The congruence group  $K$  consists of affinities and contains all translations.
- ▶ The isotropy groups  $K_x$  all have the same image  $D \subset GL(\vec{X})$ .
- ▶ The subgroup  $D \subset GL(\vec{X})$  has the property that for any two rays  $A, B \subset \vec{X}$  there are exactly two elements  $r, s \in D$  with  $r(A) = B = s(A)$ .

Let  $V$  be a twodimensional real vector space. Let  $D \subset GL(V)$  a subgroup such that for any two rays  $A, B \subset V$  there are exactly two elements  $r, s \in D$  with  $r(A) = B = s(A)$ .

- ▶ There exists a  $D$ -invariant scalar product on  $V$ .
- ▶ Any two  $D$ -invariant scalar products are scalar multiples of one another.
- ▶ The group  $D$  is the orthogonal group for one and any of these scalar products.

This proves that an incidence geometry with betweenness and congruences  $(X, G, Z, K)$  satisfying supremum and parallel axioms and having a line is isomorphic to

$$(\mathbb{R}^2, \text{affine lines, obvious } Z, O(2)_{\text{aff}})$$



Thanks!