



Glasnik
Matematički

SERIJA III

www.math.hr/glasnik

Takao Ohno and Tetsu Shimomura

*Continuity of generalized Riesz potentials for double phase functionals with
variable exponent*

Accepted manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by Glasnik Production staff.

CONTINUITY OF GENERALIZED RIESZ POTENTIALS FOR DOUBLE PHASE FUNCTIONALS WITH VARIABLE EXPONENT

TAKAO OHNO AND TETSU SHIMOMURA

Oita University and Hiroshima University, Japan

ABSTRACT. In this note, we discuss the continuity of generalized Riesz potentials $I_\rho f$ of functions in Morrey spaces $L^{\Phi, \nu(\cdot)}(G)$ of double phase functionals with variable exponents.

1. INTRODUCTION

The double phase functional introduced by Zhikov ([30]) in the 1980s has been studied intensively by many mathematicians. Regarding regularity theory of differential equations, Baroni, Colombo and Mingione [1, 4, 5] studied a double phase functional

$$\tilde{\Phi}(x, t) = t^p + a(x)t^q, \quad x \in \mathbf{R}^N, \quad t \geq 0$$

where $N \geq 2$, $1 \leq p < q$, $a(\cdot)$ is non-negative, bounded and Hölder continuous of order $\theta \in (0, 1]$. We refer to [10, 16] for Sobolev's inequality, [11] for Trudinger's inequality and e.g. [2, 7, 8] for other double phase problems.

For $0 < \alpha < N$ and a locally integrable function f on \mathbf{R}^N the Riesz potential $I_\alpha f$ of order α is defined by

$$I_\alpha f(x) = \int_{\mathbf{R}^N} |x - y|^{\alpha - N} f(y) dy.$$

In [12] we discussed the continuity of Riesz potentials $I_\alpha f$ of functions in Morrey spaces $L^{\tilde{\Phi}, \nu}(\mathbf{R}^N)$ of the double phase functionals $\tilde{\Phi}(x, t)$. We refer to [16, Section 5] for the $L^{\tilde{\Phi}}$ case and [14] for the $L^{p(\cdot), \nu(\cdot)}$ case.

2000 *Mathematics Subject Classification.* 31B15, 46E35.

Key words and phrases. Riesz potentials, Morrey spaces, double phase functionals, continuity.

In the present note, we consider the case $\Phi(x, t)$ is a double phase functional given by

$$\Phi(x, t) = t^{p(x)} + (b(x)t)^{q(x)},$$

where $p(x) < q(x)$ and $b(\cdot)$ is non-negative, bounded and Hölder continuous of order $\theta \in (0, 1]$ ([10], cf. [3, 26]).

To obtain general results, we consider the family (ρ) of all functions ρ satisfying the following conditions: $\rho : (0, \infty) \rightarrow (0, \infty)$ is a measurable function such that

$$\int_0^r \rho(s) \frac{ds}{s} < \infty$$

for all sufficiently small $r > 0$ and there exists constants $0 < k < 1$, $0 < k_1 < k_2$ and $C_\rho > 0$ such that

$$(1.1) \quad \sup_{kr \leq s \leq r} \rho(s) \leq C_\rho \int_{k_1 r}^{k_2 r} \rho(s) \frac{ds}{s}$$

for all $r > 0$ (e.g. [6, 27]). We do not postulate the doubling condition on ρ .

EXAMPLE 1.1. If ρ satisfies the doubling condition, that is, there exists a constant $C > 0$ such that $C^{-1} \leq \rho(r)/\rho(s) \leq C$ for $1/2 \leq r/s \leq 2$, then ρ satisfies (1.1) whenever $k = 1/2$ and $2k_1 = k_2$. If ρ is increasing, then ρ satisfies (1.1) with $k = 1/2$, $k_1 = 1$ and $k_2 = 2$. If $\alpha \in \mathbf{R}$ such that $\rho(r) = r^\alpha e^{-1/r}$, then ρ satisfies (1.1) with $k = 1/2$, $k_1 = 1/4$ and $k_2 = 1/2$. See also [21, Lemma 2.5], [24, 27] and [28, Remark 2.2].

Let G be an open bounded set in \mathbf{R}^N . For a function $\rho \in (\rho)$, we define the generalized Riesz potential $I_\rho f$ of f by

$$I_\rho f(x) = \int_G \frac{\rho(|x-y|)f(y)}{|x-y|^N} dy,$$

where $f \in L^1(G)$. We write $I_\rho f = I_\alpha f$ when $\rho(r) = r^\alpha$, $0 < \alpha < N$. We refer to [15, 22, 23, 25, 29] etc. for the study of $I_\rho f$.

Our aim in this note is to study the continuity of generalized Riesz potential $I_\rho f$ of functions f in Morrey spaces $L^{\Phi, \nu(\cdot)}(G)$ of the double phase functionals with variable exponents (Theorem 2.2), as an extension of [12, Theorem 4.1].

2. DEFINITIONS AND THE MAIN THEOREM

Throughout this paper, let C denote various constants independent of the variables in question and \log be a natural logarithm.

Let $p(\cdot)$ be a measurable functions on G such that

$$(P1) \quad 1 \leq p^- := \inf_{x \in G} p(x) \leq \sup_{x \in G} p(x) =: p^+ < \infty,$$

(P2) $p(\cdot)$ is log-Hölder continuous on G , namely

$$|p(x) - p(y)| \leq \frac{C_p}{\log(e + 1/|x - y|)} \quad (x, y \in G)$$

with a constant $C_p \geq 0$.

Let $\nu(\cdot)$ be a measurable functions on G such that

$$0 < \nu^- := \inf_{x \in G} \nu(x) \leq \sup_{x \in G} \nu(x) =: \nu^+ < \infty.$$

Let $B(x, r)$ denote the open ball centered at $x \in \mathbf{R}^N$ with radius $r > 0$. For a set $E \subset \mathbf{R}^N$, $|E|$ denotes the Lebesgue measure of E . Set $d_G = \sup\{|x - y| : x, y \in G\}$. Morrey space with variable exponents $L^{p(\cdot), \nu(\cdot)}(G)$ is the family of measurable functions f on G satisfying

$$L^{p(\cdot), \nu(\cdot)}(G) = \left\{ f \in L^1_{loc}(G); \right. \\ \left. \sup_{x \in G, 0 < r < d_G} \frac{r^{\nu(x)}}{|B(x, r)|} \int_{G \cap B(x, r)} |f(y)|^{p(y)} dy < \infty \right\}.$$

It is a Banach space with respect to the norm

$$\|f\|_{L^{p(\cdot), \nu(\cdot)}(G)} = \inf \left\{ \lambda > 0; \right. \\ \left. \sup_{x \in G, 0 < r < d_G} \frac{r^{\nu(x)}}{|B(x, r)|} \int_{G \cap B(x, r)} \left(\frac{|f(y)|}{\lambda} \right)^{p(y)} dy \leq 1 \right\}$$

(cf. see [19]).

We consider a function $\Phi(x, t) : G \times [0, \infty) \rightarrow [0, \infty)$ satisfying the following conditions ($\Phi 1$) and ($\Phi 2$):

- ($\Phi 1$) $\Phi(\cdot, t)$ is measurable on G for each $t \geq 0$ and $\Phi(x, \cdot)$ is convex on $[0, \infty)$ for each $x \in G$;
- ($\Phi 2$) there exists a constant $A_1 \geq 1$ such that $A_1^{-1} \leq \Phi(x, 1) \leq A_1$ for all $x \in G$.

The Musielak-Orlicz-Morrey space $L^{\Phi, \nu(\cdot)}(G)$ is defined by

$$L^{\Phi, \nu(\cdot)}(G) \\ = \left\{ f \in L^1_{loc}(G) : \right. \\ \left. \sup_{x \in G, 0 < r < d_G} \frac{r^{\nu(x)}}{|B(x, r)|} \int_{G \cap B(x, r)} \Phi \left(y, \frac{|f(y)|}{\lambda} \right) dy < \infty \text{ for some } \lambda > 0 \right\}.$$

It is a Banach space with respect to the norm

$$\|f\|_{L^{\Phi, \nu(\cdot)}(G)} = \inf \left\{ \lambda > 0 : \sup_{x \in G, 0 < r < d_G} \frac{r^{\nu(x)}}{|B(x, r)|} \int_{G \cap B(x, r)} \Phi \left(y, \frac{|f(y)|}{\lambda} \right) dy \leq 1 \right\}$$

(see [9, 20]).

Let $q(\cdot)$ be a measurable function on G such that

$$(Q1) \quad 1 \leq q^- := \inf_{x \in G} q(x) \leq \sup_{x \in G} q(x) =: q^+ < \infty,$$

(Q2) $q(\cdot)$ is log-Hölder continuous on G , namely

$$|q(x) - q(y)| \leq \frac{C_q}{\log(e + 1/|x - y|)} \quad (x, y \in G)$$

with a constant $C_q \geq 0$.

In what follows, set

$$\Phi(x, t) = t^{p(x)} + (b(x)t)^{q(x)},$$

where $p(x) < q(x)$ and $b(\cdot)$ is non-negative, bounded and Hölder continuous of order $\theta \in (0, 1]$ ([10], cf. [3, 26]).

REMARK 2.1. Let $f \in L^{\Phi, \nu(\cdot)}(G)$ be a measurable function on G . Then note that $f \in L^{p(\cdot), \nu(\cdot)}(G)$ and $bf \in L^{q(\cdot), \nu(\cdot)}(G)$.

We state the following, as an extension of [12, Theorem 4.1].

THEOREM 2.2. *Let $\rho \in (\rho)$. Assume that there are constants $\eta_1 > 0, \eta_2 > 0, \tau > 0$ and $C_0 > 0$ such that*

$$(2.2) \quad \left| \frac{\rho(|x - y|)}{|x - y|^N} - \frac{\rho(|z - y|)}{|z - y|^N} \right| \leq C_0 \frac{|x - z|^{\eta_1} \rho(\tau|x - y|)}{|x - y|^{\eta_2} |x - y|^N}$$

whenever $x, y, z \in G$ and $|x - z| \leq |x - y|/2$. Abbreviate

$$\begin{aligned} & \psi(x, z, r) \\ & \equiv \int_0^{4k_2 r} s^{-\nu(x)/p(x)+\theta} \rho(s) \frac{ds}{s} + \int_0^{4k_2 r} s^{-\nu(x)/q(x)} \rho(s) \frac{ds}{s} \\ & + \int_0^{6k_2 r} s^{-\nu(z)/p(z)+\theta} \rho(s) \frac{ds}{s} + \int_0^{6k_2 r} s^{-\nu(z)/q(z)} \rho(s) \frac{ds}{s} \\ & + r^\theta \int_{k_1 r}^{4k_2 d_G} s^{-\nu(z)/p(z)} \rho(s) \frac{ds}{s} \\ & + r^{\eta_1} \int_{2k_1 \tau r}^{4k_2 \tau d_G} s^{-\nu(x)/p(x)-\eta_2+\theta} \rho(s) \frac{ds}{s} + r^{\eta_1} \int_{2k_1 \tau r}^{4k_2 \tau d_G} s^{-\nu(x)/q(x)-\eta_2} \rho(s) \frac{ds}{s} \end{aligned}$$

for $x, z \in G$ and $0 < r \leq d_G$, where k_1 and k_2 are constants in (ρ) . Assume that $\psi(x, z, r) < \infty$ for all $x, z \in G$ and $0 < r \leq d_G$. Then there exists a constant $C > 0$ such that

$$|b(x)I_\rho f(x) - b(z)I_\rho f(z)| \leq C\psi(x, z, |x - z|)$$

for all $x, z \in G$ and measurable functions f on G with $\|f\|_{L^{\Phi, \nu(\cdot)}(G)} \leq 1$.

REMARK 2.3. Let $\rho(r) = r^\alpha e^{-1/r}$ be as in Example 1.1. Then the mean value property implies that (2.2) holds for $\eta_1 = 1, \eta_2 = 2$ and $\tau = 3/2$. Note here that there exists a constant $C \geq 1$ such that

$$C^{-1} \frac{\rho(r)}{r^{N+2}} \leq \frac{d}{dr} \left(\frac{\rho(r)}{r^N} \right) \leq C \frac{\rho(r)}{r^{N+2}}$$

for all $0 < r < d_G$ and $|x - y|/2 \leq |x - y + t(z - x)| \leq 3|x - y|/2$ for $0 \leq t \leq 1$ and $|x - z| \leq |x - y|/2$.

REMARK 2.4. If $\rho(r)r^a$ is increasing for some $a \geq 0$ and $\rho(r)/r^b$ is decreasing for some $b \geq 0$, then ρ satisfies the doubling condition and

$$\left| \frac{\rho(r)}{r^N} - \frac{\rho(s)}{s^N} \right| \leq C_0 |r - s| \frac{\rho(r)}{r^{N+1}}, \quad \text{for } \frac{1}{2} \leq \frac{s}{r} \leq 2.$$

See [22].

3. COROLLARIES

In this section, we give consequences of our theorem.

COROLLARY 3.1. Let $\rho(r) = r^\alpha (\log(e + 1/r))^\beta$ for $\alpha > 0$ and $\beta \in \mathbf{R}$. Suppose $\inf_{x \in G} (\nu(x) - \alpha p(x)) > 0$ and $\inf_{x \in G} ((\alpha + \theta)p(x) - \nu(x)) > 0$. Further suppose $\inf_{x \in G} (\nu(x) - (\alpha - 1)q(x)) > 0$ and $\inf_{x \in G} (\alpha q(x) - \nu(x)) > 0$. Then there exists a constant $C > 0$ such that

$$\begin{aligned} & |b(x)I_\rho f(x) - b(z)I_\rho f(z)| \\ & \leq C \{ |x - z|^{\alpha - \nu(x)/p(x) + \theta} + |x - z|^{\alpha - \nu(x)/q(x)} \\ & \quad + |x - z|^{\alpha - \nu(z)/p(z) + \theta} + |x - z|^{\alpha - \nu(z)/q(z)} \} (\log(e + 1/|x - z|))^\beta \end{aligned}$$

for all $x, z \in G$ and measurable functions f on G with $\|f\|_{L^{\Phi, \nu(\cdot)}(G)} \leq 1$.

PROOF. Since $\inf_{x \in G} ((\alpha + \theta)p(x) - \nu(x)) > 0$, taking ε_1 such that $0 < \varepsilon_1 < \alpha - \nu(x)/p(x) + \theta$, there exists a constant $c_1 > 0$ such that

$$s_1^{\alpha - \nu(x)/p(x) + \theta - \varepsilon_1} (\log(e + 1/s_1))^\beta \leq c_1 s_2^{\alpha - \nu(x)/p(x) + \theta - \varepsilon_1} (\log(e + 1/s_2))^\beta$$

whenever $0 < s_1 \leq s_2$ (see e.g. [17, 18]). Therefore we have

$$\begin{aligned} & \int_0^{4k_2r} s^{\alpha-\nu(x)/p(x)+\theta} (\log(e+1/s))^\beta \frac{ds}{s} \\ & \leq c_1 (4k_2r)^{\alpha-\nu(x)/p(x)+\theta-\varepsilon_1} (\log(e+1/(4k_2r)))^\beta \int_0^{4k_2r} s^{\varepsilon_1} \frac{ds}{s} \\ & \leq Cr^{\alpha-\nu(x)/p(x)+\theta} (\log(e+1/r))^\beta \end{aligned}$$

and, similarly we obtain

$$\int_0^{6k_2r} s^{\alpha-\nu(z)/p(z)+\theta} (\log(e+1/s))^\beta \frac{ds}{s} \leq Cr^{\alpha-\nu(z)/p(z)+\theta} (\log(e+1/r))^\beta.$$

Since $\inf_{x \in G} (\alpha q(x) - \nu(x)) > 0$, we obtain

$$\int_0^{4k_2r} s^{\alpha-\nu(x)/q(x)} (\log(e+1/s))^\beta \frac{ds}{s} \leq Cr^{\alpha-\nu(x)/q(x)} (\log(e+1/r))^\beta$$

and

$$\int_0^{6k_2r} s^{\alpha-\nu(z)/q(z)} (\log(e+1/s))^\beta \frac{ds}{s} \leq Cr^{\alpha-\nu(z)/q(z)} (\log(e+1/r))^\beta.$$

Since $\inf_{x \in G} (\nu(x) - \alpha p(x)) > 0$, taking ε_2 such that $0 < \varepsilon_2 < \nu(z)/p(z) - \alpha$, there exists a constant $c_2 > 0$ such that

$$s_2^{\alpha-\nu(z)/p(z)+\varepsilon_2} (\log(e+1/s_2))^\beta \leq c_2 s_1^{\alpha-\nu(z)/p(z)+\varepsilon_2} (\log(e+1/s_1))^\beta$$

whenever $0 < s_1 \leq s_2$, so that

$$\begin{aligned} & r^\theta \int_{k_1r}^{4k_2d_G} s^{\alpha-\nu(z)/p(z)} (\log(e+1/s))^\beta \frac{ds}{s} \\ & \leq c_2 r^\theta (k_1r)^{\alpha-\nu(z)/p(z)+\varepsilon_2} (\log(e+1/(k_1r)))^\beta \int_{k_1r}^{4k_2d_G} s^{-\varepsilon_2} \frac{ds}{s} \\ & \leq Cr^{\alpha-\nu(z)/p(z)+\theta} (\log(e+1/r))^\beta. \end{aligned}$$

We also have

$$r \int_{2k_1\tau r}^{4k_2\tau d_G} s^{\alpha-\nu(x)/p(x)-1+\theta} (\log(e+1/s))^\beta \frac{ds}{s} \leq Cr^{\alpha-\nu(x)/p(x)+\theta} (\log(e+1/r))^\beta$$

since $\inf_{x \in G} (\nu(x) - (\alpha + \theta - 1)p(x)) \geq \inf_{x \in G} (\nu(x) - \alpha p(x)) > 0$, and

$$r \int_{2k_1\tau r}^{4k_2\tau d_G} s^{\alpha-\nu(x)/q(x)-1} (\log(e+1/s))^\beta \frac{ds}{s} \leq Cr^{\alpha-\nu(x)/q(x)} (\log(e+1/r))^\beta$$

since $\inf_{x \in G} (\nu(x) - (\alpha - 1)q(x)) > 0$.

Collecting these facts, we obtain by our assumptions

$$\begin{aligned} \psi(x, z, r) & \leq C \{ r^{\alpha-\nu(x)/p(x)+\theta} + r^{\alpha-\nu(x)/q(x)} \\ & \quad + r^{\alpha-\nu(z)/p(z)+\theta} + r^{\alpha-\nu(z)/q(z)} \} (\log(e+1/r))^\beta < \infty \end{aligned}$$

for $x, z \in G$ and $0 < r \leq d_G$. By Theorem 2.2, we obtain the required result. \square

COROLLARY 3.2. *Suppose $\inf_{x \in G}(\nu(x) - \alpha p(x)) > 0$ and $\inf_{x \in G}((\alpha + \theta)p(x) - \nu(x)) > 0$. Further suppose $\inf_{x \in G}(\nu(x) - (\alpha - 1)q(x)) > 0$ and $\inf_{x \in G}(\alpha q(x) - \nu(x)) > 0$. Then there exists a constant $C > 0$ such that*

$$|b(x)I_\alpha f(x) - b(z)I_\alpha f(z)| \leq C \left\{ |x - z|^{\alpha - \nu(x)/p(x) + \theta} + |x - z|^{\alpha - \nu(x)/q(x)} \right. \\ \left. + |x - z|^{\alpha - \nu(z)/p(z) + \theta} + |x - z|^{\alpha - \nu(z)/q(z)} \right\}$$

for all $x, z \in G$ and measurable functions f on G with $\|f\|_{L^{\Phi, \nu(\cdot)}(G)} \leq 1$.

PROOF. This is the case $\beta = 0$ in Corollary 3.1. \square

Compare Corollaries 3.1 and 3.2 with [12, Theorem 4.1].

COROLLARY 3.3. *Let $\rho(r) = r^\alpha e^{-1/r}$ be as in Example 1.1. Then there exists a constant $C > 0$ such that*

$$|b(x)I_\rho f(x) - b(z)I_\rho f(z)| \leq C|x - z|^\theta$$

for all $x, z \in G$ and measurable functions f on G with $\|f\|_{L^{\Phi, \nu(\cdot)}(G)} \leq 1$.

PROOF. Since, for $a \in \mathbf{R}$, there exists a constant $c > 0$ such that

$$\int_0^r s^a e^{-1/s} \frac{ds}{s} \leq cr^\theta$$

for all $0 < r \leq d_G$, it follows from Remark 2.3 that

$$\psi(x, z, r) \leq C(r + r^\theta) \leq Cr^\theta$$

for all $x, z \in G$ and $0 < r \leq d_G$, since $\theta \in (0, 1]$. Hence, we obtain the required inequality. \square

COROLLARY 3.4. *Let $\rho(r) = r^\alpha (\log(e + 1/r))^\beta$ for $\alpha > 0$ and $\beta \in \mathbf{R}$. Suppose $\inf_{x \in G}(\nu(x) - (\alpha - 1)p(x)) > 0$ and $\inf_{x \in G}(\alpha p(x) - \nu(x)) > 0$. Then there exists a constant $C > 0$ such that*

$$|I_\rho f(x) - I_\rho f(z)| \\ \leq C \left\{ |x - z|^{\alpha - \nu(x)/p(x)} + |x - z|^{\alpha - \nu(z)/p(z)} \right\} (\log(e + 1/|x - z|))^\beta$$

for all $x, z \in G$ and measurable functions f on G with $\|f\|_{L^{p(\cdot), \nu(\cdot)}(G)} \leq 1$.

PROOF. To show this, we take $b(\cdot) \equiv 1$ and $q(\cdot) = p(\cdot)$ in the proof of Theorem 2.2. As in the proof of Corollary 3.1, we obtain the result. \square

4. LEMMAS

Before giving a proof of Theorem 2.2, we prepare two lemmas. To prove the following lemma, (P2) and (Q2) were used.

LEMMA 4.1 ([13, Lemma 2.1], cf. [14, Lemma 2.7]). *There exists a constant $C > 0$ such that*

$$\frac{r^{\nu(x)/p(x)}}{|B(x, r)|} \int_{G \cap B(x, r)} |f(y)| dy \leq C$$

for all $x \in G$, $0 < r < d_G$ and measurable functions f on G with $\|f\|_{L^{p(\cdot), \nu(\cdot)}(G)} \leq 1$.

LEMMA 4.2. *Let $\tau > 0, \beta \in \mathbf{R}$ and $\rho \in (\rho)$. Let f be a nonnegative function on G such that $\|f\|_{L^{p(\cdot), \nu(\cdot)}(G)} \leq 1$. Then there exists a constant $C > 0$ such that*

$$(4.3) \quad \int_{G \cap B(x, r)} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \leq C \int_0^{2k_2\tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s}$$

and

$$(4.4) \quad \int_{G \setminus B(x, r)} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \leq C \int_{k_1\tau r}^{4k_2\tau d_G} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s}$$

for all $x \in G$ and $0 < r \leq d_G$, where k_1 and k_2 are constants in (ρ) .

PROOF. Let f be a nonnegative function on G such that $\|f\|_{L^{p(\cdot), \nu(\cdot)}(G)} \leq 1$. Take $\gamma \in \mathbf{R}$ such that $1 < \gamma \leq \min\{1/k, 2\}$. If $y \in G \cap (B(x, \gamma^j r) \setminus B(x, \gamma^{j-1} r))$ for $j \in \mathbf{Z}$, then a geometric observation and (1.1) show

$$\begin{aligned} \frac{\rho(\tau|x-y|)}{|x-y|^{N+\beta}} &\leq \frac{\max\{1, \gamma^{N+\beta}\}}{(\gamma^j r)^{N+\beta}} \sup_{\gamma^{j-1}\tau r \leq s \leq \gamma^j \tau r} \rho(s) \\ &\leq \frac{\max\{1, \gamma^{N+\beta}\}}{(\gamma^j r)^{N+\beta}} \sup_{k\gamma^j \tau r \leq s \leq \gamma^j \tau r} \rho(s) \\ &\leq \frac{C_\rho \max\{1, \gamma^{N+\beta}\}}{(\gamma^j r)^{N+\beta}} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} \rho(s) \frac{ds}{s} \end{aligned}$$

by $\gamma \leq 1/k$. By Lemma 4.1, we have

$$\frac{1}{|B(x, \gamma^j r)|} \int_{G \cap B(x, \gamma^j r)} f(y) dy \leq C_1 (\gamma^j r)^{-\nu(x)/p(x)}$$

for some constant $C_1 > 0$, so that

$$\begin{aligned}
& \int_{G \cap (B(x, \gamma^j r) \setminus B(x, \gamma^{j-1} r))} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \\
& \leq \frac{C_\rho \sigma_N \max\{1, \gamma^{N+\beta}\}}{(\gamma^j r)^\beta} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} \rho(s) \frac{ds}{s} \cdot \frac{1}{|B(x, \gamma^j r)|} \int_{G \cap B(x, \gamma^j r)} f(y) dy \\
& \leq \frac{C_\rho \sigma_N \max\{1, \gamma^{N+\beta}\}}{(\gamma^j r)^\beta} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} \rho(s) \frac{ds}{s} \cdot C_1 (\gamma^j r)^{-\nu(x)/p(x)} \\
& \leq C_1 C_\rho \sigma_N \max\{1, 2^{N+\beta}\} (\gamma^j r)^{-\nu(x)/p(x)-\beta} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} \rho(s) \frac{ds}{s} \\
& \leq C_1 C_\rho \sigma_N \max\{1, 2^{N+\beta}\} \\
& \times \max\left\{(\tau k_1)^{\nu(x)/p(x)+\beta}, (\tau k_2)^{\nu(x)/p(x)+\beta}\right\} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s} \\
& \leq C_2 \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s},
\end{aligned}$$

where σ_N denotes the volume of the unit ball $B(0, 1)$ and

$$\begin{aligned}
C_2 &= C_1 C_\rho \sigma_N \max\{1, 2^{N+\beta}\} \\
& \times \max\left\{(\tau k_1)^{\nu^+/p^- + \beta}, (\tau k_1)^{\nu^-/p^+ + \beta}, (\tau k_2)^{\nu^+/p^- + \beta}, (\tau k_2)^{\nu^-/p^+ + \beta}\right\}.
\end{aligned}$$

Therefore we have

$$(4.5) \quad \int_{G \cap (B(x, \gamma^j r) \setminus B(x, \gamma^{j-1} r))} \leq C_2 \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s}.$$

Let j_0 be the smallest integer such that $k_2/k_1 \leq \gamma^{j_0}$. Using (4.5), we obtain

$$\begin{aligned}
& \int_{G \cap B(x, r)} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy = \sum_{j=0}^{\infty} \int_{G \cap (B(x, \gamma^{-j} r) \setminus B(x, \gamma^{-j-1} r))} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \\
& \leq C_2 \sum_{j=0}^{\infty} \int_{\gamma^{-j} k_1 \tau r}^{\gamma^{-j} k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s} \leq C_2 \sum_{j=0}^{\infty} \int_{\gamma^{-j} k_1 \tau r}^{\gamma^{-j+j_0} k_1 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s} \\
& \leq j_0 C_2 \int_0^{2k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s},
\end{aligned}$$

which proves (4.3).

Let j_1 be the smallest integer such that $d_G \leq \gamma^{j_1} r$. If we use (4.5),

$$\begin{aligned}
& \int_{G \setminus B(x,r)} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \leq \sum_{j=1}^{j_1} \int_{G \cap (B(x,\gamma^j r) \setminus B(x,\gamma^{j-1} r))} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\beta}} dy \\
& \leq C_2 \sum_{j=1}^{j_1} \int_{\gamma^j k_1 \tau r}^{\gamma^j k_2 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s} \\
& \leq C_2 \sum_{j=1}^{j_1} \int_{\gamma^j k_1 \tau r}^{\gamma^{j+j_0} k_1 \tau r} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s} \\
& \leq C_2 j_0 \int_{k_1 \tau r}^{4k_2 \tau d_G} s^{-\nu(x)/p(x)-\beta} \rho(s) \frac{ds}{s}.
\end{aligned}$$

Thus, (4.4) follows. \square

5. PROOF OF THEOREM

Without loss of generality, we can assume that f is a nonnegative function on G such that $\|f\|_{L^{\Phi, \nu(\cdot)}(G)} \leq 1$. First note from (2.2) that for $x, z \in G$ and $r = |x - z|$

$$\begin{aligned}
& |b(x)I_\rho f(x) - b(z)I_\rho f(z)| \\
& \leq b(x) \int_{G \cap B(x,2r)} \frac{\rho(|x-y|)f(y)}{|x-y|^N} dy + b(z) \int_{G \cap B(x,2r)} \frac{\rho(|z-y|)f(y)}{|z-y|^N} dy \\
& + |b(x) - b(z)| \int_{G \setminus B(x,2r)} \frac{\rho(|z-y|)f(y)}{|z-y|^N} dy \\
& + b(x) \int_{G \setminus B(x,2r)} \left| \frac{\rho(|x-y|)}{|x-y|^N} - \frac{\rho(|z-y|)}{|z-y|^N} \right| f(y) dy \\
& \leq C \left\{ b(x) \int_{G \cap B(x,2r)} \frac{\rho(|x-y|)f(y)}{|x-y|^N} dy + b(z) \int_{G \cap B(z,3r)} \frac{\rho(|z-y|)f(y)}{|z-y|^N} dy \right. \\
& \left. + r^\theta \int_{G \setminus B(z,r)} \frac{\rho(|z-y|)f(y)}{|z-y|^N} dy + r^{\eta_1} b(x) \int_{G \setminus B(x,2r)} \frac{\rho(\tau|x-y|)f(y)}{|x-y|^{N+\eta_2}} dy \right\} \\
& = C \left\{ I_1(x) + \tilde{I}_1(z) + I_2(z) + I_3(x) \right\}.
\end{aligned}$$

For $I_1(x)$, we have

$$\begin{aligned} I_1(x) &\leq \int_{G \cap B(x, 2r)} \frac{\rho(|x-y|)}{|x-y|^N} |b(x) - b(y)| f(y) dy \\ &\quad + \int_{G \cap B(x, 2r)} \frac{\rho(|x-y|)}{|x-y|^N} b(y) f(y) dy \\ &\leq C \int_{G \cap B(x, 2r)} \frac{\rho(|x-y|) f(y)}{|x-y|^{N-\theta}} dy + \int_{G \cap B(x, 2r)} \frac{\rho(|x-y|) \{b(y) f(y)\}}{|x-y|^N} dy \\ &= CI_{11}(x) + I_{12}(x). \end{aligned}$$

By (4.3), we obtain

$$I_{11}(x) \leq C \int_0^{4k_2 r} s^{-\nu(x)/p(x)+\theta} \rho(s) \frac{ds}{s},$$

and

$$I_{12}(x) \leq C \int_0^{4k_2 r} s^{-\nu(x)/q(x)} \rho(s) \frac{ds}{s}.$$

For $\tilde{I}_1(z)$, we have by (4.3)

$$\tilde{I}_1(z) \leq C \left\{ \int_0^{6k_2 r} s^{-\nu(z)/p(z)+\theta} \rho(s) \frac{ds}{s} + \int_0^{6k_2 r} s^{-\nu(z)/q(z)} \rho(s) \frac{ds}{s} \right\}$$

as in the estimate of $I_{11}(x)$ and $I_{12}(x)$.

For $I_2(z)$, we have by (4.4)

$$I_2(z) \leq Cr^\theta \int_{k_1 r}^{4k_2 d_G} s^{-\nu(z)/p(z)} \rho(s) \frac{ds}{s}.$$

Finally, for $I_3(x)$ we have

$$\begin{aligned} I_3(x) &\leq r^{\eta_1} \int_{G \setminus B(x, 2r)} \frac{\rho(\tau|x-y|)}{|x-y|^{N+\eta_2}} |b(x) - b(y)| f(y) dy \\ &\quad + r^{\eta_1} \int_{G \setminus B(x, 2r)} \frac{\rho(\tau|x-y|)}{|x-y|^{N+\eta_2}} b(y) f(y) dy \\ &\leq Cr^{\eta_1} \int_{G \setminus B(x, 2r)} \frac{\rho(\tau|x-y|) f(y)}{|x-y|^{N-\theta+\eta_2}} dy \\ &\quad + r^{\eta_1} \int_{G \setminus B(x, 2r)} \frac{\rho(\tau|x-y|) \{b(y) f(y)\}}{|x-y|^{N+\eta_2}} dy \\ &= CI_{31}(x) + I_{32}(x). \end{aligned}$$

Note from (4.4) that

$$I_{31}(x) \leq Cr^{\eta_1} \int_{2k_1 \tau r}^{4k_2 \tau d_G} s^{-\nu(x)/p(x)-\eta_2+\theta} \rho(s) \frac{ds}{s}$$

and

$$I_{32}(x) \leq C r^{\eta_1} \int_{2k_1 \tau r}^{4k_2 \tau d_G} s^{-\nu(x)/q(x) - \eta_2} \rho(s) \frac{ds}{s}.$$

Collecting these facts, we obtain

$$|b(x)I_\rho f(x) - b(z)I_\rho f(z)| \leq C\psi(x, z, r).$$

Thus this theorem is proved.

ACKNOWLEDGEMENTS.

We would like to express our thanks to the referees for their kind comments and helpful suggestions.

REFERENCES

- [1] P. Baroni, M. Colombo and G. Mingione, *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations **57** (2018), no. 2, Paper No. 62, 48 pp.
- [2] S. S. Byun, S. Liang and S. Zheng, *Nonlinear gradient estimates for double phase elliptic problems with irregular double obstacles*, Proc. Amer. Math. Soc. **147** (2019), 3839–3854.
- [3] S. S. Byun and H. S. Lee, *Calderón-Zygmund estimates for elliptic double phase problems with variable exponents*, J. Math. Anal. Appl. **501** (2021), 124015.
- [4] M. Colombo and G. Mingione, *Regularity for double phase variational problems*, Arch. Rat. Mech. Anal. **215** (2015), 443–496.
- [5] M. Colombo and G. Mingione, *Bounded minimizers of double phase variational integrals*, Arch. Rat. Mech. Anal. **218** (2015), 219–273.
- [6] Eridani, H. Gunawan, E. Nakai and Y. Sawano, *Characterizations for the generalized fractional integral operators on Morrey spaces*, Math. Ineq. Appl. **17** (2014), no. 2, 761–777.
- [7] C. De Filippis and G. Mingione, *Manifold constrained non-uniformly elliptic problems*, J. Geom. Anal. **30** (2020), no. 2, 1661–1723.
- [8] P. Hästö and J. Ok, *Maximal regularity for local minimizers of non-autonomous functionals*, to appear in J. Eur. Math. Soc.
- [9] F.-Y. Maeda, Y. Mizuta, T. Ohno and T. Shimomura, *Boundedness of maximal operators and Sobolev's inequality on Musielak-Orlicz-Morrey spaces*, Bull. Sci. Math., **137** (2013), 76–96.
- [10] F. - Y. Maeda, Y. Mizuta, T. Ohno and T. Shimomura, *Sobolev's inequality for double phase functionals with variable exponents*, Forum Math. **31** (2019), 517–527.
- [11] F. - Y. Maeda, Y. Mizuta, T. Ohno and T. Shimomura, *Trudinger's inequality for double phase functionals with variable exponents*, Czechoslovak Math. J. **71** (2021), 511–528.
- [12] Y. Mizuta, E. Nakai, T. Ohno and T. Shimomura, *Campanato-Morrey spaces for the double phase functionals*, Rev. Mat. Complut. **33** (2020), 817–834.
- [13] Y. Mizuta, E. Nakai, T. Ohno and T. Shimomura, *Campanato-Morrey spaces for the double phase functionals with variable exponents*, Nonlinear Anal. **197** (2020), article no. 111827, 19 pp.
- [14] Y. Mizuta, E. Nakai, T. Ohno and T. Shimomura, *Riesz potentials and Sobolev embeddings on Morrey spaces of variable exponent*, Complex Var. Elliptic Equ. **56**, No. 7-9, (2011), 671–695.

- [15] Y. Mizuta, E. Nakai, Y. Sawano and T. Shimomura, *Gagliardo-Nirenberg inequality for generalized Riesz potentials of functions in Orlicz-Musielak spaces*, Arch. Math. **98** (2012), 253–263.
- [16] Y. Mizuta, T. Ohno and T. Shimomura, *Sobolev’s theorem for double phase functionals*, Math. Ineq. Appl. **23** (2020), 17–33.
- [17] T. Shimomura and Y. Mizuta, *Taylor expansion of Riesz potentials*, Hiroshima Math. J. **25** (1995), 595–621.
- [18] Y. Mizuta and T. Shimomura, *Differentiability and Hölder continuity of Riesz potentials of Orlicz functions*, Analysis (Munich) **20** (2000), no. 3, 201–223.
- [19] C. B. Morrey, *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43** (1938), 126–166.
- [20] J. Musielak, *Orlicz spaces and modular spaces*, Lecture Notes in Math. **1034**, Springer-Verlag, 1983.
- [21] S. Nagayasu and H. Wadade, *Characterization of the critical Sobolev space on the optimal singularity at the origin*, J. Funct. Anal. **258** (2010), no. 11, 3725–3757.
- [22] E. Nakai, *On generalized fractional integrals*, Taiwan. J. Math. **5** (2001), 587–602.
- [23] A. Ourraoui and M. A. Ragusa, *An existence result for a class of $p(x)$ -anisotropic type equations*, Symmetry **13** (2021), no. 4, 633.
- [24] C. Pérez, *Sharp L^p -weighted Sobolev inequalities*, Ann. Inst. Fourier (Grenoble) **45** (1995), 809–824.
- [25] E. Pustylnik, *Generalized potential type operators on rearrangement invariant spaces*, Israel Math. Conf. Proc. **13** (1999), no. 3, 161–171.
- [26] M. A. Ragusa and A. Tachikawa, *Regularity for minimizers for functionals of double phase with variable exponents*, Adv. Nonlinear Anal. **9** (2020), no. 1, 710–728.
- [27] Y. Sawano and T. Shimomura, *Boundedness of the generalized fractional integral operators on generalized Morrey spaces over metric measure spaces*, Zeit. Anal. Anwend. **36** (2017), 159–190.
- [28] Y. Sawano, S. Sugano and H. Tanaka, *Orlicz-Morrey spaces and fractional operators*, Potential Anal. **36** (2012), no. 4, 517–556.
- [29] A. Tachikawa, *Boundary regularity of minimizers of double phase functionals*, J. Math. Anal. Appl. **501** (2021), no. 1, 123946, 34 pp.
- [30] V. V. Zhikov, *Averaging of functionals of the calculus of variations and elasticity theory*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), 675–710.

T. Ohno
Faculty of Education
Oita University
DannoHaru Oita-city 870-1192, Japan
E-mail: t-ohno@oita-u.ac.jp

T. Shimomura
Department of Mathematics
Graduate School of Humanities and Social Sciences
Hiroshima University
Higashi-Hiroshima 739-8524, Japan
E-mail: tshimo@hiroshima-u.ac.jp

T. Ohno
Faculty of Education
Oita University
DannoHaru Oita-city 870-1192, Japan
E-mail: t-ohno@oita-u.ac.jp

T. Shimomura
Department of Mathematics
Graduate School of Humanities and Social Sciences
Hiroshima University
Higashi-Hiroshima 739-8524, Japan
E-mail: `tshimo@hiroshima-u.ac.jp`