



Glasnik
Matematički

SERIJA III

www.math.hr/glasnik

Mario Osvin Pavčević and Kristijan Tabak
CZ-groups with nonabelian normal subgroup of order p^4

Accepted manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by Glasnik Production staff.

***CZ*-GROUPS WITH NONABELIAN NORMAL SUBGROUP OF ORDER p^4**

MARIO OSVIN PAVČEVIĆ AND KRISTIJAN TABAK

University of Zagreb, Croatia and Rochester Institute of Technology

ABSTRACT. A p -group G with the property that its every nonabelian subgroup has a trivial centralizer (namely only its center) is called a *CZ*-group. In Berkovich's monograph (see [1]) the description of the structure of a *CZ*-group was posted as a research problem. Here we provide further progress in this topic based on results proved in [5]. In this paper we have described the structure of *CZ*-groups G that possess a nonabelian normal subgroup of order p^4 which is contained in the Frattini subgroup $\Phi(G)$. We manage to prove that such a group of order p^4 is unique and that the order of the entire group G is less or equal p^7 , p being a prime. Additionally, all such groups G are shown to be of class less than maximal.

1. INTRODUCTION

A p -group G is a group of order p^n , where p is prime. The conjugation of x by y is given by $x^y = y^{-1}xy$, where $x, y \in G$. If $x^y = x$, then x and y commute, i.e. $[x, y] = x^{-1}y^{-1}xy = 1$. Let $H \leq G$ be a subgroup of G . The centralizer of H in G is $C_G(H) = \{g \in G \mid h^g = h, \forall h \in H\}$. The center of G is given by $Z(G) = \{g \in G \mid x^g = x, \forall x \in G\}$. The center $Z(H)$ of a subgroup $H \leq G$ is defined in the same way.

A finite group G is called a **CZ-group** (this abbreviated form comes from the words centralizer and Zentrum) if $C_G(H) = Z(H)$ for all nontrivial $H \leq G$. The set of all *CZ*-groups that are at the same time p -groups will be denoted by CZ_p and sometimes we will call such a group a CZ_p -group. The question of determining the general structure of $G \in CZ_p$ was posted in [1] as one of the open problems in the theory of p -groups. More on p -groups can be

2020 *Mathematics Subject Classification.* 20D15, 20D25.

Key words and phrases. p -group, center, centralizer, Frattini subgroup, minimal non-abelian subgroup.

found in [2] and [3]. The first results about groups $G \in CZ_p$ were published in [5], where it was shown that a minimal CZ_p -group has order at least p^5 . Additionally, the structure of maximal abelian subgroups of a minimal CZ_p group has been described in that paper as well.

In this paper, we assume that the Frattini subgroup $\Phi(G)$, which is defined as the intersection of all maximal subgroups of G , contains a normal nonabelian subgroup of order p^4 . A subgroup $H \leq G$ that is a normal subgroup of G will be sometimes called G -invariant (if we want to point out this fact, we will denote it by $H \trianglelefteq G$). The existence of a normal subgroup of order p^4 in $\Phi(G)$ doesn't appear as a limitation, since we can always find normal subgroups in p -groups of any given order. What however appears as a true assumption is that we in addition assume for this subgroup of order p^4 to be nonabelian.

In the next result we will determine the lower bound for the order $|G|$ of $G \in CZ_p$.

LEMMA 1.1. *If $G \in CZ_p$, then $|G : Z(G)| \geq p^3$ and $|G| \geq p^5$.*

PROOF. Let's assume the opposite, so let $|G : Z(G)| \leq p^2$. Then immediately we get $|G : Z(G)| = p^2$, since otherwise G would be abelian. The factor group $G/Z(G)$ can't be cyclic, otherwise G would be abelian again. Thus, $G/Z(G) \cong E_{p^2}$ (the elementary abelian group of order p^2). Since the Frattini subgroup is the smallest subgroup such that its factor group is elementary abelian, we get $\Phi(G) \leq Z(G)$. If $\Phi(G) < Z(G)$, there is some maximal subgroup M such that $Z(G) \not\leq M$. Hence, M must be abelian, since otherwise, we would be able to find some $g \in Z(G) \setminus M$, leading further to $g \in C_G(M)$, which is a contradiction since $G \in CZ_p$. Therefore, $MZ(G) = G$ and G is abelian, which is a contradiction again. So, $|G : Z(G)| \geq p^3$ and $|G| \geq p^4$ (since $|Z(G)| \geq p$). If $|G| = p^4$, then $|Z(G)| = p$ and $Z(G) \leq \Phi(G)$. This implies that any maximal subgroup of G is minimally nonabelian, thus G is a minimal CZ group, from which follows that $|G| \geq p^5$ (as it was proved in [5]). This is a contradiction. Therefore, the only remaining option is $|G| \geq p^5$. \square

LEMMA 1.2. *Let $G \in CZ_p$ and $M < G$, $M \in CZ_p$. Then $|G : Z(G)| \geq p^4$ and $|G| \geq p^5$.*

PROOF. Lemma 1.1 states that $|G : Z(G)| \geq p^3$. Let $M \in CZ_p$ and $M < G$. Then again by Lemma 1.1, $|M : Z(M)| \geq p^3$. It was proved in [5] that $Z(G) \leq Z(M)$. Thus $|G : Z(G)| > |M : Z(G)| \geq |M : Z(M)| \geq p^3$. Therefore, $|G : Z(G)| \geq p^4$ and $|G| \geq p^5$. \square

The following statement establishes a connection between CZ_p -groups and the maximality of class.

THEOREM 1.3. *Let $G \in CZ_p$ and $B < G$ be a nonabelian group of order p^3 . Then, G is a group of maximal class.*

PROOF. Let $B < G$, where $|B| = p^3$ and B nonabelian. Then $C_G(B) < B$. Therefore, $Z(G) \leq Z(B)$. Clearly, $|Z(B)| = p$ and $Z(G) = Z(B)$.

It is known that if $H \in \text{Syl}_p(\text{Aut}(B))$ (a Sylow p -group), then $|H| = p^3$ and H is nonabelian. Therefore, $N_G(B)/C_G(B) \leq \text{Aut}(B)$ is a p -group. Also, $N_G(B) > B$ and $C_G(B) = Z(B)$. Therefore, $|N_G(B)/C_G(B)| \geq p^3$ since $|N_G(B) : C_G(B)| = |N_G(B) : B| \cdot |B : Z(B)| \geq p \cdot p^2 = p^3$. Thus, it is necessary that $N_G(B)/Z(G) \cong H \in \text{Syl}_p(\text{Aut}(B))$. Also, $N_G(B)/Z(G) < G/Z(G)$ (since $|Z(G)| = p$ and $|G| \geq p^5$ and H nonabelian of order p^3).

Obviously $C_G(N_G(B)/Z(G)) \leq N_G(B)/Z(G)$. Inductively, $G/Z(G)$ is of maximal class, where $|Z(G)| = p$. From here we deduce that G is of maximal class. \square

2. CZ-GROUPS WITH NONABELIAN G -INVARIANT SUBGROUP $N \leq \Phi(G)$ OF ORDER p^4

Let us now we introduce the main assumption. We will assume further that G is a CZ_p group possessing a subgroup $N \leq \Phi(G)$ which is a nonabelian G -invariant subgroup of order p^4 . The nilpotency class of a group G will be denoted by $cl(G)$. If the class is maximal, we will put $cl(G) = max$, otherwise $cl(G) < max$. If the group is generated by at least k elements, we shall say that it is a k -generated group and write $d(G) = k$.

We will make use of the following result. Its proof can be found in [1] (Lemma 1.4.).

LEMMA 2.1. *Let G be a p -group for $p > 2$ and $N \trianglelefteq G$. If N has no abelian G -invariant subgroups of type (p, p) , then N is cyclic.*

The structure of a p -subgroup N satisfying the properties mentioned above is partially described in the following result.

LEMMA 2.2. *Let $G \in CZ_p$ where $p > 2$ and $cl(G) < max$. Let $N \leq \Phi(G)$ be a G -invariant nonabelian group of order p^4 . Then $\Phi(N) = Z(N) \cong E_{p^2}$ and N is a 2-generated group of exponent p^2 .*

PROOF. Assume that $Z(N)$ is cyclic. Let $A \trianglelefteq G$ and $A \leq N$ of order p^2 . Then $|N_G(A)/C_G(A)| = |G/C_G(A)| \leq |\text{Aut}(A)|_p = p$, where $|\text{Aut}(A)|_p$ is the maximal power of p that divides $|\text{Aut}(A)|$. Hence, $N \leq \Phi(G) \leq C_G(A)$ and $A \leq C_G(A)$. Thus, $A \leq Z(\Phi(G)) \cap N$ and $A \leq Z(N)$ (since $N \leq \Phi(G)$). Therefore, A is cyclic. Then, according to Lemma 2.1, N must be cyclic, which is a contradiction. Therefore, $Z(N)$ is not cyclic. If $d(Z(N)) \geq 3$, then $|Z(N)| \geq p^3$ and $|N : Z(N)| \leq p$. This would imply that N is abelian. Therefore, $d(Z(N)) = 2$ and $Z(N) \cong E_{p^2}$. Clearly, $d(N) \geq 2$. Assume that $d(N) = 4$. Then $N/\Phi(N) \cong E_{p^4}$ and $\Phi(N) = 1$. On the other hand, $\Phi(N) = N'\mathcal{U}_1(N)$ and $N' = 1$. This is a contradiction. Thus, $d(N) \leq 3$.

If $d(N) = 3$, then $\Phi(N) \cong C_p$ and $1 < N' \leq \Phi(N)$. Thus, $\Phi(N) = N'$. Clearly, $N' \cap Z(N) > 1$ since $N' \triangleleft N$. Put $Z(N) = \langle x \rangle \times \langle y \rangle \cong C_p \times C_p$ such that $N' = \langle x \rangle$. Thus, there is $y \in Z(N) - \Phi(N)$.

Thus, y is a generator of N and the order of y is p . Then there is some maximal subgroup $M < N$ such that $y \notin M$. Therefore, $N = \langle M, y \rangle = M \times \langle y \rangle$. If $w^p = y$ for some w , then $y \in \mathcal{U}_1(N) \leq \Phi(N)$. This is a contradiction since y is a generator. Because N is nonabelian, M must be nonabelian, otherwise $N = M \times \langle y \rangle$ would be abelian. Therefore, $M' > 1$ and $|M| = p^3$. Then, according to the Theorem 1.3, the class of the group G is maximal. This is a contradiction with our assumption. Therefore, $d(N) = 2$ and $N/\Phi(N) \cong E_{p^2}$ where $|\Phi(N)| = p^2$. If $Z(N) \not\leq \Phi(N)$, then there is some maximal $M \triangleleft_p N$ such that $Z(N) \not\leq M$ then $M' = 1$. Otherwise, by Theorem 1.3 we would get $cl(G) = max$. Thus, it is necessary that $Z(N) \leq \Phi(N)$. Since both groups have order p^2 , we get $Z(N) = \Phi(N)$.

Let $exp(G) = p$. Then $|\mathcal{U}_1(G)| = 1$ and $\Phi(N) = N'\mathcal{U}_1(N) = N' \cong C_p \times C_p$. Since N has a maximal abelian subgroup, then $p^4 = |N| = p \cdot |N'| \cdot |Z(N)| = p \cdot p^2 \cdot p^2$. This is a contradiction. Thus, $exp(N) > p$.

If $exp(N) = p^3$, then $N \cong M_{p^4}$, where M_{p^4} is a minimal nonabelian group with a maximal cyclic subgroup. Then, there is some $w \in N$ of order p^3 . Hence $\mathcal{U}_1(N) = \langle w^p \rangle \cong C_{p^2}$ and $\mathcal{U}_1(N) = \Phi(N) = Z(N)$ and $d(N) = 1$. Again, this is a contradiction. Thus, the only remaining option is $exp(N) = p^2$. \square

The following result shows the uniqueness of the nonabelian G -invariant subgroup $N \leq \Phi(G)$, where $cl(G) < max$, $G \in CZ_p$ and $|N| = p^4$.

LEMMA 2.3. *Let $G \in CZ_p$, $p > 2$ and $cl(G) < max$. Let $N \leq \Phi(G)$ be a G -invariant nonabelian subgroup of order p^4 . Then N is uniquely determined by its generators and relations with $N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$.*

PROOF. From Lemma 2.2 we have $Z(N) = \Phi(N) \cong E_{p^2}$. Also, $exp(N) = p^2$. If $M \triangleleft N$ is maximal, then $Z(N) < M$ and $M' = 1$. Thus, $|N| = p \cdot |N'| \cdot |Z(N)|$. Hence, $|N'| = p$. We can put $Z(N) = \langle a \rangle \times \langle b \rangle$. We can assume $N' = \langle a \rangle$. There are $x, y \in N$ such that $x^p = a$, $y^p = b$. Otherwise, $a \notin \Phi(N) = N'\mathcal{U}_1(N)$ and $b \notin \Phi(N)$. Now, take $[x, y] = a = x^{-1}y^{-1}xy = x^{-1}x^y = x^p$. This gives us $x^y = x^{1+p}$. \square

LEMMA 2.4. *Let the group N be defined as $N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. Let $z^{-1} = x^p$. Then for all integers i, j, n the following relations hold: $y^j x = xy^j z^j$, $y^j x^i = x^i y^j z^{ij}$. Furthermore, $(x^i y^j)^n = x^{ni} y^{nj} z^{\binom{n}{2} ij}$ and the order $o(g) = p^2$, for all $g \in N - \Phi(N)$. The subgroup $\langle x^i y^{pj} \rangle \leq N$ is normal in N .*

PROOF. Since $z^{-1} = x^p \in Z(N)$, it follows $x^y = xz^{-1}$ and $xyz = yx$. Then, $y^j x = y^{j-1}(yx) = y^{j-1}(xy)z = y^{j-2}(yx)yz = y^{j-2}(xy)yz^2 = y^{j-2}xy^2z^2 = \dots = xy^j z^j$. We have $y^j x^i = y^j x x^{i-1} = xy^j x^{i-1} z^j = xy^j x x^{i-2} z^j =$

$xx^j x^{i-2} z^{2j} = \dots = x^i y^j z^{ij}$. We will use induction to prove the claim about $(x^i y^j)^n$. For $n = 1$ the claim is trivial. Assume that $(x^i y^j)^n = x^{ni} y^{nj} z^{\binom{n}{2}ij}$. Now we proceed with the induction step by computing

$$\begin{aligned} (x^i y^j)^{n+1} &= x^{ni} y^{nj} z^{\binom{n}{2}ij} x^i y^j = x^{ni} y^{nj} x^i y^j z^{\binom{n}{2}ij} = x^{ni} x^i y^{nj} y^j z^{ni j} z^{\binom{n}{2}ij} = \\ &= x^{(n+1)i} y^{(n+1)j} z^{\binom{n+1}{2}ij}. \end{aligned}$$

Let $g \notin \Phi(N)$. Then $g = x^i y^j$, where either i or j is not divisible by p . Otherwise, $g \in \langle x^p, y^p \rangle = \Phi(N) = Z(N)$, (see Lemma 2.2). Since $z^p = 1$ and $p \mid \binom{p}{2}$, we obtain $(x^i y^j)^p = x^{pi} y^{pj} z^{\binom{p}{2}ij} = x^{pi} y^{pj}$. If $g^p = 1$, then $x^{pi} = y^{-pj} \in \langle x \rangle \cap \langle y \rangle$, which implies $x^{pi} = 1$ and $i \equiv (0 \pmod p)$. In the other case $j \equiv (0 \pmod p)$. This is a contradiction. Therefore $o(g) = p^2$.

Look now at $x^i y^{pj}$, where i and j are not divisible by p . Since $y^p \in Z(N)$, it follows $(x^i y^{pj})^x = x^i y^{pj}$. Let us assume that there is some integer n such that $(x^i y^{pj})^y = (x^i y^{pj})^n$. This would imply $\langle x^i y^{pj} \rangle \trianglelefteq N$. If such an n exists, this would imply $(xz^{-1})^i y^{pj} = (x^i y^{pj})^n$. Then, $x^i y^{pj} z^{-i} = x^{ni} y^{npj}$ and $y^{npj-pj} \in \langle x \rangle$. Thus $pj(n-1) \equiv 0 \pmod{p^2}$ and $n-1 \equiv 0 \pmod p$ since $j \not\equiv 0 \pmod p$. Let $n = 1 + mp$, for some integer m . Then $x^{i(1-n)} = z^i$ and $x^{-mpi} = z^i$. Therefore, $z^{mi} = z^i$. Take $m = 1$ and $n = 1 + p$. We conclude, such n exists and $\langle x^i y^{pj} \rangle \trianglelefteq N$. \square

LEMMA 2.5. *Let $G \in CZ_p$, $p > 2$ and $cl(G) < \max$. Let $N \trianglelefteq G$ and $N \leq \Phi(G)$ be nonabelian of order p^4 . Then $G/Z(N)$ is isomorphic to some subgroup of $Aut(N)$.*

PROOF. Since $N_G(N) = G$ and $C_G(N) \leq N$, we get $C_G(N) = Z(N)$. Then by the N/C -theorem, $N_G(N)/C_G(N) \lesssim Aut(N)$. \square

The following results is from [4] (Theorem 12.2.2, page 178).

THEOREM 2.6. *Let $|G| = p^n$ and $d(G) = d$. Then $|Aut(G)|$ divides $|Aut(E_{p^d}) \times \Phi(G)^d|$.*

The following result establishes an upper bound for the order of a group G with conditions we are studying here.

THEOREM 2.7. *Let $G \in CZ_p$, $cl(G) < \max$ and let $N \leq \Phi(G)$ be a normal nonabelian subgroup of G of order p^4 . Then $|G| \leq p^7$.*

PROOF. According to Lemma 2.2 and Lemma 2.3, $\Phi(N) = Z(N) \cong E_{p^2}$ and N is uniquely determined by $N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. Applying Lemma 2.5 and Theorem 2.6, we have $G/Z(N) \lesssim Aut(N)$ and $|G/Z(N)|_p$ divides $|Aut(E_{p^2}) \times \Phi(N)^2|_p = |(p^2-1)(p^2-p) \cdot p^4|_p = p^5$. Since $|Z(N)| = p^2$, we get $|G| \leq p^7$. \square

COROLLARY 2.8. *Let $G \in CZ_p$, $cl(G) < \max$ and $N \leq \Phi(G)$ a normal nonabelian subgroup of order p^4 . Then $|G| \in \{p^5, p^6, p^7\}$.*

We conclude this section with a technical result that we shall need.

LEMMA 2.9. *Let $\langle x \rangle \cong C_{p^2}$, $p > 2$ and let $\varphi \in \text{Aut}(\langle x \rangle)$ be an automorphism of $\langle x \rangle$ of order p . Then, there is some $m \in \mathbb{N}$ such that $x^\varphi = x^{1+mp}$.*

3. THE CASE $|G| = p^6$ AND $cl(G) < \max$

We shall continue with the same assumption that G possesses a non-abelian subgroup $N \leq \Phi(G)$ of order p^4 . Additionally, we shall assume that G is **not of maximal class**. By Corollary 2.8, the order of G is at least p^5 . If $|G| = p^5$, then $|G : \Phi(G)| = p$ and G is cyclic. Therefore, from this moment on, we can assume that $|G| \geq p^6$. If $|G| = p^6$, then $|G : \Phi(G)| = p^2$ and G is a 2-generated group.

We now prove additional results about the structure of the group N .

LEMMA 3.1. *Let $N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. Then $Z(N) = \Phi(N) = \langle x^p, y^p \rangle \cong E_{p^2}$ and $\langle y^{p^{i+1}} \rangle \not\triangleleft N$, $i = 0, 1, \dots, p-1$.*

Proof: From $(x^p)^y = (x^y)^p = (x^{1+p})^p = x^p$ and $[x, x^p] = 1$ we have $x^p \in Z(N)$. Furthermore, $x^{y^p} = x^{(1+p)^p} = x^p$ (since $(1+p)^p \equiv p \pmod{p^2}$). Therefore, $\langle x^p, y^p \rangle \leq Z(N)$. Since $|N : Z(N)| \geq p^2$ and $\langle x^p \rangle \cap \langle y^p \rangle = 1$, we have $Z(N) = \langle x^p, y^p \rangle$. Since N is 2-generated and $N/Z(N) \cong E_{p^2}$, it follows that $Z(N) = \Phi(N)$.

Now we shall prove the second claim. We firstly use the following: $(y^{p^{i+1}})^x = (y^x)^{p^{i+1}} = (y^{1+p})^{p^{i+1}} = y^{p^{i+1}} x^{-p}$. If $\langle y^{p^{i+1}} \rangle$ is N -invariant, then $y^{p^{i+1}} x^{-p} \in \langle y^{p^{i+1}} \rangle$. This implies $x^{-p} \in \langle y^{p^{i+1}} \rangle \leq \langle y \rangle$ and $\langle x \rangle \cap \langle y \rangle > 1$, which is a contradiction. Therefore, $\langle y^{p^{i+1}} \rangle \not\triangleleft N$. \square

The following two results have been proved in [1]. We shall present them here with slightly different proofs. We will use the following notation: if H is a normal subgroup of index p^i of G , then we shall write this as $H \triangleleft_{p^i} G$.

THEOREM 3.2. *Let G be a p -group and let $K \trianglelefteq G$ contain a abelian maximal subgroup. Then K contains a maximal abelian subgroup that is G -invariant.*

PROOF. If G is an abelian group, the claim is true. Let G be a nonabelian group, and let $A \triangleleft_p K \trianglelefteq G$, where A is an abelian subgroup. If $\{T \mid T \triangleleft_p K, T' = 1\} = \{A\}$, then $A^g \triangleleft_p K^g = K$ for all $g \in G$ (A^g is abelian as well). Therefore, $A^g = A$ for all $g \in G$. This implies that A is G -invariant.

Now assume that A_1 and A_2 are distinct maximal abelian subgroups of K . Then $A_i \triangleleft K$ and $A_1 A_2 = K$. Since $A_1 \cap A_2 \triangleleft_p A_i$, we have $A_1 \cap A_2 \leq C_K(A_1) \cap C_K(A_2)$. This implies $A_1 \cap A_2 \leq Z(K)$. Let K be a non-abelian group. Then $K/Z(K) \cong E_{p^2}$. There is a subgroup $C \leq K$ such that $C/Z(K) \cong C_p$. Then $C/Z(K) \triangleleft K/Z(K)$. There is a one-to-one map between $\{C/Z(K) \mid C/Z(K) \triangleleft K/Z(K)\}$ and $\{C/Z(K) \mid Z(K) \triangleleft_p C \triangleleft_p K\}$. Note

that $|\{C/Z(K) \mid C/Z(K) \triangleleft_p K/Z(K)\}| = \begin{bmatrix} 2 \\ 1 \end{bmatrix}_p = \frac{p^2 - 1}{p - 1} = p + 1$. This implies that K has $p + 1$ abelian maximal subgroups. The group G acts via conjugation on $p + 1$ maximal abelian subgroups of K . Orbits of this action have lengths $\equiv 0 \pmod{p}$. This implies that there is at least one fixed subgroup and that one is G -invariant. The proof is identical in the case when $K' = 1$. \square

THEOREM 3.3. *Let $N \trianglelefteq G$ and $|N| > p^3$, where G is a p -group. Then there is some abelian $D < N$ of order p^3 such that $D \trianglelefteq G$.*

PROOF. There is a composition series that goes through each normal subgroup of G . It implies that there is a G -invariant subgroup $M < N$ of order p^4 . Let $A < M$ be of order p^2 . Then, A is abelian and $|M : A| = p^2$. By Theorem 3.2, there is a $B \trianglelefteq M$ of order p^2 . Note that B is abelian as well. Since $|Aut(B)_p| = p$, it is necessary that $|N_M(B) : C_M(B)| \leq p$, where $N_M(B) = M$. If $M = C_M(B)$, then $B \leq Z(M)$. This implies that there is $g \in M - B$ such that $g^p \in B$ and $\langle B, g \rangle < M$ is an abelian group of order p^3 .

If $C_M(B) \triangleleft_p M$, then $C_M(B)$ is abelian of order p^3 . Thus, we can always find an abelian M -invariant subgroup of M the order of which is p^3 . The claim follows from Theorem 3.2. \square

PROPOSITION 3.4. *Let $G \in CZ_p$ be a 2-generated group of order p^6 . Let $\Phi(G) = N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$ and $exp(G) \leq p^2$. Then $\mathcal{U}_1(G) = \mathcal{U}_1(N) = \langle x^p, y^p \rangle$ and $|G'| \geq p^3$.*

PROOF. From Lemma 3.1 the center of N is $Z(N) = \langle x^p, y^p \rangle = \Phi(N)$. Therefore, $\langle x^p, y^p \rangle \leq \mathcal{U}_1(N) \leq \Phi(N) = \langle x^p, y^p \rangle$. This implies $\mathcal{U}_1(N) = \langle x^p, y^p \rangle$. Thus, $\langle x^p, y^p \rangle \leq \mathcal{U}_1(G)$. Since $exp(G) = p^2$, there is $x \in G$ and $o(x) = p^2$. Furthermore, $x^p \in \Phi(G) = N$ and $o(x^p) = p$. By Lemma 2.4, it follows that if $t \in N \setminus \Phi(N)$, then $o(t) = p^2$. Therefore, $x^p \in \Phi(N) = \langle x^p, y^p \rangle$. Thus, $\mathcal{U}_1(G) \leq \langle x^p, y^p \rangle$ and finally $\mathcal{U}_1(G) = \langle x^p, y^p \rangle$.

Since $\langle x^p \rangle = N' \leq G'$ and $\langle x^p \rangle \leq \mathcal{U}_1(G)$, we have $|\mathcal{U}_1(G) \cap G'| \geq p$ and $p \leq |\mathcal{U}_1(G) \cap G'| = \frac{|\mathcal{U}_1(G)||G'|}{|\mathcal{U}_1(G)G'|} = \frac{p^2|G'|}{|\Phi(G)|} = \frac{|G'|}{p^2}$, yielding $|G'| \geq p^3$. \square

THEOREM 3.5. *Let $G \in CZ_p$ be of order p^6 and $\Phi(G) = N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. Let $exp(G) = p^2$ and $|G'| = p^4$. Then $Z(G) = \langle x^p \rangle$ and $Z_2(G) = Z(N) = \langle x^p, y^p \rangle$.*

PROOF. By Proposition 3.4, $|G'| \geq p^3$. Since $|G : G'| \leq p^2$, the only options are $|G'| = p^3$ or $|G'| = p^4$. Let $|G'| = p^4$. Since $G' \leq \Phi(G) = \mathcal{U}_1(G)G'$, we have $G' = N$. By Gr\"unns's theorem (see [1]), we have $[G' : Z_2(G)] = 1$. Therefore, $[N, Z_2(G)] = 1$. Since $G \in CZ_p$, we have $Z_2(G) \leq C_G(N) = Z(N) = \langle x^p, y^p \rangle$ (see Lemma 3.1). This implies $Z_2(G)/Z_1(G) =$

$Z(G/Z_1(G)) > 1$. Therefore, $Z_2(G) > Z_1(G) > 1$. Since $|Z_2(G)| = |\langle x^p, y^p \rangle| = p^2$, we have $|Z_1(G)| = |Z(G)| = p$. Note that $N' = \langle x^p \rangle$ is a characteristic subgroup of N , and N is a characteristic subgroup of G . It follows that $N' \trianglelefteq G$ is of order p . Therefore, $|N' \cap Z(G)| > 1$ and $N' = Z(G) = \langle x^p \rangle$. \square

THEOREM 3.6. *Let $G \in CZ_p$ be a group of order p^6 . Let $\Phi(G) = N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$ and $\exp(G) \leq p^2$. If $|G'| = p^4$, then G is a group of maximal class.*

PROOF. If we assume that $cl(G) < max$, then, by Theorem 3.3, there is a G -invariant, abelian subgroup $A \leq N$ of order p^3 . This implies $Z(N) = \langle x^p, y^p \rangle \leq A$. Otherwise, N would be an abelian group. Note that $\mathcal{U}_1(N) = Z(N) \leq A$. Since $\mathcal{U}_1(N)$ is a characteristic subgroup of A , we have $\langle x^p, y^p \rangle \triangleleft G$. We know that $|G : \Phi(G)| = p^2$ and $G = \langle a, b \rangle$ for some $a, b \in G$. By Theorem 3.5, we have $Z(G) = \langle x^p \rangle$. This yields $[x^p, a] = [x^p, b] = 1$. If $(y^p)^a = (y^p)^b = y^p$, then $y^p \in Z(G) = \langle x^p \rangle$, which is a contradiction. Therefore, we have $(y^p)^a \neq y^p$. Since $\langle x^p, y^p \rangle \triangleleft G$, we have $(y^p)^a \in \langle x^p, y^p \rangle$. Also, $o(a^p) \leq p$ and $a^p \in N$. Therefore, $a^p \in \Omega_1(N) = \langle x^p, y^p \rangle$. It follows that $\langle x^p, y^p, a \rangle$ is a nonabelian group of order p^3 and by Theorem 1.3 we have $cl(G) = max$. This is the final contradiction which proves the theorem. \square

THEOREM 3.7. *Let $G \in CZ_p$, $cl(G) < max$, $\exp(G) \leq p^2$ and let $N \leq \Phi(G)$ be a nonabelian G -invariant subgroup of order p^4 . Then $|G| = p^7$.*

PROOF. By Corollary 2.8, we have $p^5 \leq |G| \leq p^7$. Since $|\Phi(G)| \geq p^4$, it follows $|G| \geq p^6$ (since otherwise $d(G) = 1$). By Lemma 2.3, we know the structure of the group N .

Let $|G| = p^6$. By Proposition 3.4, we have $|G'| \geq p^3$. Since $|G| = p^6$ and $d(G) = 2$, it follows that $G' \leq \Phi(G)$ and $|G'| \leq p^4$. If $|G'| = p^4$, then by Theorem 3.6, the class of G would be maximal, contradicting the assumption. Hence $|G'| = p^3$. By Proposition 3.4, we have $\mathcal{U}_1(G) = \mathcal{U}_1(N) = Z(N) = \langle x^p, y^p \rangle = \Phi(N)$. Since $|G'| = p^3$, we have $G' \leq N = \Phi(G)$. On the other hand, G' is a maximal subgroup of N . Therefore $Z(N) = \mathcal{U}_1(G) = \Phi(N) \leq G'$. This implies $\Phi(G) = \mathcal{U}_1(G)G' = G' < N = \Phi(G)$. This is a contradiction. So, the only remaining possibility is $|G| = p^7$. \square

4. THE CASE $|G| = p^7$ AND $cl(G) < max$

We shall continue with the same assumption that there is a nonabelian $N \leq \Phi(G)$ of order p^4 . Additionally, we shall assume that G is **not of maximal class** and $|G| = p^7$, $\exp(G) = p^2$. Note that $\exp(G) \leq p^3$. We begin with the following result on the size of G' .

LEMMA 4.1. *Let $G \in CZ_p$ be a group of order p^7 and $\exp(G) = p^2$ where $N = \Phi(G) = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. Then $\mathcal{U}_1(G) = \mathcal{U}_1(N) = \langle x^p, y^p \rangle$ and $|G'| \geq p^3$.*

PROOF. Notice that $\mathcal{U}_1(N) \leq \mathcal{U}_1(G)$ and $\exp(\mathcal{U}_1(G)) = p$. By Lemma 3.1, we have $\mathcal{U}_1(N) = \Phi(N) = \langle x^p, y^p \rangle$. We also have $\mathcal{U}_1(G) \leq \Phi(G) = N$. Then $\mathcal{U}_1(G) \leq \Phi(N) = \mathcal{U}_1(N)$. This implies $\mathcal{U}_1(G) = \mathcal{U}_1(N) = \langle x^p, y^p \rangle$.

By Lemma 3.4, we have $\langle x^p \rangle = N' \leq G' \cap \mathcal{U}_1(G)$. Therefore $p \leq |\mathcal{U}_1(G) \cap G'| = \frac{|\mathcal{U}_1(G)| \cdot |G'|}{|\mathcal{U}_1(G) \cdot G'|} = \frac{p^2 \cdot |G'|}{|\Phi(G)|} = \frac{|G'|}{p^2}$. This yields $|G'| \geq p^3$. \square

THEOREM 4.2. *Let $G \in CZ_p$ be a group of order p^7 with $\exp(G) = p^2$ where $N = \Phi(G) = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$. If $|G'| = p^4$, then $Z(G) = \langle x^p \rangle$ and $Z_2(G) = Z(N) = \langle x^p, y^p \rangle$.*

PROOF. By Lemma 4.1, we have $|G'| \geq p^3$. Since $G' \leq \Phi(G) = N$, it follows $|G'| \leq p^4$. The rest of the proof follows the proof of Theorem 3.5. \square

Now we shall present the main result.

THEOREM 4.3. *Let $G \in CZ_p$ be of exponent p^2 , where $cl(G) < \max$. Let $N \leq \Phi(G)$ be a G -invariant nonabelian subgroup of order p^4 . Then $|G| = p^7$ and $N = \langle x, y \mid x^{p^2} = y^{p^2} = 1, x^y = x^{1+p} \rangle$ is of index p in $\Phi(G)$.*

PROOF. Assume that $N = \Phi(G)$ and $|G'| = p^4$. As in Theorem 3.5, we have $Z(G) = \langle x^p \rangle$ and $Z_2(G) = Z(N) = \langle x^p, y^p \rangle$. By Theorem 3.3, there is an abelian group $A \trianglelefteq G$ such that $A \leq N$ and $|A| = p^3$. Therefore $Z(N) \leq A$, since otherwise $AZ(N) = N$ and N would be an abelian group. By Lemma 4.1, we have $\mathcal{U}_1(G) = \mathcal{U}_1(N) = \langle x^p, y^p \rangle \leq A$. Note that $\mathcal{U}_1(N) = Z(N)$. By Lemma 3.1, we have $Z(N) = \Phi(N) = \langle x^p, y^p \rangle \trianglelefteq G$ (since $\mathcal{U}_1(G) = \langle x^p, y^p \rangle$ is a characteristic subgroup of G). From $G/\Phi(G) \cong E_{p^3}$, we have $G = \langle a, b, c \rangle$ for some generators $a, b, c \in G$. Since $x^p \in Z(G)$, it follows $[x^p, a] = [x^p, b] = [x^p, c]$.

If $(y^p)^a = (y^p)^b = (y^p)^c = y^p$, then $y^p \in Z(G)$. This is a contradiction. Thus, we may assume $(y^p)^a \neq y^p$. Furthermore, $o(a^p) \leq p$ (since $\exp(G) = p^2$) and $a^p \mathcal{U}_1(G) = \langle x^p, y^p \rangle$. This implies that $\langle x^p, y^p, a \rangle$ is a nonabelian group of order p^3 and by Theorem 1.3 the group G has maximal class. This is a contradiction. Therefore, by Lemma 4.1, we have $|G'| = p^3$. By Lemma 4.1, it follows that $\mathcal{U}_1(G) = \mathcal{U}_1(N) = Z(N) = \Phi(N) = \langle x^p, y^p \rangle$. From $|G'| = p^3$, we have $G' \leq N = G' \Phi(G)$. Since G' is maximal in N , it implies $\Phi(N) < G'$. Since $\mathcal{U}_1(G) = \mathcal{U}_1(N) = \Phi(N) < G'$, it follows that $\Phi(G) = \mathcal{U}_1(G)G' \leq G' < N$. This yields now $\Phi(G) < N = \Phi(G)$, which is a contradiction. By Theorem 3.7, we have $|G| = p^7$. It follows $N < \Phi(G)$. The description of the group N is given by Lemma 2.3. Since $|G : \Phi(G)| \geq p^2$, we have $|\Phi(G) : N| = p$. \square

ACKNOWLEDGEMENTS.

This work has been fully supported by Croatian Science Foundation under the project 6732 and 9752.

REFERENCES

- [1] Y. Berkovich, Groups of Prime Power Order Vol. 1., Walter de Gruyter, Berlin - New York, 2008.
- [2] Y. Berkovich, Z. Janko, Groups of Prime Power Order Vol. 2., Walter de Gruyter, Berlin - New York, 2008.
- [3] Y. Berkovich, Z. Janko, Groups of Prime Power Order Vol. 3., Walter de Gruyter, Berlin - New York, 2010.
- [4] M- Hall, Theory of Groups, The Macmillan Company, New York, 1959.
- [5] M. O. Pavčević, K. Tabak, *CZ-groups*, Glas. Mat. Ser. III **51(71)** (2016), 345–358.

M.O. Pavčević
Department of applied mathematics
Faculty of Electrical Engineering and Computing
University of Zagreb
10 000 Zagreb
Croatia
E-mail: mario.pavcevic@fer.hr

K. Tabak
Rochester Institute of Technology
Zagreb Campus
D.T. Gavrana 15
10000 Zagreb
Croatia
E-mail: kxtcad@rit.edu