

# The Dixmier property and weak centrality for $C^*$ -algebras

Ilja Gogić

Department of Mathematics, Faculty of Science  
University of Zagreb

Seminar za algebro in funkcionalno analizo  
Fakulteta za matematiko in fiziko  
Ljubljana, January 8, 2026

*based on a joint work with Robert J. Archbold and Leonel Robert  
(IMRN 2022 & 2023)*

# $C^*$ -algebras - definition and basic properties

A  **$C^*$ -algebra** is a complex Banach  $*$ -algebra  $A$  whose norm  $\|\cdot\|$  satisfies the  $C^*$ -identity

$$\|a^*a\| = \|a\|^2, \quad \text{for all } a \in A.$$

## $C^*$ -algebras - definition and basic properties

A  **$C^*$ -algebra** is a complex Banach  $*$ -algebra  $A$  whose norm  $\|\cdot\|$  satisfies the  $C^*$ -identity

$$\|a^*a\| = \|a\|^2, \quad \text{for all } a \in A.$$

The  $C^*$ -identity is a very strong requirement. For instance, for any  $a \in A$  let  $\sigma(a)$  denote the spectrum of  $a$ , i.e.

$$\sigma(a) := \{\lambda \in \mathbb{C} : \lambda 1 - a \text{ is not invertible in } A\}.$$

Then the  $C^*$ -identity combined with the spectral radius formula

$$r(a) := \max\{|\lambda| : \lambda \in \sigma(a)\} = \lim_{n \rightarrow \infty} \|a^n\|^{\frac{1}{n}},$$

implies that the  $C^*$ -norm is uniquely determined by the algebraic structure:

$$\|a\|^2 = \|a^*a\| = r(a^*a) = \max\{|\lambda| : \lambda \in \sigma(a^*a)\}.$$

In the category of  $C^*$ -algebras, the natural candidates for morphisms are the  **$*$ -homomorphisms**, i.e. the algebra homomorphisms which preserve the involution. Basic properties:

- they are automatically contractive (isometric if injective), and
- their image is a  $C^*$ -subalgebra of the codomain  $C^*$ -algebra.

## Basic examples

- To any LCH (locally compact Hausdorff) space one can associate a commutative  $C^*$ -algebra  $C_0(X)$  of all continuous functions  $f : X \rightarrow \mathbb{C}$  that vanish at infinity, with respect to the pointwise operations, involution  $f^*(x) := \overline{f(x)}$ , and sup-norm  $\|f\|_\infty := \sup_{x \in X} |f(x)|$ .
- The set  $B(\mathcal{H})$  of bounded linear operators on a Hilbert space  $\mathcal{H}$  becomes a  $C^*$ -algebra with respect to the standard operations, usual adjoint and operator norm. In particular, the complex matrix algebras  $M_n(\mathbb{C})$  are  $C^*$ -algebras. In fact, the finite direct sums of matrix algebras over  $\mathbb{C}$  make up all finite-dimensional  $C^*$ -algebras.
- To any LC group  $G$ , one can associate a  $C^*$ -algebra  $C^*(G)$ . Everything about the representation theory of  $G$  is encoded in  $C^*(G)$ .
- The category of  $C^*$ -algebras is closed under the formation of direct products, direct sums, extensions, direct limits, pullbacks, pushouts, ( $C^*$ -)tensor products, etc.

In fact, all commutative  $C^*$ -algebras arise as in previous example:

### **Theorem (Commutative Gelfand-Naimark theorem, 1943)**

*The (contravariant) functor  $X \rightsquigarrow C_0(X)$  defines an equivalence of categories of LCH spaces (with proper continuous maps as morphisms) and commutative  $C^*$ -algebras (with non-degenerate  $*$ -homomorphisms as morphisms).*

In fact, all commutative  $C^*$ -algebras arise as in previous example:

### Theorem (Commutative Gelfand-Naimark theorem, 1943)

*The (contravariant) functor  $X \rightsquigarrow C_0(X)$  defines an equivalence of categories of LCH spaces (with proper continuous maps as morphisms) and commutative  $C^*$ -algebras (with non-degenerate  $*$ -homomorphisms as morphisms).*

In other words: By passing from the space  $X$  to the function algebra  $C_0(X)$ , no information is lost. In fact,  $X$  can be recovered from  $C_0(X)$ . Thus, topological properties of  $X$  can be translated into algebraic properties of  $C_0(X)$ , and vice versa. Therefore, the theory of  $C^*$ -algebras is often thought of as **noncommutative topology**.

## Representations of $C^*$ -algebras

A **representation** of a  $C^*$ -algebra  $A$  is a  $*$ -homomorphism  $\pi : A \rightarrow \mathbb{B}(\mathcal{H})$  for some Hilbert space  $\mathcal{H}$ . A representation  $\pi$  is said to be **irreducible** if it has no nontrivial (closed) invariant subspaces (i.e. if  $\mathcal{K}$  is a (closed) subspace of  $\mathcal{H}$  such that  $\pi(A)\mathcal{K} \subseteq \mathcal{K}$ , then  $\mathcal{K} = \{0\}$  or  $\mathcal{K} = \mathcal{H}$ ).

## Representations of $C^*$ -algebras

A **representation** of a  $C^*$ -algebra  $A$  is a  $*$ -homomorphism  $\pi : A \rightarrow \mathbb{B}(\mathcal{H})$  for some Hilbert space  $\mathcal{H}$ . A representation  $\pi$  is said to be **irreducible** if it has no nontrivial (closed) invariant subspaces (i.e. if  $\mathcal{K}$  is a (closed) subspace of  $\mathcal{H}$  such that  $\pi(A)\mathcal{K} \subseteq \mathcal{K}$ , then  $\mathcal{K} = \{0\}$  or  $\mathcal{K} = \mathcal{H}$ ).

## Theorem (General Gelfand-Naimark theorem, 1943)

*Any  $C^*$ -algebra admits an injective (hence isometric) representation on some Hilbert space.*

## Representations of $C^*$ -algebras

A **representation** of a  $C^*$ -algebra  $A$  is a  $*$ -homomorphism  $\pi : A \rightarrow \mathbb{B}(\mathcal{H})$  for some Hilbert space  $\mathcal{H}$ . A representation  $\pi$  is said to be **irreducible** if it has no nontrivial (closed) invariant subspaces (i.e. if  $\mathcal{K}$  is a (closed) subspace of  $\mathcal{H}$  such that  $\pi(A)\mathcal{K} \subseteq \mathcal{K}$ , then  $\mathcal{K} = \{0\}$  or  $\mathcal{K} = \mathcal{H}$ ).

## Theorem (General Gelfand-Naimark theorem, 1943)

*Any  $C^*$ -algebra admits an injective (hence isometric) representation on some Hilbert space.*

Because of the previous theorem,  $C^*$ -algebras can be concretely defined to be norm closed self-adjoint subalgebras of bounded operators on some Hilbert space  $\mathcal{H}$ .

## The primitive spectrum of a $C^*$ -algebra

Let  $A$  be  $C^*$ -algebra.

- A **primitive ideal** of  $A$  is an ideal which is the kernel of an irreducible representation of  $A$ .
- The **primitive spectrum** of  $A$  is the set  $\text{Prim}(A)$  of primitive ideals of  $A$  equipped with the **Jacobson (hull-kernel) topology**: if  $S$  is a set of primitive ideals, its closure is

$$\overline{S} := \left\{ P \in \text{Prim}(A) : \ker S = \bigcap_{Q \in S} Q \subseteq P \right\}.$$

## The primitive spectrum of a $C^*$ -algebra

Let  $A$  be  $C^*$ -algebra.

- A **primitive ideal** of  $A$  is an ideal which is the kernel of an irreducible representation of  $A$ .
- The **primitive spectrum** of  $A$  is the set  $\text{Prim}(A)$  of primitive ideals of  $A$  equipped with the **Jacobson (hull-kernel) topology**: if  $S$  is a set of primitive ideals, its closure is

$$\overline{S} := \left\{ P \in \text{Prim}(A) : \ker S = \bigcap_{Q \in S} Q \subseteq P \right\}.$$

### Example - commutative case

If  $A = C_0(X)$  and  $x \in X$ , let  $P_x := \{f \in C_0(X) : f(x) = 0\}$ . Then  $\text{Prim}(C_0(X)) = \{P_x : x \in X\}$ . Moreover, the correspondence  $x \mapsto P_x$  defines a homeomorphism between  $X$  and  $\text{Prim}(C_0(X))$ .

## Properties of $\text{Prim}(A)$

- $\text{Prim}(A)$  is always a locally compact and is compact if  $A$  is unital.
- If  $A$  is separable,  $\text{Prim}(A)$  is second countable.
- However, as a topological space,  $\text{Prim}(A)$  is in general badly behaved and may satisfy only the  $T_0$ -separation axiom.

## Properties of $\text{Prim}(A)$

- $\text{Prim}(A)$  is always a locally compact and is compact if  $A$  is unital.
- If  $A$  is separable,  $\text{Prim}(A)$  is second countable.
- However, as a topological space,  $\text{Prim}(A)$  is in general badly behaved and may satisfy only the  $T_0$ -separation axiom.

When a  $C^*$ -algebra  $A$  is unital, the Jacobson topology on  $\text{Prim}(A)$  not only describes the ideal structure of  $A$ , but also allows us to completely describe its centre  $Z(A) = \{z \in A : za = az\}$ :

### Dauns-Hofmann Theorem, 1968

Let  $A$  be a unital  $C^*$ -algebra. Then there is a  $*$ -isomorphism  $\Psi_A : Z(A) \rightarrow C(\text{Prim}(A))$  such that

$$z + P = \Psi_A(z)(P)1 + P$$

for all  $f \in C(\text{Prim}(A))$ ,  $a \in A$  and  $P \in \text{Prim}(A)$ .

# The Dixmier property and weak centrality

## Preliminaries

- Throughout  $A$  will be a  $C^*$ -algebra with centre  $Z(A)$  and unitary group  $\mathcal{U}(A) = \{u \in A : u^*u = uu^* = 1\}$  (if  $A$  is unital).
- By an ideal of  $A$  we always mean a closed two-sided ideal. We denote by  $\text{Ideal}(A)$  the set of all (closed two-sided) ideals of  $A$ .
- By  $\mathcal{S}(A)$  we denote the set of all states on  $A$  (i.e. positive linear functionals  $\omega : A \rightarrow \mathbb{C}$  of norm 1) equipped with the relative  $w^*$ -topology.
- A state  $\tau \in \mathcal{S}(A)$  is said to be **tracial** if  $\tau(xy) = \tau(yx) \ \forall x, y \in A$ .
- By  $\mathcal{T}(A)$  we denote the set of all tracial states on  $A$ . If  $A$  is unital then  $\mathcal{T}(A)$  is a convex  $w^*$ -compact subset of  $\mathcal{S}(A)$ .
- By  $\partial_e \mathcal{T}(A)$  we denote the extreme boundary of  $\mathcal{T}(A)$ , so that  $\mathcal{T}(A)$  is equal to the closed convex hull of  $\partial_e \mathcal{T}(A)$  (by the Krein-Milman theorem).

- A **unitary mixing operator** on  $A$  is a map  $\phi: A \rightarrow A$  of the form

$$\phi(x) = \sum_{i=1}^n t_i u_i^* x u_i,$$

where  $n$  is a positive integer,  $u_1, \dots, u_n \in \mathcal{U}(A)$  and  $t_1, \dots, t_n$  non-negative real numbers such that  $t_1 + \dots + t_n = 1$ . The set of all such maps is denoted by  $\text{UM}(A)$ .

## The Dixmier property and weak centrality

Let  $A$  be a unital  $C^*$ -algebra.

- For an element  $a \in A$  the **Dixmier set**  $D_A(a)$  is defined as the norm-closure of the set  $\{\phi(a) : \phi \in \text{UM}(A)\}$ . Then  $A$  is said to have the **Dixmier property** (DP) if

$$D_A(a) \cap Z(A) \neq \emptyset, \quad \forall a \in A.$$

- $A$  is said to be **weakly central** (WC) if for any pair of maximal ideals  $M_1$  and  $M_2$  of  $A$ ,  $M_1 \cap Z(A) = M_2 \cap Z(A)$  implies  $M_1 = M_2$ .

## Important properties

- $\text{DP} \implies \text{WC}$  (Archbold 1972).
- All von Neumann algebras satisfy DP (Dixmier 1949, Misonou 1952).
- A unital simple  $C^*$ -algebra satisfies DP iff it admits at most one tracial state (Haagerup-Zsidó 1984). In particular,  $\text{WC} \not\implies \text{DP}$ .

## Important properties

- $DP \implies WC$  (Archbold 1972).
- All von Neumann algebras satisfy DP (Dixmier 1949, Misonou 1952).
- A unital simple  $C^*$ -algebra satisfies DP iff it admits at most one tracial state (Haagerup-Zsidó 1984). In particular,  $WC \not\implies DP$ .

A complete characterization of  $C^*$ -algebras with DP was obtained recently.

### Theorem (Archbold-Robert-Tikuisis, 2017)

A unital  $C^*$ -algebra  $A$  has DP iff all of the following hold:

- $A$  is WC.
- Every simple quotient of  $A$  has at most one tracial state.
- Every extreme tracial state of  $A$  factors through some simple quotient.

## Important properties

- $DP \implies WC$  (Archbold 1972).
- All von Neumann algebras satisfy DP (Dixmier 1949, Misonou 1952).
- A unital simple  $C^*$ -algebra satisfies DP iff it admits at most one tracial state (Haagerup-Zsidó 1984). In particular,  $WC \not\implies DP$ .

A complete characterization of  $C^*$ -algebras with DP was obtained recently.

### Theorem (Archbold-Robert-Tikuisis, 2017)

A unital  $C^*$ -algebra  $A$  has DP iff all of the following hold:

- $A$  is WC.
- Every simple quotient of  $A$  has at most one tracial state.
- Every extreme tracial state of  $A$  factors through some simple quotient.

### Corollary

A unital postliminal  $C^*$ -algebra has DP iff it is WC.

## Corollary

For a unital  $C^*$ -algebra  $A$  the following conditions are equivalent:

- $Z(A) = \mathbb{C}1$  and  $A$  has DP.
- $A$  has a unique maximal ideal  $M$ ,  $A$  (or  $A/M$ ) has at most one tracial state and  $M$  has no tracial states.

## Corollary

For a unital  $C^*$ -algebra  $A$  the following conditions are equivalent:

- $Z(A) = \mathbb{C}1$  and  $A$  has DP.
- $A$  has a unique maximal ideal  $M$ ,  $A$  (or  $A/M$ ) has at most one tracial state and  $M$  has no tracial states.

## The Dixmier's example

Let  $\mathcal{H}$  be a separable infinite-dimensional Hilbert space and let  $p \in B(\mathcal{H})$  be any projection with infinite-dimensional kernel and image. Set

$$A := K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1-p) \subset B(\mathcal{H}).$$

Then  $Z(A) = \mathbb{C}1$ ,  $A$  has precisely two maximal ideals, namely

$$M_1 := K(\mathcal{H}) + \mathbb{C}p \quad \text{and} \quad M_2 := K(\mathcal{H}) + \mathbb{C}(1-p),$$

and obviously  $M_1 \cap Z(A) = M_2 \cap Z(A) = \{0\}$ . Hence,  $A$  is not WC.

In 2008 Magajna gave a characterisation of weak centrality in terms of more general averaging which are defined as follows.

In 2008 Magajna gave a characterisation of weak centrality in terms of more general averaging which are defined as follows.

## EUCP operators and the Magajna set

- By an **elementary unital completely positive operator** on a unital  $C^*$ -algebra  $A$  we mean a map  $\phi : A \rightarrow A$  of the form

$$\phi(x) = \sum_{i=1}^n a_i^* x a_i,$$

where  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in A$  such that  $\sum_{i=1}^n a_i^* a_i = 1$ . The set of all such maps on  $A$  is denoted by  $\text{EUCP}(A)$ .

- For  $a \in A$  we define the **Magajna set**  $M_A(a)$  as the norm-closure of the set  $\{\phi(a) : \phi \in \text{EUCP}(A)\}$  (i.e. the closed  $C^*$ -convex hull of  $a$ ). Obviously  $D_A(a) \subseteq M_A(a)$  for any  $a \in A$ .

In 2008 Magajna gave a characterisation of weak centrality in terms of more general averaging which are defined as follows.

## EUCP operators and the Magajna set

- By an **elementary unital completely positive operator** on a unital  $C^*$ -algebra  $A$  we mean a map  $\phi : A \rightarrow A$  of the form

$$\phi(x) = \sum_{i=1}^n a_i^* x a_i,$$

where  $n \in \mathbb{N}$ ,  $a_1, \dots, a_n \in A$  such that  $\sum_{i=1}^n a_i^* a_i = 1$ . The set of all such maps on  $A$  is denoted by  $\text{EUCP}(A)$ .

- For  $a \in A$  we define the **Magajna set**  $M_A(a)$  as the norm-closure of the set  $\{\phi(a) : \phi \in \text{EUCP}(A)\}$  (i.e. the closed  $C^*$ -convex hull of  $a$ ). Obviously  $D_A(a) \subseteq M_A(a)$  for any  $a \in A$ .

## Theorem (Magajna, 2008)

A unital  $C^*$ -algebra  $A$  is WC iff  $M_A(a) \cap Z(A) \neq \emptyset$  for all  $a \in A$ .

On the other hand, in 1971 Vesterstrøm gave a different characterization of weak centrality of  $A$  in terms of the quotient images of  $Z(A)$ .

On the other hand, in 1971 Vesterstrøm gave a different characterization of weak centrality of  $A$  in terms of the quotient images of  $Z(A)$ .

### The centre-quotient property

- If  $J \in \text{Ideal}(A)$  it is immediate that

$$(Z(A) + J)/J = q_J(Z(A)) \subseteq Z(A/J),$$

where  $q_J : A \rightarrow A/J$  is the canonical map.

On the other hand, in 1971 Vesterstrøm gave a different characterization of weak centrality of  $A$  in terms of the quotient images of  $Z(A)$ .

### The centre-quotient property

- If  $J \in \text{Ideal}(A)$  it is immediate that

$$(Z(A) + J)/J = q_J(Z(A)) \subseteq Z(A/J),$$

where  $q_J : A \rightarrow A/J$  is the canonical map.

- $A$  is said to have the **centre-quotient property** (CQP) if the equality holds above for any  $J \in \text{Ideal}(A)$ .

On the other hand, in 1971 Vesterstrøm gave a different characterization of weak centrality of  $A$  in terms of the quotient images of  $Z(A)$ .

### The centre-quotient property

- If  $J \in \text{Ideal}(A)$  it is immediate that

$$(Z(A) + J)/J = q_J(Z(A)) \subseteq Z(A/J),$$

where  $q_J : A \rightarrow A/J$  is the canonical map.

- $A$  is said to have the **centre-quotient property** (CQP) if the equality holds above for any  $J \in \text{Ideal}(A)$ .

### Theorem (Vesterstrøm, Math. Scand. 1971)

A unital  $C^*$ -algebra is WC iff it has CQP.

On the other hand, in 1971 Vesterstrøm gave a different characterization of weak centrality of  $A$  in terms of the quotient images of  $Z(A)$ .

### The centre-quotient property

- If  $J \in \text{Ideal}(A)$  it is immediate that

$$(Z(A) + J)/J = q_J(Z(A)) \subseteq Z(A/J),$$

where  $q_J : A \rightarrow A/J$  is the canonical map.

- $A$  is said to have the **centre-quotient property** (CQP) if the equality holds above for any  $J \in \text{Ideal}(A)$ .

### Theorem (Vesterstrøm, Math. Scand. 1971)

A unital  $C^*$ -algebra is WC iff it has CQP.

#### Example

If  $A = K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1 - p) \subset B(\mathcal{H})$  is the Dixmier  $C^*$ -algebra, then  $Z(A) = \mathbb{C}1$ , while  $Z(A/K(\mathcal{H})) = A/K(\mathcal{H}) \cong \mathbb{C} \oplus \mathbb{C}$ .

We would like to investigate in what extent an arbitrary  $C^*$ -algebra fails to satisfy WC/CQP and DP. We proceed in two different ways:

We would like to investigate in what extent an arbitrary  $C^*$ -algebra fails to satisfy WC/CQP and DP. We proceed in two different ways:

## Global approach

Show that any  $C^*$ -algebra  $A$  has the largest WC ideal  $J_{wc}(A)$  and the largest ideal  $J_{dp}(A)$  with DP, and obtain their concrete descriptions.

We would like to investigate in what extent an arbitrary  $C^*$ -algebra fails to satisfy WC/CQP and DP. We proceed in two different ways:

## Global approach

Show that any  $C^*$ -algebra  $A$  has the largest WC ideal  $J_{wc}(A)$  and the largest ideal  $J_{dp}(A)$  with DP, and obtain their concrete descriptions.

## Local approach

Consider individual elements of  $A$  which witness DP and WC/CQP. We define an element  $a \in A$  to be:

- a **Dixmier element** if  $D_A(a) \cap Z(A) \neq \emptyset$ ;
- a **Magajna element** if  $M_A(a) \cap Z(A) \neq \emptyset$ ;
- a **CQ-element** if for any ideal  $J$  of  $A$ ,  $a + J \in Z(A/J)$  implies  $a \in Z(A) + J$ .

We would like to investigate in what extent an arbitrary  $C^*$ -algebra fails to satisfy WC/CQP and DP. We proceed in two different ways:

## Global approach

Show that any  $C^*$ -algebra  $A$  has the largest WC ideal  $J_{wc}(A)$  and the largest ideal  $J_{dp}(A)$  with DP, and obtain their concrete descriptions.

## Local approach

Consider individual elements of  $A$  which witness DP and WC/CQP. We define an element  $a \in A$  to be:

- a **Dixmier element** if  $D_A(a) \cap Z(A) \neq \emptyset$ ;
- a **Magajna element** if  $M_A(a) \cap Z(A) \neq \emptyset$ ;
- a **CQ-element** if for any ideal  $J$  of  $A$ ,  $a + J \in Z(A/J)$  implies  $a \in Z(A) + J$ .

By  $\text{Dix}(A)$ ,  $\text{Mag}(A)$  and  $\text{CQ}(A)$  we respectively denote the sets of all Dixmier, Magajna and CQ-elements of  $A$ . Obviously  $A$  has DP iff  $\text{Dix}(A) = A$ , while  $A$  is WC/has CQP iff  $\text{Mag}(A) = \text{CQ}(A) = A$ .

## Global approach

We begin by extending the definition of WC and DP for non-unital  $C^*$ -algebras in the obvious way: We say that a non-unital  $C^*$ -algebra  $A$  is WC/has DP if its minimal unitization  $A^\sharp = A \oplus \mathbb{C}1$  has the same property.

# Global approach

We begin by extending the definition of WC and DP for non-unital  $C^*$ -algebras in the obvious way: We say that a non-unital  $C^*$ -algebra  $A$  is WC/has DP if its minimal unitization  $A^\sharp = A \oplus \mathbb{C}1$  has the same property.

## Modular maximal ideals

- An ideal  $J$  of  $A$  is said to be **modular** if the algebra  $A/J$  is unital.
- Any proper modular ideal of  $A$  (if such exists) is contained in some modular maximal ideal of  $A$  and all modular maximal ideals of  $A$  are primitive. By  $\text{Max}(A)$  we denote the set of all modular maximal ideals of  $A$ , so that  $\text{Max}(A) \subseteq \text{Prim}(A)$ .
- $\text{Max}(A)$  can be empty (e.g. the algebra  $A = K(\mathcal{H})$  of compact operators on a separable infinite-dimensional Hilbert space  $\mathcal{H}$ ).
- If  $A$  is unital, both spaces  $\text{Prim}(A)$  and  $\text{Max}(A)$  are compact.
- For any  $J \in \text{Ideal}(A)$  we define  $\text{Max}^J(A)$  for the set of all modular maximal ideals of  $A$  that contain  $J$ . The space  $\text{Max}^J(A)$  is canonically homeomorphic to  $\text{Max}(A/J)$  via the assignment  $M \mapsto M/J$ .

## Theorem (Archbold-G, 2022)

For any  $C^*$ -algebra  $A$  the following conditions are equivalent:

- $A$  is WC.
- No modular maximal ideal of  $A$  contains  $Z(A)$  and for all  $M_1, M_2 \in \text{Max}(A)$ ,  $M_1 \cap Z(A) = M_2 \cap Z(A)$  implies  $M_1 = M_2$ .
- $A$  has CQP.

Further, the class of WC  $C^*$ -algebras is closed under forming ideals, quotients, direct sums and  $C^*$ -tensor product. Moreover if  $A_1$  and  $A_2$  are  $C^*$ -algebras then  $A_1 \otimes_{\beta} A_2$  is WC for some/every  $C^*$ -norm  $\beta$  iff both  $A_1$  and  $A_2$  are WC.

## Theorem (Archbold-G, 2022)

For any  $C^*$ -algebra  $A$  the following conditions are equivalent:

- $A$  is WC.
- No modular maximal ideal of  $A$  contains  $Z(A)$  and for all  $M_1, M_2 \in \text{Max}(A)$ ,  $M_1 \cap Z(A) = M_2 \cap Z(A)$  implies  $M_1 = M_2$ .
- $A$  has CQP.

Further, the class of WC  $C^*$ -algebras is closed under forming ideals, quotients, direct sums and  $C^*$ -tensor product. Moreover if  $A_1$  and  $A_2$  are  $C^*$ -algebras then  $A_1 \otimes_{\beta} A_2$  is WC for some/every  $C^*$ -norm  $\beta$  iff both  $A_1$  and  $A_2$  are WC.

It is possible to show that every  $C^*$ -algebra contains a largest ideal with CQP by using Zorn's lemma and the fact that the sum of two ideals with CQP has CQP.

## Theorem (Archbold-G, 2022)

For any  $C^*$ -algebra  $A$  the following conditions are equivalent:

- $A$  is WC.
- No modular maximal ideal of  $A$  contains  $Z(A)$  and for all  $M_1, M_2 \in \text{Max}(A)$ ,  $M_1 \cap Z(A) = M_2 \cap Z(A)$  implies  $M_1 = M_2$ .
- $A$  has CQP.

Further, the class of WC  $C^*$ -algebras is closed under forming ideals, quotients, direct sums and  $C^*$ -tensor product. Moreover if  $A_1$  and  $A_2$  are  $C^*$ -algebras then  $A_1 \otimes_{\beta} A_2$  is WC for some/every  $C^*$ -norm  $\beta$  iff both  $A_1$  and  $A_2$  are WC.

It is possible to show that every  $C^*$ -algebra contains a largest ideal with CQP by using Zorn's lemma and the fact that the sum of two ideals with CQP has CQP.

However, we take a different approach that has the merit of obtaining a formula for this ideal in terms of the set of those modular maximal ideals of  $A$  which witness the failure of the weak centrality of  $A$ .

## Theorem (Archbold-G, 2022)

Let  $A$  be a  $C^*$ -algebra and  $T_A$  the set of all  $M \in \text{Max}(A)$  such that either

- $Z(A) \subseteq M$ , or
- there is  $N \in \text{Max}(A)$  such that  $M \neq N$ ,  $Z(A) \not\subseteq M, N$  and  $M \cap Z(A) = N \cap Z(A)$ .

Then  $J_{wc}(A) := \ker T_A$  is the largest weakly central ideal of  $A$ .

## Theorem (Archbold-G, 2022)

Let  $A$  be a  $C^*$ -algebra and  $T_A$  the set of all  $M \in \text{Max}(A)$  such that either

- $Z(A) \subseteq M$ , or
- there is  $N \in \text{Max}(A)$  such that  $M \neq N$ ,  $Z(A) \not\subseteq M, N$  and  $M \cap Z(A) = N \cap Z(A)$ .

Then  $J_{wc}(A) := \ker T_A$  is the largest weakly central ideal of  $A$ .

### Example

- If  $A = K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1-p) \subset B(\mathcal{H})$  is the Dixmier's example, then  $J_{wc}(A) = (K(\mathcal{H}) + \mathbb{C}p) \cap (K(\mathcal{H}) + \mathbb{C}(1-p)) = K(\mathcal{H})$ .
- If  $G$  is either the free group on two generators  $\mathbb{F}_2$  or the discrete three-dimensional Heisenberg group

$$H_3 = \left\{ \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in \mathbb{Z} \right\},$$

then for  $A = C^*(G)$  we have  $J_{wc}(A) = \{0\}$ .

By using Zorn's lemma, in 1972 Archbold showed that any unital  $C^*$ -algebra  $A$  contains the largest ideal  $J_{dp}(A)$  with DP. We now describe  $J_{dp}(A)$  more explicitly. But first we recall the notion of Glimm ideals.

By using Zorn's lemma, in 1972 Archbold showed that any unital  $C^*$ -algebra  $A$  contains the largest ideal  $J_{dp}(A)$  with DP. We now describe  $J_{dp}(A)$  more explicitly. But first we recall the notion of Glimm ideals.

### Glimm ideals in unital $C^*$ -algebras

The **Glimm ideals** of a unital  $C^*$ -algebra  $A$  are the ideals of  $A$  generated by the maximal ideals of  $Z(A)$ . Notation:  $\text{Glimm}(A)$ .

By using Zorn's lemma, in 1972 Archbold showed that any unital  $C^*$ -algebra  $A$  contains the largest ideal  $J_{dp}(A)$  with DP. We now describe  $J_{dp}(A)$  more explicitly. But first we recall the notion of Glimm ideals.

### Glimm ideals in unital $C^*$ -algebras

The **Glimm ideals** of a unital  $C^*$ -algebra  $A$  are the ideals of  $A$  generated by the maximal ideals of  $Z(A)$ . Notation:  $\text{Glimm}(A)$ .

The correspondence

$$\text{Glimm}(A) \ni N \longmapsto N \cap Z(A) \in \text{Max}(Z(A))$$

is a bijection. Its inverse is given by

$$\text{Max}(Z(A)) \ni J \longmapsto JA \in \text{Glimm}(A),$$

and no closure operation is required, by the Hewitt–Cohen factorization theorem. Via this identification, we equip  $\text{Glimm}(A)$  with the compact Hausdorff topology inherited from  $\text{Max}(Z(A))$ .

- Now consider the set  $X \subseteq \text{Glimm}(A)$  of Glimm ideals  $N$  such that  $A/N$  has DP and a trivial centre.
- This is equivalent to saying that  $N$  is contained in a unique maximal ideal  $M_N$  of  $A$ , that  $A/N$  has at most one tracial state and that if  $A/N$  does have a tracial state then it factors through  $A/M_N$ .
- For  $N \in \text{Glimm}(A) \setminus X$  define

$$I_N := \ker \text{Max}^N(A) \cap \ker\{I_\tau : \tau \in \mathcal{T}(A/N)\}$$

where, for  $\tau \in \mathcal{T}(A/N)$ ,  $I_\tau$  is the corresponding trace-kernel, i. e.

$$I_\tau := \{a \in A : \tau(a^*a + N) = 0\}.$$

- Now consider the set  $X \subseteq \text{Glimm}(A)$  of Glimm ideals  $N$  such that  $A/N$  has DP and a trivial centre.
- This is equivalent to saying that  $N$  is contained in a unique maximal ideal  $M_N$  of  $A$ , that  $A/N$  has at most one tracial state and that if  $A/N$  does have a tracial state then it factors through  $A/M_N$ .
- For  $N \in \text{Glimm}(A) \setminus X$  define

$$I_N := \ker \text{Max}^N(A) \cap \ker\{I_\tau : \tau \in \mathcal{T}(A/N)\}$$

where, for  $\tau \in \mathcal{T}(A/N)$ ,  $I_\tau$  is the corresponding trace-kernel, i. e.

$$I_\tau := \{a \in A : \tau(a^*a + N) = 0\}.$$

### Theorem (Archbold-G-Robert, 2023)

We have

$$J_{dp}(A) = \ker\{I_N : N \in \text{Glimm}(A) \setminus X\}.$$

## Local approach

Recall, if  $a \in A$  then:

- $a \in \text{Dix}(A)$  if  $D_A(A) \cap Z(A) \neq \emptyset$
- $a \in \text{Mag}(A)$  if  $M_A(A) \cap Z(A) \neq \emptyset$ .
- $a \in \text{CQ}(A)$  if for any ideal  $J$  of  $A$ ,  $a + J \in Z(A/J)$  implies  $a \in Z(A) + J$ .

## Local approach

Recall, if  $a \in A$  then:

- $a \in \text{Dix}(A)$  if  $D_A(A) \cap Z(A) \neq \emptyset$
- $a \in \text{Mag}(A)$  if  $M_A(A) \cap Z(A) \neq \emptyset$ .
- $a \in \text{CQ}(A)$  if for any ideal  $J$  of  $A$ ,  $a + J \in Z(A/J)$  implies  $a \in Z(A) + J$ .

We always have

$$\text{Dix}(A) \subseteq \text{Mag}(A) \subseteq \text{CQ}(A).$$

- $\text{Dix}(A)$  always contains  $Z(A) + J_{dp}(A)$ , all self-commutators  $[a^*, a]$  and all quasinilpotents. In particular,  $\text{Dix}(A) = Z(A)$  iff  $A$  is abelian.
- $\text{Mag}(A)$  always contains  $Z(A) + J_{wc}(A)$  and all products  $ab$  where  $a$  or  $b$  is quasinilpotent.
- $\text{CQ}(A)$  always contains all commutators  $[a, b]$ . There are  $C^*$ -algebras  $A$  such that  $[a, b] \notin \text{Mag}(A)$  for some  $a, b \in A$ .

We always have

$$\overline{\text{span}(\text{Mag}(A))} = \overline{\text{span}(\text{CQ}(A))} = Z(A) + \text{Ideal}([A, A]).$$

On the other hand, when sets  $\text{CQ}(A)$  and  $\text{Mag}(A)$  coincide,  $A$  is not far from being WC. On the other hand, when this fails, both sets dramatically fail to be  $C^*$ -subalgebras of  $A$ :

### **Theorem (Archbold-G, 2022 & Archbold-G-Robert, 2023)**

*The following conditions are equivalent:*

- $\text{Mag}(A) = \text{CQ}(A)$ .
- $\text{Mag}(A) = \text{CQ}(A) = Z(A) + J_{wc}(A)$ .
- $A/J_{wc}(A)$  is abelian.
- $\text{Mag}(A)$  and/or  $\text{CQ}(A)$  is closed under addition.
- $\text{Mag}(A)$  and/or  $\text{CQ}(A)$  is closed under multiplication.
- $\text{Mag}(A)$  is closed under EUCP operators.
- $\text{CQ}(A)$  is norm-closed.

We also exhibited examples of (separable continuous trace)  $C^*$ -algebras  $A$  for which  $J_{wc}(A) = \{0\}$ , while  $CQ(A)$  is norm-dense in  $A$ .

We also exhibited examples of (separable continuous trace)  $C^*$ -algebras  $A$  for which  $J_{wc}(A) = \{0\}$ , while  $CQ(A)$  is norm-dense in  $A$ .

In order to identify the set  $CQ(A)$  we shall need the following result:

### Theorem (Archbold-G, 2022)

Let  $A$  be a unital  $C^*$ -algebra and let  $J$  be an ideal of  $A$ . A central element  $\dot{z}$  of  $A/J$  can be lifted to a central element of  $A$  iff

$$\Psi_{A/J}(\dot{z})(P_1/J) = \Psi_{A/J}(\dot{z})(P_2/J)$$

for all  $P_1, P_2 \in \text{Prim}(A)$  that contain  $J$  and  $P_1 \cap Z(A) = P_2 \cap Z(A)$ .

We also exhibited examples of (separable continuous trace)  $C^*$ -algebras  $A$  for which  $J_{wc}(A) = \{0\}$ , while  $CQ(A)$  is norm-dense in  $A$ .

In order to identify the set  $CQ(A)$  we shall need the following result:

### Theorem (Archbold-G, 2022)

Let  $A$  be a unital  $C^*$ -algebra and let  $J$  be an ideal of  $A$ . A central element  $\dot{z}$  of  $A/J$  can be lifted to a central element of  $A$  iff

$$\Psi_{A/J}(\dot{z})(P_1/J) = \Psi_{A/J}(\dot{z})(P_2/J)$$

for all  $P_1, P_2 \in \text{Prim}(A)$  that contain  $J$  and  $P_1 \cap Z(A) = P_2 \cap Z(A)$ .

### Theorem (Archbold-G, 2022)

An element  $a \in A$  belongs to  $A \setminus CQ(A)$  iff one of the following holds:

- there exists  $M \in \text{Max}(A)$  such that  $Z(A) \subseteq M$  and  $a + M$  is a non-zero scalar in  $A/M$ ;
- there exist  $M_1, M_2 \in \text{Max}(A)$  and scalars  $\lambda_1 \neq \lambda_2$  such that  $Z(A) \neq M_1 \cap Z(A) = M_2 \cap Z(A)$  and  $a + M_i = \lambda_i 1_{A/M_i}$  ( $i = 1, 2$ ).

On the other hand, a complete description of  $\text{Dix}(A)$  and  $\text{Mag}(A)$  is in general difficult to obtain. This has led us to also consider the sets

$$\overline{\text{Mag}}(A) = \{a \in A : \text{dist}(M_A(a), Z(A)) = 0\},$$

$$\overline{\text{Dix}}(A) = \{a \in A : \text{dist}(D_A(a), Z(A)) = 0\}.$$

These are more tractable sets (e.g. they are norm-closed). We have

$$\begin{array}{ccc} \overline{\text{Dix}}(A) & & \\ \cup\!\!\!/\!\!\! \cup & & \cup\!\!\!/\!\!\! \cup \\ \text{Dix}(A) & & \overline{\text{Mag}}(A) \subseteq \text{CQ}(A). \\ \cup\!\!\!/\!\!\! \cup & & \cup\!\!\!/\!\!\! \cup \\ & & \text{Mag}(A) \end{array}$$

Also note that  $A$  has DP iff  $\overline{\text{Dix}}(A) = A$  and  $A$  is WC iff  $\overline{\text{Mag}}(A) = A$ .

## Numerical range

Given  $a \in A$  the **(algebraic) numerical range** of  $a$  is defined as  $W_A(a) := \{\omega(a) : \omega \in \mathcal{S}(A)\}$ . It is a compact convex subset of  $\mathbb{C}$  that contains  $\sigma(a)$ . If  $a$  is normal then  $W_A(a)$  is the convex hull of  $\sigma(a)$ .

## Numerical range

Given  $a \in A$  the **(algebraic) numerical range** of  $a$  is defined as  $W_A(a) := \{\omega(a) : \omega \in \mathcal{S}(A)\}$ . It is a compact convex subset of  $\mathbb{C}$  that contains  $\sigma(a)$ . If  $a$  is normal then  $W_A(a)$  is the convex hull of  $\sigma(a)$ .

### Theorem (Magajna, 2000)

Let  $a \in A$ . A normal element  $b \in A$  belongs to  $M_A(a)$  iff  $W_{A/P}(b + P) \subseteq W_{A/P}(a + P)$  for each  $P \in \text{Prim}(A)$ .

## Numerical range

Given  $a \in A$  the **(algebraic) numerical range** of  $a$  is defined as  $W_A(a) := \{\omega(a) : \omega \in \mathcal{S}(A)\}$ . It is a compact convex subset of  $\mathbb{C}$  that contains  $\sigma(a)$ . If  $a$  is normal then  $W_A(a)$  is the convex hull of  $\sigma(a)$ .

## Theorem (Magajna, 2000)

Let  $a \in A$ . A normal element  $b \in A$  belongs to  $M_A(a)$  iff  $W_{A/P}(b + P) \subseteq W_{A/P}(a + P)$  for each  $P \in \text{Prim}(A)$ .

## Theorem (Archbold-G-Robert, 2023)

For any  $a \in A$  we have  $a \in \overline{\text{Mag}}(A)$  iff for all  $N \in \text{Glimm}(A)$ ,

$$\Lambda_a(N) := \bigcap_{M \in \text{Max}^N(A)} W_{A/M}(a + M) \neq \emptyset.$$

Further, if  $a$  is self-adjoint, then  $a \in \overline{\text{Mag}}(A)$  iff  $a \in \text{Mag}(A)$ .

## Example

Let  $B = K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1 - p)$  be the Dixmier  $C^*$ -algebra and let  $A = C([-1, 1], M_2(\mathbb{C})) \otimes B$ . We define elements  $a, b \in A$  as:

$$a(t) := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad b(t) := \begin{pmatrix} \alpha(t) & 0 \\ 0 & \beta(t) \end{pmatrix},$$

where  $\alpha(t)$  and  $\beta(t)$  are curves in the plane such that:

- From  $t = -1$  to  $t = 0$  the interval  $[\alpha(t), \beta(t)]$  starts at  $[-1, -1 + 2i]$ , remains pinned at  $-1$  while rotating till it is flat and equal to  $[-1, 1]$  at  $t = 0$ .
- Then from  $t = 0$  to  $t = 1$  the interval  $[\alpha(t), \beta(t)]$  is pinned at  $1$ , and rotates till it stops at  $[1, 1 + 2i]$ .

If  $c \in A$  defined as

$$c := a \otimes p + b \otimes (1 - p),$$

then  $c$  is a normal element of  $A$  such that  $c \in \overline{\text{Mag}}(A) \setminus \text{Mag}(A)$ .

We now describe the set  $\overline{\text{Dix}}(A)$ . Define

$$Y := \{N \in \text{Glimm}(A) : \mathcal{T}(A/N) \neq \emptyset\}.$$

It is not difficult to see that  $Y$  is a closed subset of  $\text{Glimm}(A)$ .

We now describe the set  $\overline{\text{Dix}}(A)$ . Define

$$Y := \{N \in \text{Glimm}(A) : \mathcal{T}(A/N) \neq \emptyset\}.$$

It is not difficult to see that  $Y$  is a closed subset of  $\text{Glimm}(A)$ .

### Theorem (Archbold-G-Robert, 2023)

For an element  $a \in A$  consider the following conditions:

- (i)  $a \in \text{Dix}(A)$ .
- (ii)  $a \in \overline{\text{Dix}}(A)$ .
- (iii) (a) There is a function  $f_a: Y \rightarrow \mathbb{C}$  such that
  - (a1) for all  $N \in Y$  and  $\tau \in \mathcal{T}(A/N)$ ,  $f_a(N) = \tau(a + N)$ ,
  - (a2) for all  $N \in Y$ ,  $f_a(N) \in \Lambda_a(N)$ .
- (b) For all  $N \in \text{Glimm}(A) \setminus Y$ ,  $\Lambda_a(N) \neq \emptyset$ .

Then (i)  $\implies$  (ii)  $\iff$  (iii). Further, if (iii) holds then  $f_a$  is unique and it is continuous on  $Y$ . Finally, if  $a = a^*$ , then (i), (ii), and (iii) are equivalent.

## Example

Let  $B = K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1 - p)$  be the Dixmier  $C^*$ -algebra and  $A := C([-1, 1], \mathcal{O}_2) \otimes B$  ( $\mathcal{O}_2$  is the Cuntz algebra). Then  $\mathcal{T}(A) = \emptyset$ , so that  $\text{Dix}(A) = \text{Mag}(A)$  and there is  $a \in \overline{\text{Dix}(A)} \setminus \text{Dix}(A)$ .

## Example

Let  $B = K(\mathcal{H}) + \mathbb{C}p + \mathbb{C}(1 - p)$  be the Dixmier  $C^*$ -algebra and  $A := C([-1, 1], \mathcal{O}_2) \otimes B$  ( $\mathcal{O}_2$  is the Cuntz algebra). Then  $\mathcal{T}(A) = \emptyset$ , so that  $\text{Dix}(A) = \text{Mag}(A)$  and there is  $a \in \overline{\text{Dix}(A)} \setminus \text{Dix}(A)$ .

## Theorem (Archbold-G-Robert, 2023)

The set  $Z(A) + \overline{[A, A]}$  contains  $\overline{\text{Dix}(A)}$  and is equal to the closed linear span of  $\text{Dix}(A)$ . Further, the following conditions are equivalent:

- (i)  $\text{Dix}(A) = Z(A) + \overline{[A, A]}$ .
- (ii)  $\text{Dix}(A)$  is closed under unitary mixing operators.
- (iii)  $\text{Dix}(A)$  is closed under addition.
- (iv) (a) For all  $N \in Y$  and  $M \in \text{Max}^N(A)$ ,  $\mathcal{T}(A/M) \neq \emptyset$ .  
(b) For all  $N \in \text{Glimm}(A) \setminus Y$ ,  $\text{Max}^N(A)$  is a singleton set.

Moreover, when these equivalent conditions hold,  $\text{Dix}(A) = \overline{\text{Dix}(A)}$ .

## Problem

Is  $\overline{\text{Mag}(A)} = \overline{\text{Mag}}(A)$  and  $\overline{\text{Dix}(A)} = \overline{\text{Dix}}(A)$ ?

## Problem

Is  $\overline{\text{Mag}(A)} = \overline{\text{Mag}}(A)$  and  $\overline{\text{Dix}(A)} = \overline{\text{Dix}}(A)$ ?

## Theorem (Archbold-G-Robert, 2023)

The following conditions are equivalent:

- (i)  $A/J_{dp}(A)$  is abelian.
- (ii)  $\text{Dix}(A) = Z(A) + J_{dp}(A)$ ,
- (iii)  $\text{Dix}(A)$  is closed under multiplication.

Moreover, under these equivalent conditions  $J_{dp}(A) = J_{wc}(A)$ , so that

$$\text{Dix}(A) = \text{Mag}(A) = Z(A) + J_{dp}(A) = Z(A) + \overline{[A, A]}.$$

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

An algebra  $A$  over a field  $\mathbb{F}$  is said to be **centrally stable (CS)** if for any ideal  $J$  of  $A$  we have  $(Z(A) + J)/I = Z(A/J)$  (Brešar-G., 2019).

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

An algebra  $A$  over a field  $\mathbb{F}$  is said to be **centrally stable (CS)** if for any ideal  $J$  of  $A$  we have  $(Z(A) + J)/I = Z(A/J)$  (Brešar-G., 2019).

In contrast with the CQ-property for  $C^*$ -algebras, central stability is, as expected, a more delicate notion. For example:

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

An algebra  $A$  over a field  $\mathbb{F}$  is said to be **centrally stable (CS)** if for any ideal  $J$  of  $A$  we have  $(Z(A) + J)/I = Z(A/J)$  (Brešar-G., 2019).

In contrast with the CQ-property for  $C^*$ -algebras, central stability is, as expected, a more delicate notion. For example:

- For a unital noncommutative algebra  $A$ , it may occur that the centrally stable elements of  $A$  are precisely the central ones. This cannot happen for CQ-elements in the  $C^*$ -algebra setting.

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

An algebra  $A$  over a field  $\mathbb{F}$  is said to be **centrally stable (CS)** if for any ideal  $J$  of  $A$  we have  $(Z(A) + J)/I = Z(A/J)$  (Brešar-G., 2019).

In contrast with the CQ-property for  $C^*$ -algebras, central stability is, as expected, a more delicate notion. For example:

- For a unital noncommutative algebra  $A$ , it may occur that the centrally stable elements of  $A$  are precisely the central ones. This cannot happen for CQ-elements in the  $C^*$ -algebra setting.
- Central stability does not, in general, pass to ideals.

## Purely algebraic context – Central Stability

An analogous definition of the CQ-property makes sense for purely algebraic objects like groups, rings or algebras.

An algebra  $A$  over a field  $\mathbb{F}$  is said to be **centrally stable (CS)** if for any ideal  $J$  of  $A$  we have  $(Z(A) + J)/I = Z(A/J)$  (Brešar-G., 2019).

In contrast with the CQ-property for  $C^*$ -algebras, central stability is, as expected, a more delicate notion. For example:

- For a unital noncommutative algebra  $A$ , it may occur that the centrally stable elements of  $A$  are precisely the central ones. This cannot happen for CQ-elements in the  $C^*$ -algebra setting.
- Central stability does not, in general, pass to ideals.

### Theorem (Brešar-G., 2019)

If  $A$  is a finite-dimensional unital algebra over a perfect field  $\mathbb{F}$ , then  $A$  is CS iff there exist finite field extensions  $\mathbb{F}_1, \dots, \mathbb{F}_r$  of  $\mathbb{F}$ , commutative unital  $\mathbb{F}_i$ -algebras  $C_1, \dots, C_r$ , and central simple  $\mathbb{F}_i$ -algebras  $A_1, \dots, A_r$  such that  $A \cong (C_1 \otimes_{\mathbb{F}_1} A_1) \times \dots \times (C_r \otimes_{\mathbb{F}_r} A_r)$ .